

Title	A Fuzzy Three Sphere and Fuzzy Tori
Creators	Dolan, Brian P. and O'Connor, Denjoe
Date	2003
Citation	Dolan, Brian P. and O'Connor, Denjoe (2003) A Fuzzy Three Sphere and Fuzzy Tori. (Preprint)
URL	https://dair.dias.ie/id/eprint/620/
DOI	DIAS-STP-03-05

A Fuzzy Three Sphere and Fuzzy Tori

Brian P. Dolan^{a),b)*} and Denjoe O'Connor^{b)†}

^{a)}Dept. of Mathematical Physics, NUI, Maynooth, Ireland

^{b)}School of Theoretical Physics,
Dublin Institute for Advanced Studies,
10 Burlington Rd., Dublin 8, Ireland

June 11, 2003

Abstract

A fuzzy circle and a fuzzy 3-sphere are constructed as subspaces of fuzzy complex projective spaces, of complex dimension one and three, by modifying the Laplacians on the latter so as to give unwanted states large eigenvalues. This leaves only states corresponding to fuzzy spheres in the low energy spectrum. The construction of a fuzzy circle opens the way to fuzzy tori of any dimension, thus circumventing the problem of power law corrections in possible numerical simulations on these spaces.

1 Introduction

One of the principal goals of the study of field theories on fuzzy spaces is to develop an alternative non-perturbative technique to the familiar lattice one [1]. To date, this new approach in the case of four dimensional field theories has been limited to studies of Euclidean Field theory on $S^2 \times S^2$ [2], \mathbb{CP}^2 [3] and S^4 [4]. All but $S^2 \times S^2$ have additional complications. For example, \mathbb{CP}^2 is not spin but spin^c and S^4 is really a squashed \mathbb{CP}^3 and includes many unwanted massive Kaluza-Klein type modes. Even $S^2 \times S^2$

*bdolan@thphys.may.ie

†denjoe@stp.dias.ie

is not ideal since it has curvature effects that drop off as power corrections rather than exponentially as in the case of toroidal geometries.

The fuzzy approach does, however, have the advantage of preserving continuous symmetries such as the $SU(2)$ symmetry of a round S^2 and does not suffer from fermion doubling [5]. The advantages are gained at the cost of introducing a non-locality associated with the non-commutativity of the fuzzy sphere. There is therefore a balance of advantages and disadvantages associated with the fuzzy approach. The final decision on whether the approach has real advantages over the standard lattice approach should be determined by doing genuine simulations. For this reason Monte Carlo simulations of the fuzzy approach are now in progress. In the lattice approach non-locality is also a problem when Fermions are included. So our expectation is that as far as Monte Carlo simulations are concerned the fuzzy approach will not be competitive with the lattice one until Fermions are included. The approach will gain further advantages in situations where symmetries are more important, so we expect the true power of the approach to emerge when supersymmetry and chiral symmetry are present in a model.

A radically different alternative to the Euclidean Monte Carlo approach becomes available once one has a fuzzy three dimensional space. Such a space has the advantage that it allows one to develop very different non-perturbative methods, since now one can address the non-perturbative questions from a Hamiltonian point of view.

The purpose of this article is to introduce precisely such fuzzy three dimensional spaces. We will begin by presenting a fuzzy version of the circle S_F^1 , from which one can obtain tori of arbitrary dimension. We will then present a fuzzy approximation to the three sphere, S_F^3 . Unfortunately, both of these spaces are still not ideal in that they involve many unwanted additional degrees of freedom which we suppress so that they do not contribute to the low energy physics. The presence of additional degrees of freedom is probably unavoidable as it seems to be the price one pays for the classical space not being a phase space. The three sphere is also curved and hence the results obtained from studies of field theories on this space will approach those of a flat three dimensional space with polynomial corrections. It has, however, the advantages of preserving the full $SO(4)$ symmetry of a round S^3 . From the construction it seems clear that both of these spaces will also be free of fermion doubling problems.

We will restrict our focus here to scalar field theories and demonstrate how the unwanted degrees of freedom can be suppressed so that the limiting large matrix theory of a scalar field theory recovers field theory on the commutative spaces. We will argue that the data specifying the geometries can be cleanly specified by giving a suitable Laplace type operator for the scalar

field, which together with the matrix algebra and its Hilbert space structure give a spectral triple.

Aside from our personal motivations, non-commutative geometry has recently become a very popular area of research from both the point of view of possible new physics in string theory and D -brane theory, [6, 7], and as a new regularisation technique in ordinary quantum field theory, [2]-[4] and [8]-[11]. In both these endeavours “fuzzy” spaces play an important rôle. Roughly speaking a fuzzy space is a finite matrix approximation to the algebra of functions on a continuous manifold, the seminal example being the fuzzy two-sphere, [12]. It has the important property of preserving the isometries of the space that it is approximating. As such the idea can serve as a source of examples related to matrix models in string theory and as a regularisation technique for ordinary quantum field theory. As a regularisation method it provides one that preserves the underlying space-time symmetries and is amenable to numerical computation.

Fuzzy spheres in dimensions other than two were analysed in [13]-[16] but the construction there was incomplete. They also advocate projecting out the unwanted modes and working with a non-associative algebra which we consider unsatisfactory. Also the case of odd spheres works very differently to that of even spheres. An alternative approach for the fuzzy four-sphere, S_F^4 , was given in [4], based on the fact that fuzzy \mathbb{CP}^3 and $\mathbb{CP}^1 \cong S^2$ are well understood [17], and, in the continuum, limit \mathbb{CP}^3 is an S^2 bundle over S^4 .

In this paper we show how the odd dimensional fuzzy spheres S_F^1 and S_F^3 can be extracted from the matrix algebras associated with the fuzzy complex projective spaces \mathbb{CP}_F^1 and \mathbb{CP}_F^3 (an alternative approach to obtaining a finite approximation to $S^3 \cong SU(2)$, based on conformal field theory, was presented in [18]). Our method uses the same idea as [4], although there is no closed finite dimensional matrix algebra for S_F^N unless $N = 2$, the relevant degrees of freedom when $N = 1$ and $N = 3$ are contained in the matrix algebras for \mathbb{CP}^1 and \mathbb{CP}^3 respectively. One can therefore obtain functional integrals for field theories over S_F^1 and S_F^3 by starting with functional integrals over \mathbb{CP}_F^1 and \mathbb{CP}_F^3 and then suppressing the unwanted modes in the exponential so that they do not contribute to the functional integral. Because of the high degree of symmetry inherent in the construction, the unwanted modes can be suppressed simply by using appropriate quadratic Casimirs in the Laplacian. In this way we by-pass the problems associated with the fact that the algebra of matrices associated with functions on the sphere does not close on the sphere, but necessarily lifts into the enveloping complex projective space. In a similar fashion we expect that when a Hamiltonian approach to field theory is developed using these spaces the unwanted modes will cause no difficulties since they can be made arbitrarily difficult to excite.

The paper is organized as follows. In section 2 we summarize how a given geometry is captured in the fuzzy approach. Section 3 then gives our construction of a fuzzy circle, S_F^1 . Section 4 summarises the construction of S_F^4 presented in [4] and in section 5 we present our fuzzy three sphere, S_F^3 . Section 6 gives an alternative construction of S_F^3 which lends itself to a generalisation to S_F^N for any N , [19].

2 Encoding the geometry of a fuzzy space

Fröhlich and Gawędzki [18] (following Connes, [20]) have demonstrated that the abstract triple $(H, \mathcal{A}, \Delta_\gamma)$, where H is the Hilbert space of square integrable functions on the manifold, \mathcal{M} , with Laplace-Beltrami operator Δ_γ , γ being the metric, and $\mathcal{A} = C^\infty(\mathcal{M})$ is the algebra of smooth bounded functions on \mathcal{M} , captures a topological space together with its metrical geometry.

In a similar fashion one can specify a fuzzy space, \mathcal{M}_F , as the sequence of triples

$$\mathcal{M}_F := (H_L, \text{Mat}_{d_L}, \Delta) \quad (1)$$

parameterized by L , where $H = \mathbb{C}^{d_L}$ is the Hilbert space acted on by the complete matrix algebra Mat_{d_L} of dimension d_L with inner product $\langle M, N \rangle = \frac{1}{d_L} \text{Tr}(M^\dagger N)$ and Δ is a suitable Laplacian acting on matrices. One can readily extract information such as the dimension of the space, from these data. The Laplacian comes with a cutoff and so the dimension can be read from the growth of the number of eigenvalues. Sub-leading corrections give such quantities as the Euler characteristic and other information about the space.

The data contained in the triple $(H, \mathcal{A}, \Delta_\gamma)$ are precisely the data that go into the Euclidean action for a scalar field theory on the space M and hence specifying the scalar action is a convenient method of prescribing these data.

In the fuzzy approach the algebra will always be a matrix algebra and we will retain the Hilbert space inner product specified above so the only data from the triple, $(H_L, \text{Mat}_{d_L}, \Delta)$, remaining to be supplied are the permitted matrix dimension, d_L and a realization of the Laplacian, Δ . Once this information is given the fuzzy geometry is specified.

Though it may be convenient to give a map to functions this is not necessary. Once the Laplacian is given its eigenmatrices and spectrum can be used to provide such a map if needed. Suppose for example that a complete set of eigenmatrices is given by $\hat{\Psi}_\lambda$ and the corresponding commutative eigenfunctions are Ψ_λ , then

$$D = \sum_{\lambda}^{d_L^2} \Psi_\lambda \hat{\Psi}_\lambda \quad (2)$$

provides a map to functions with

$$f_M = \frac{1}{d_L} \text{Tr}(DM) \quad (3)$$

the function corresponding to the matrix M . One then gets a $*$ product given by

$$f_M *_D f_N = \frac{1}{d_L} \text{Tr}(DMN) \quad (4)$$

which represents matrix multiplication in terms of an operation on the image functions. The $*$ product depends on D , a different but equivalent one could be obtained by giving a nonzero weighting $c_\lambda(L)$ to the different terms in the sum (2). In the case of \mathbb{CP}^N a particular choice of the $c_\lambda(L)$ will give the diagonal coherent state prescription¹ as discussed in [17].

If the symbol map (2) has the property that

$$\Delta_\gamma f_M = \frac{1}{d_L} \text{Tr}(D\Delta M) \quad (5)$$

where Δ_γ is a natural Laplacian for the space to be approximated, then the spectrum of the fuzzy space will be precisely a cutoff version of that of the commutative space \mathcal{M} . This is precisely what happens in the case of \mathbb{CP}_F^N , see [17].

However, it is convenient to extend the definition of fuzzy space to the case where the spectrum coincides for low lying eigenvalues but deviates for a family of eigenvalues that can be given arbitrarily high value and which correspond to degrees of freedom that have no counterpart in the commutative space \mathcal{M} . This allows us to obtain fuzzy approximations to additional spaces — in particular, as we will see, to tori and the three sphere.

If one takes the Euclidean quantum field theory point of view then the desired geometry appears as that associated with the accessible configurations of the field theory and the deviations are suppressed in a probabilistic fashion.

A successful method of suppressing the unwanted modes would be to add to the scalar action a term $S_I[\Phi]$ which is non-negative for any Φ : zero only for matrices that correspond to functions on \mathcal{M} , and positive for those that do not. The modified action would therefore be of the form $S[\Phi] + \hbar S_I[\Phi]$.

¹In the case where the symbol map is the projector of coherent states the function f_M is referred to as the covariant symbol of the matrix M and since the coefficients $c_\lambda(L)$ are not one it will differ from the corresponding contravariant symbol, see Berezin [21]. The symbol-map is referred to as symmetric when its covariant and contravariant symbols are equal and coincides with the case of $c_\lambda(L) = 1$.

The parameter h should then be chosen large and positive. The probability of any given matrix configuration then takes the form

$$\mathcal{P}[\Phi] = \frac{e^{-S[\Phi]-hS_I[\Phi]}}{Z} \quad (6)$$

where

$$Z = \int d[\Phi] e^{-S[\Phi]-hS_I[\Phi]} \quad (7)$$

is the partition function of the model. If the prescription is to work for free field theories then $S_I[\Phi]$ should be at most quadratic in Φ . This can then be thought of as a modification of the Laplacian in the triple (1).

Furthermore the problem of UV/IR mixing in scalar theories can be removed by including a higher derivative operator in the quadratic term of the field theory that renders all diagrams finite when the matrix size is sent to infinity. With such a prescription it is clear that each diagram has a limiting commutative value and all diagrams must limit to the same value, hence no mixing can occur. This prescription is simpler than the normal ordering prescription proposed in [22] and works for any dimension.

It is therefore clear that the entire problem of constructing a fuzzy approximation to a space is the problem of giving a suitable prescription for the matrix Laplacian.

3 Approximating a circle from a fuzzy sphere

Consider the finite matrix algebra representation of the fuzzy sphere S_F^2 , [12]. The algebra of $(L+1) \times (L+1)$ matrices, which will be denoted by Mat_{L+1} , has the same dimension as the number of degrees of freedom in a spherical harmonic expansion of a function on S^2 , truncated at angular momentum L ,

$$f_L(\theta, \phi) = \sum_{l=0}^L \sum_{m=-l}^l f_{lm} Y_{lm}(\theta, \phi). \quad (8)$$

That is

$$\sum_{l=0}^L (2l+1) = (L+1)^2. \quad (9)$$

The precise identification between a matrix $\Phi \in \text{Mat}_{L+1}$ and a cut-off function $f_L(\theta, \phi)$, as discussed in the preceding section, is not unique, but the possible maps can be given in terms of coherent states or the symmetric symbol map D of (2), and the resulting product of functions is non-commutative for

finite L . It is crucial to our construction that only maps for which the product of functions becomes commutative in the limit $L \rightarrow \infty$ be considered. The symbol map (2) associates orthonormal $(L+1) \times (L+1)$ polarisation tensors \hat{Y}_{lm} with spherical harmonics $Y_{lm}(\theta, \phi)$. The conventions used here will be that

$$\hat{Y}_{lm} = \frac{1}{\sqrt{L+1}} \hat{T}_{lm} \quad (10)$$

where the polarisation tensors \hat{T}_{lm} are those of [23].

The $SO(3)$ symmetric Laplacian on the fuzzy sphere acts on matrices Φ and is represented by the second order Casimir corresponding to the adjoint action of the angular momentum generators L_i in the $(L+1) \times (L+1)$ representation:

$$\mathcal{L}^2 \Phi = [L_i, [L_i, \Phi]]. \quad (11)$$

Hence the action can be taken to be

$$S[\Phi] = \frac{1}{L+1} \text{Tr} \left(\frac{1}{2} \Phi^\dagger \mathcal{L}^2 \Phi + V(\Phi) \right) \quad (12)$$

for some scalar potential $V(\Phi)^\dagger = V(\Phi)$, which is assumed to be bounded below. This action can then be used in a partition function which involves ordinary integration over $(L+1)^2$ degrees of freedom

$$Z = \int \mathcal{D}\Phi e^{-S[\Phi]}. \quad (13)$$

The probability distribution for field configurations is then

$$\mathcal{P}[\Phi] = \frac{e^{-S[\Phi]}}{Z} \quad (14)$$

where $S[\Phi]$ given is by (12). This probability distribution is associated with the geometry $(H_L, \text{Mat}_{L+1}, \mathcal{L}^2)$ which specifies a round fuzzy sphere. This field theory, however, suffers from UV/IR mixing problems [22, 24]. If we add the term $h\mathcal{L}^4$ to the Laplacian and use the triple $(H_L, \text{Mat}_{L+1}, \mathcal{L}^2 + h\mathcal{L}^4)$ the UV/IR mixing problem is removed and we recover field theory on the commutative S^2 in the infinite matrix size limit.

There is no finite matrix approximation to the algebra of functions on S^1 . Nevertheless, the degrees of freedom relevant to a circle are certainly contained in Mat_{L+1} . Focusing on the top harmonic in (8), with $l = L$, the Y_{Lm} contain all $-L \leq m \leq L$ and thus reproduce functions on the circle as $m \rightarrow \infty$. This implies that the partition function and correlation functions for a field theory on a circle can be extracted from that of the fuzzy sphere

by suppressing all the modes with $l < L$ in (13). One way of achieving this is to penalise modes with $l < L$ by giving them a large positive weight in the action. To this end we modify the action (12) to

$$S_h[\Phi] = \frac{1}{L+1} \text{Tr} \left\{ \frac{1}{2} \Phi^\dagger [L_3, [L_3, \Phi]] + \frac{h}{2} \Phi^\dagger (L(L+1) - \mathcal{L}^2) \Phi + V(\Phi) \right\}. \quad (15)$$

All modes with $l < L$ now have the wrong sign for the kinetic energy and, when h is very large, are heavily penalised in the partition function (13), contributing nothing as $h \rightarrow \infty$. In this limit only the modes with $l = L$ remain and these have the correct sign for their kinetic energy, because the term linear in h vanishes on these and only these modes. The wrong sign kinetic energy here would be analogous to an anti-ferromagnetic coupling in a lattice theory and just as in the lattice theory with a anti-ferromagnetic coupling the action here is also bounded below. That the action remains bounded from below is intimately related to the fact that there is an ultraviolet cutoff in the model and therefore a maximum eigenvalue for the Laplacian or equivalently a shortest wavelength.

To see that the commutative algebra of functions on S^1 is recovered in the $l = L$ sector of the fuzzy sphere as $L \rightarrow \infty$, we first decompose the matrix Φ using the basis of polarisation tensors

$$\Phi = \sum_{l=L}^{\infty} \sum_{m=-l}^l \Phi_{lm} \hat{Y}_{lm}. \quad (16)$$

In our conventions (10) the commutator of the polarisation tensors is given by (see e.g. [23] page 191, equation (46))

$$\begin{aligned} [\hat{Y}_{l_1 m_1}, \hat{Y}_{l_2 m_2}] &= \sqrt{\frac{(2l_1+1)(2l_2+1)}{L+1}} \sum_{l=0}^L (-1)^{L-l} \{1 - (-1)^{l_1+l_2+l}\} \\ &\quad \times \left\{ \begin{matrix} l_1 & l_2 & l \\ L/2 & L/2 & L/2 \end{matrix} \right\} C_{l_1 m_1, l_2 m_2}^{lm} \hat{Y}_{lm}, \end{aligned} \quad (17)$$

where $\left\{ \begin{matrix} l_1 & l_2 & l \\ L/2 & L/2 & L/2 \end{matrix} \right\}$ are $6j$ -symbols and $C_{l_1 m_1, l_2 m_2}^{lm}$ are Clebsch-Gordon co-efficients. Now for large L

$$\left\{ \begin{matrix} l_1 & l_2 & l \\ L/2 & L/2 & L/2 \end{matrix} \right\} \approx \frac{1}{\sqrt{L+1}} C_{l_1 0, l_2 0}^{l 0} \quad (18)$$

and $C_{l_1 0, l_2 0}^{l 0} = 0$ when $l_1 + l_2 + l$ is odd. Thus the algebra

$$[\hat{Y}_{l_1 m_1}, \hat{Y}_{l_2 m_2}] \rightarrow 0 \quad (19)$$

is commutative as $L \rightarrow 0$ as promised. In particular

$$[\hat{Y}_{L m_1}, \hat{Y}_{L m_2}] \rightarrow 0 \quad (20)$$

and the top harmonic alone reproduces the commutative algebra of functions on S^1 in the continuum.

To summarize we can encode the geometry specifying a fuzzy circle by the triple

$$S_F^1 := (\mathcal{H}_L, \text{Mat}_{L+1}, \mathcal{L}_3^2 + h(L(L+1) - \mathcal{L}^2)). \quad (21)$$

This picks out the fuzzy circle from the top angular momentum polarization tensor $\hat{Y}_{L,m}$. One could equally pick it out from a lower one, $\hat{Y}_{L_0,m}$ by modifying the term proportional to h to $(L_0(L_0+1) - \mathcal{L}^2)^2$. This latter choice may have advantages for the suppression of UV/IR mixing effects. It corresponds roughly in the fuzzy context to a mixture of ‘nearest neighbour’ and next nearest neighbour ferromagnetic and anti-ferromagnetic couplings.

Having constructed a fuzzy circle it is now clear that there is no obstacle to constructing fuzzy tori of arbitrary dimension, simply by taking products of fuzzy circles. This has the obvious advantage for numerical simulation of avoiding power-law curvature effects.

4 Approximating S^4 from \mathbb{CP}_F^3

We can use a similar procedure to approximate S^3 from a finite approximation to S^4 but first, in this section, we summarise the construction of the fuzzy S^4 from fuzzy \mathbb{CP}^3 given in [4]. The construction utilises the fact that \mathbb{CP}^3 is an S^2 bundle over S^4 and there is a well defined matrix approximation to $\mathbb{CP}^3 \cong SU(3)/U(3)$. The harmonic expansion of a function on \mathbb{CP}^3 requires representations of $SU(4)$ that contain singlets of $U(3)$ under $SU(4) \rightarrow SU(3) \times U(1)$: in terms of Young tableaux these are the representations

$$\begin{array}{|c|c|c|c|} \hline \overbrace{\quad \quad \quad}^n & \overbrace{\quad \quad \quad}^n & & \\ \hline \cdot & \cdot & & \\ \hline & \cdot & & \\ \hline & \cdot & & \\ \hline \end{array} \quad (22)$$

of dimension $\frac{1}{12}(2n+3)(n+2)^2(n+1)^2$. All such representation, for $n \leq L$, appear in the tensor product

$$\begin{array}{|c|c|c|} \hline & & \\ \hline & \cdots & \\ \hline & & \\ \hline & & \\ \hline \end{array} \times \begin{array}{|c|c|c|} \hline & & \\ \hline & \cdots & \\ \hline & & \\ \hline & & \\ \hline \end{array} = \mathbf{1} + \mathbf{15} + \mathbf{84} + \cdots + \frac{(2L+3)(L+2)^2(L+1)^2}{12}. \quad (23)$$

Since the dimension of $\begin{array}{|c|c|c|} \hline & & \\ \hline & \cdots & \\ \hline & & \\ \hline & & \\ \hline \end{array}$ is $d_L := \frac{1}{6}(L+3)(L+2)(L+1)$ the representations in (23) constitute a $d_L \times d_L$ matrix and are thus in one-to-one correspondence with elements Φ of Mat_{d_L} .

Fuzzy \mathbb{CP}^3 is now identified with Mat_{d_L} with an appropriate Laplacian. The most natural Laplacian on \mathbb{CP}_F^3 is the $SU(4)$ invariant one which is the the second order Casimir corresponding to the adjoint action of the $SU(4)$ generators in the $d_L \times d_L$ representation. For future convenience we shall use the fact that $SU(4) \approx Spin(6)$ and denote the generators by J_{AB} , with $A, B = 1, \dots, 6$ and $J_{AB} = -J_{BA}$. The $Spin(6)$ invariant Laplacian on Mat_{d_L} is then

$$\mathcal{L}_{(6)}^2 \Phi = \frac{1}{2} [J_{AB}, [J_{AB}, \Phi]]. \quad (24)$$

As $L \rightarrow \infty$ this corresponds to the continuum $Spin(6)$ invariant Laplacian on \mathbb{CP}^3 .

In the notation of [25] we shall label the $Spin(6)$ irreducible representations by their highest weights (n_1, n_2, n_3) , with n_1, n_2, n_3 either all integers or all half-integers and $n_1 \geq n_2 \geq n_3$. The dimensions of these representations are

$$d^{(6)} = (n_1, n_2, n_3) = \frac{1}{12} \left((n_1+2)^2 - (n_2+1)^2 \right) \left((n_1+2)^2 - n_3^2 \right) \left((n_2+1)^2 - n_3^2 \right) \quad (25)$$

and the eigenvalues of $\mathcal{L}_{(6)}^2$ are

$$C_2^{(6)}(n_1, n_2, n_3) = \left(n_1(n_1+4) + n_2(n_2+2) + n_3^2 \right). \quad (26)$$

The irreducible representations that appear in a harmonic expansion of function on \mathbb{CP}^3 (22) are $(n, n, 0)$ with n an integer, so

$$C_2^{(6)}(n, n, 0) = 2n(n+3). \quad (27)$$

In this notation (23) translates to

$$\overline{\left(\frac{L}{2}, \frac{L}{2}, \frac{L}{2} \right)} \times \left(\frac{L}{2}, \frac{L}{2}, \frac{L}{2} \right) = \sum_{n=0}^L (n, n, 0). \quad (28)$$

The extraction of a fuzzy S^4 from this algebra further relies on the curious fact that there is another possibility for the Laplacian on \mathbb{CP}_F^3 that has a lower symmetry, $SO(5)$, coming from the fact that it is also possible to represent \mathbb{CP}^3 as the coset space $SO(5)/(SU(2) \times U(1))$. In this representation the harmonic expansion of a function on \mathbb{CP}^3 requires all representations of $Spin(5)$ that contain singlets of $SU(2) \times U(1)$ under the embedding

$$\begin{aligned} Spin(5) &\rightarrow SU(2) \times U(1) \\ \mathbf{4} &\rightarrow \mathbf{2}_0 + \mathbf{1}_1 + \mathbf{1}_{-1} \\ \mathbf{5} &\rightarrow \mathbf{2}_1 + \mathbf{2}_{-1} + \mathbf{1}_0 \\ \mathbf{10} &\rightarrow \mathbf{3}_0 + \mathbf{2}_1 + \mathbf{2}_{-1} + \mathbf{1}_2 + \mathbf{1}_{-2} + \mathbf{1}_0. \end{aligned} \tag{29}$$

Irreducible representations of $Spin(5)$ can be labelled by two numbers (n_1, n_2) , either both integers or both half-integers, and $n_1 \geq n_2$. They have dimension

$$d^{(5)}(n_1, n_2) = \frac{1}{6}(n_1 - n_2 + 1)(n_1 + n_2 + 2)(2n_1 + 3)(2n_2 + 1) \tag{30}$$

and second order Casimirs

$$C_2^{(5)}(n_1, n_2) = \left(n_1(n_1 + 3) + n_2(n_2 + 1) \right). \tag{31}$$

From (29) we see that the $Spin(5)$ representations that contain singlets of $SU(2) \times U(1)$ are those with (n_1, n_2) both integers — these are all the tensor representations $T_{a_1 \dots a_n}$, with $a_j = 1, \dots, 5$, and are therefore really representations of $SO(5)$. They are related to the $SO(6)$ representations appearing in (23) by

$$\begin{aligned} SO(6) &\rightarrow SO(5) \\ \mathbf{15} &\rightarrow \mathbf{5} + \mathbf{10} \\ \mathbf{84} &\rightarrow \mathbf{14} + \mathbf{35} + \mathbf{35}', \end{aligned} \tag{32}$$

or, in general,

$$(n, n, 0) \rightarrow \sum_{m=0}^n (n, m). \tag{33}$$

The fact that $\mathbb{CP}^3 \cong SO(5)/(SU(2) \times U(1))$ means that $SO(5)$ acts transitively on \mathbb{CP}^3 and functions on \mathbb{CP}^3 can be expanded in terms of $SO(5)$ irreducible representations (n, m) with an $SO(5)$ invariant Laplacian. As discussed in [4], there is no unique $SO(5)$ invariant Laplacian on Mat_{d_L} but rather any linear combination of the restriction of (24) to $SO(5)$

$$\mathcal{L}_{(5)}^2 \Phi = \frac{1}{2} [J_{ab}, [J_{ab}, \Phi]] \tag{34}$$

with $a, b = 1, \dots, 5$ and

$$\mathcal{L}_{(v)}^2 \Phi = [J_a, [J_a, \Phi]], \quad (35)$$

where $J_a = J_{a6}$, can be used as a Laplacian provided the combination has positive eigenvalues.

The fuzzy S^4 can now be extracted from this by noting that the harmonic expansion of functions on $S^4 \cong SO(5)/SO(4)$ require irreducible representations of $SO(5)$ that contain singlets of $SO(4)$ under $SO(5) \rightarrow SO(4)$ and these are of course the symmetric tensor representations of $SO(5)$, labelled by $(n, 0)$ in the notation above. A Laplacian whose low lying modes are those of S_F^4 can be constructed by penalizing the modes (n, m) in (33) with $m \neq 0$. From (27) and (31) we see that

$$2C_2^{(5)}(n, m) - C_2^{(6)}(n, n, 0) = 2m(m+1), \quad (36)$$

so the Laplacian

$$\mathcal{L}_h^2 = \mathcal{L}_{(6)}^2 + h(2\mathcal{L}_{(5)}^2 - \mathcal{L}_{(6)}^2) \quad (37)$$

has eigenvalues $2n(n+3) + 2hm(m+1)$ and states with $m > 0$ will be suppressed in a functional integral for large h . The parameter h here is acting like a “squashing” parameter, $h = 0$ is the “round” $SO(6)$ invariant metric on \mathbb{CP}^3 , while $h \neq 0$ breaks this symmetry down to $SO(5)$. The lowest permitted value for h is $h = -1$ and we have $\mathcal{L}_{-1}^2 = 2\mathcal{L}_v^2$, we see that this Laplacian is rather singular in the large L limit as the representation (n, n) for large n develops a zero eigenvalue.

The family of actions

$$S_h[\Phi] = \frac{1}{d_L} \text{Tr} \{ \Phi^\dagger \mathcal{L}_h^2 \Phi + V(\Phi) \} \quad (38)$$

give a field theory on squashed \mathbb{CP}_F^3 for $h > -1$. Furthermore as $h \rightarrow \infty$ modes with $m > 0$ are completely suppressed in a functional integral and (38) corresponds to a field theory on S_F^4 . Note that it does not matter whether we use $\mathcal{L}_{(6)}^2$, $2\mathcal{L}_{(5)}^2$ or $2\mathcal{L}_{(v)}^2$ for the first term on the right-hand side of (37) — when the constraint $m = 0$ is imposed all three are the same operator.

5 S_F^3 from S_F^4

We can build on the construction of the last section to get a Laplacian whose low lying modes are those associated with a field on S_F^3 by using the same trick as in section 3 to pick out the top mode, $n = L$, of the S_F^4 . As an

irreducible representation of $SO(5)$ this is the representation $(L, 0)$ with dimension

$$d^{(5)}(L, 0) = \frac{1}{6}(2L+3)(L+2)(L+1). \quad (39)$$

The harmonic expansion of a function on $S^3 \cong SU(2)$ requires all irreducible representations of $SU(2)$, both integral and half-integral,

$$f_L(\theta, \phi, \psi) = \sum_{j=0, \frac{1}{2}, \dots}^{L/2} \sum_{\bar{m}=-j}^j \sum_{m=-j}^j f_{\bar{m}, m}^j D_{\bar{m}, m}^j(\theta, \phi, \psi), \quad (40)$$

where (θ, ϕ, ψ) are Euler angles and $D^j(\theta, \phi, \psi)$ are the Wigner D -matrices. The key to extracting S_F^3 from S_F^4 is the observation that the total number of degrees of freedom in (40) is

$$\sum_{j=0, \frac{1}{2}, \dots}^{L/2} (2j+1)^2 = \frac{1}{6}(2L+3)(L+2)(L+1) \quad (41)$$

which is the same as $d^{(5)}(L, 0)$ in (39). This is because the top mode (or indeed any mode $(n, 0)$) of S_F^4 has the representation content of an S_F^3 .

The top $SO(5)$ mode of S_F^4 can now be picked out by penalising the modes with $n < L$ with a wrong sign kinetic-energy term, i.e. using an anti-ferromagnetic like coupling. To this end we define the Laplacian

$$\mathcal{L}_{h, h'}^2 \Phi = \frac{1}{2} [J_{\alpha\beta}, [J_{\alpha\beta}, \Phi]] + h' (2L(L+3) - \mathcal{L}_{(6)}^2) \Phi + h (2\mathcal{L}_{(5)}^2 - \mathcal{L}_{(6)}^2) \Phi, \quad (42)$$

with $\alpha, \beta = 1, \dots, 4$. For finite h and h' , both > 0 , this is a positive operator and contains all modes on \mathbb{CP}_F^3 . As $h \rightarrow \infty$ modes not relevant to S_F^4 are sent to infinity and, finally, modes not relevant to S_F^3 are sent to infinity when $h' \rightarrow \infty$. For very large h and h' the low lying eigenvalues are therefore precisely those of S_F^3 . In a functional integral for a scalar field based on this Laplacian we recover scalar field theory on S_F^3 in the large h and h' limit. Field theory on S_F^3 therefore arises from the double limit $h, h' \rightarrow \infty$ in the action

$$S_{h, h'}[\phi] = \frac{1}{d_L} \text{Tr} \{ \Phi^\dagger \mathcal{L}_{h, h'}^2 \Phi + V(\Phi) \} \quad (43)$$

with $\Phi \in \text{Mat}_{d_L}$.

6 An alternative construction of S_F^3

The constructions described up till now have relied on matrix approximations to \mathbb{CP}^N , specifically $\mathbb{CP}^1 \cong S^2$ and \mathbb{CP}^3 . There is however another construction for S_F^3 based on the orthogonal Grassmannian $SO(5)/(SO(3) \times SO(2))$.

This is a co-adjoint orbit and therefore a well-defined finite matrix approximation to the algebra of functions on this Grassmannian exists. This space is not the same as \mathbb{CP}^3 : it arises from a different embedding of $SU(2) \times U(1)$ into $Spin(5)$, characterised by

$$\begin{aligned} Spin(5) &\rightarrow SU(2) \times U(1) \\ \mathbf{4} &\rightarrow \mathbf{2}_1 + \mathbf{2}_{-1} \\ \mathbf{5} &\rightarrow \mathbf{3}_0 + \mathbf{1}_2 + \mathbf{1}_{-2} \\ \mathbf{10} &\rightarrow \mathbf{3}_2 + \mathbf{3}_{-2} + \mathbf{3}_0 + \mathbf{1}_0. \end{aligned} \tag{44}$$

In particular the $\mathbf{5}$ does not give a singlet under this embedding and so must be excluded from the harmonic expansion on this space — it is clearly not the same space as \mathbb{CP}^3 . The representation content here is such that

$$SO(5)/(SO(3) \times SO(2)) \cong Sp(2)/U(2), \tag{45}$$

which is known not to admit a spin structure [26]. The expansion of a function on the Grassmannian $SO(5)/(SO(3) \times SO(2))$ can be obtained from that on $\mathbb{CP}^3 \cong SO(5)/(SU(2) \times U(1))$ simply by omitting all the odd rank tensors from the latter. In this way $SO(5)/(SO(3) \times SO(2))$ follows from moding out the $SO(5)$ representation of \mathbb{CP}^3 by the \mathbb{Z}_2 action $T_a \rightarrow -T_a$ on the $\mathbf{5}$, so

$$SO(5)/(SO(3) \times SO(2)) \cong \mathbb{CP}^3/\mathbb{Z}_2. \tag{46}$$

As a side remark we note that $SO(5)/(SO(3) \times SO(2))$ is an S^2 bundle over the real projective space \mathbb{RP}^4 .

In the notation of section 4 the even rank tensor representations of $SO(5)$ are (n, m) with $n + m = 2l$, and these have dimension

$$d^{(5)}(2l - m, m) = \frac{1}{6}(2l - 2m + 1)(2l + 2)(4l - 2m + 3)(2m + 1). \tag{47}$$

So the total number of degrees of freedom for $m \leq l$ and $0 \leq l \leq L/2$ (L even) is

$$\sum_{l=0}^{L/2} \sum_{m=0}^l d^{(5)}(2l - m, m) = \left[\frac{(L+4)(L+3)(L+2)}{24} \right]^2, \tag{48}$$

which is the same as that of $\text{Mat}_{d'_L}$ with

$$d'_L = \frac{1}{24}(L+4)(L+3)(L+2). \quad (49)$$

The action on the fuzzy Grassmannian looks identical to that on a squashed \mathbb{CP}_F^3 ,

$$S[\Phi] = \frac{1}{d'_L} \text{Tr} \left\{ \Phi^\dagger \mathcal{L}_{(5)}^2 \Phi + V(\Phi) \right\}, \quad (50)$$

except that the matrix algebras are restricted to those of size d'_L in (49) containing only even rank tensor representations $SO(5)$, $d^{(5)}(2l-m, m)$.

The harmonic expansion of a function on S_F^3 is contained in $\text{Mat}_{d'_L}$ because the top representation for a given L , $2l = L$ with $m = 0$, has dimension (39) and, as observed in the previous section, this is a sum of the dimensions of the $SU(2)$ representations required for a harmonic expansion on S^3 , (41).

If we can penalise all modes with $2l < L$ and $m \neq 0$ for $2l = L$ in a functional integral over $SO(5)/(SO(3) \times SO(2))$ then we will really be doing a functional integral over S_F^3 . This is easily achieved since $2l = L$ and $m = 0$ has the largest second order Casimir,

$$C_2^{(5)}(L, 0) = L(L+3), \quad (51)$$

of all the $SO(5)$ representations in $\text{Mat}_{d'_L}$. In the now familiar manner the unwanted modes in the functional integral over $SO(5)/(SO(3) \times SO(2))$ can be suppressed by using the Laplacian

$$\mathcal{L}_{h'}^2 = \frac{1}{2} [J_{\alpha\beta}, [J_{\alpha\beta}, \cdot]] + h' (L(L+3) - \mathcal{L}_{(5)}^2), \quad (52)$$

which acts on fields $\Phi \in \text{Mat}_{d'_L}$ and L even. The unwanted modes are completely eliminated in the limit $h' \rightarrow \infty$, giving S_F^3 truncated at level L . The constraint that L is even does not change the fact that we get the full continuum S^3 as $L \rightarrow \infty$.

7 Conclusions

By starting with the known finite matrix algebras for \mathbb{CP}^3 and \mathbb{CP}^1 , the fuzzy \mathbb{CP}_F^3 and the fuzzy sphere $\mathbb{CP}_F^1 \cong S_F^2$, finite functional integrals for scalar field theories on S_F^3 and S_F^1 have been constructed. The geometry of a fuzzy space is specified by a triple $(H_L, \text{Mat}_{d_L}, \Delta)$ and, although there is no known closed associated algebra giving a fuzzy S^1 as a triple directly,

\mathbb{CP}_F^1 nevertheless contains the states required for a S_F^1 plus other unwanted states. The unwanted states are given large eigenvalues by modifying the Laplacian on \mathbb{CP}_F^1 , as in equation (21), leaving only the states of S_F^1 in the low energy spectrum of the Laplacian. In a similar way \mathbb{CP}_F^3 contains the states necessary for a fuzzy description of S^3 (via S_F^4) and the Laplacian on \mathbb{CP}_F^3 can be modified, as in equation (42), so that states not related to S_F^3 are given large eigenvalues, leaving only S_F^3 states in the low energy spectrum.

An alternative construction of S_F^3 , based on suppressing modes on a fuzzy version of the orthogonal Grassmannian $SO(5)/(SO(3) \times SO(2))$, has been presented in section 6. This has the advantage of having a natural extension to S_F^N for any N , [19]. Thus S_F^3 can be obtained either in two steps, via the fuzzy S_F^4 constructed in [4], $\mathbb{CP}_F^3 \rightarrow S_F^4 \rightarrow S_F^3$, or alternatively in a single step from the fuzzy version of $SO(5)/[SO(3) \times SO(2)]$ as described in section 6.

The construction of the fuzzy circle allows fuzzy tori to be defined in an obvious way, by taking products of fuzzy circles, thus opening the way to numerical simulations on tori while preserving the full $U(1) \times \dots \times U(1)$ isometry group and avoiding the fermion doubling problem [5]. One first writes down a finite functional integral for a field theory on $S_F^2 \times \dots \times S_F^2$, which contain the modes relevant for propagation $S_F^1 \times \dots \times S_F^1$ in its spectrum, and then damps the unwanted modes. This can be done, in a manner that preserves the isometries of the torus, by introducing appropriate combinations of second order Casimirs into the propagators.

It is a pleasure to acknowledge Peter Prešnajder, Harald Grosse and Daniel Roggenkamp for helpful discussions.

References

- [1] I. Montvay and G. Munster, *Quantum Field on a Lattice*, Cambridge, 1997.
- [2] S. Vaidya and B. Ydri, On the Origin of the UV-IR Mixing in Non-commutative Matrix Geometry, [hep-th/0305201](#).
- [3] G. Alexanian, A.P. Balachandran, G. Immirzi and B. Ydri, *Fuzzy \mathbb{CP}^2 J. of Geom. and Phys.* **42** (2002) 28, [hep-th/0103023](#).
- [4] Denjoe O'Connor and Julieta Medina, *Scalar Field Theory on Fuzzy S^4* , [hep-th/0212170](#).

- [5] A. P. Balachandran, T. R. Govindarajan and B. Ydri, *Mod. Phys. Lett.* **A15** (2000) 1279, hep-th/9911087; A. P. Balachandran, T. R. Govindarajan and B. Ydri, *Fermion doubling problem and noncommutative geometry II*, hep-th/0006216; A. P. Balachandran and G. Immirzi, *The Fuzzy Ginsparg-Wilson Algebra: A Solution of the Fermion Doubling Problem*, hep-th/0301242.
- [6] A. Yu. Alekseev, A. Recknagel and V. Schomerus, *Non-commutative World-volume Geometries: Branes on $SU(2)$ and Fuzzy Spheres*, *JHEP* **9909**, (1999) 023, hep-th/9908040.
- [7] Yasuaki Hikida, Masatoshi Nozaki and Yuji Sugawara, *Nucl. Phys.* **B617**, (2001) 117, hep-th/0101211.
- [8] H. Grosse, C. Klimčík and P. Prešnajder, *Int. J. Theor. Phys.* **35**, (1996) 231, hep-th/9505175; H. Grosse and A. Strohmaier, *Lett. Math. Phys.* **48**, (1999) 163, hep-th/9902138.
- [9] H. Grosse, C. Klimčík and P. Prešnajder, *Comm. Math. Phys.* **178**, (1996) 507; H. Grosse and P. Prešnajder, *Lett. Math. Phys.* **46**, (1998) 61.
- [10] P. Prešnajder, *J. Math. Phys.* **41** (2000) 2789, hep-th/9912050.
- [11] A.P. Balachandran and S. Vaidya, *Instantons and Chiral Anomaly in Fuzzy Physics Int. J. Mod. Phys.* **A16**, (2001) 17, hep-th/9910129; S. Baez, A. P. Balachandran, S. Vaidya and B. Ydri, *Monopoles and Solitons in Fuzzy Physics, Comm. Math. Phys.* **208**, (2000) 787, hep-th/9811169.
- [12] J. Madore, *Class. Quant. Grav.* **9** (1992) 69.
- [13] S. Ramgoolam, *On spherical harmonics for fuzzy spheres in diverse dimensions*, *Nucl. Phys.* **B610** (2001) 461, hep-th/0105006.
- [14] Yusuke Kimura, *On higher dimensional fuzzy spherical branes* hep-th/0301055.
- [15] S. Ramgoolam, *Higher dimensional geometries related to fuzzy odd-dimensional spheres JHEP* **0210** (2002) 064, hep-th/0207111.
- [16] P. M. Ho and S. Ramgoolam, *Higher Dimensional Geometries from Matrix Brane constructions*, *Nucl. Phys.* **B627** (2002) 266, hep-th/0111278.

- [17] A.P. Balachandran, B.P. Dolan, J. Lee, X. Martin and D. O'Connor, *Fuzzy Complex Projective Spaces and their Star-products*, *J. Geom. and Phys.* **43** (2002) 184, [hep-th/0107099](#).
- [18] J. Frohlich and K. Gawędzki, *Conformal Field Theory and Geometry of Strings*, Lectures given at Mathematical Quantum Theory Conference, Vancouver, Canada, 4-8 Aug 1993. Published in Vancouver 1993, Proceedings, Mathematical quantum theory, **Vol. 1** 57-97, [hep-th/9310187](#).
- [19] B.P. Dolan, Denjoe O'Connor and P. Prešnajder, in preparation.
- [20] A. Connes. *Noncommutative Geometry*, Academic Press, London, 1994.
- [21] F.A. Berezin, *General Concept of Quantization*, *Comm. Math. Phys.* **40**, (1975) 153.
- [22] Brian P. Dolan, Denjoe O'Connor and P. Prešnajder, *Matrix ϕ^4 Models on the Fuzzy Sphere and their Continuum Limits*, *JHEP* **0203** (2002) 013, [hep-th/0109084](#).
- [23] D.A. Varshalovich, A.N. Moskalev and V.K. Khersonskii, *Quantum Theory of Angular Momentum*, World Scientific (1988)
- [24] Chong-Sun Chu, John Madore and Harold Steinacker, *Scaling Limits of the Fuzzy Sphere at one Loop*, *JHEP* **0108** (2001) 038, [hep-th/0106205](#).
- [25] W. Fulton and J. Harris, *Representation Theory: a first course*, Graduate Texts in Mathematics 129, Springer, (1991).
- [26] B.P. Dolan and C. Nash, *JHEP* **0207** (2002) 057, [hep-th/0207007](#).