

Title	On the Generalised Random Energy Model
Creators	Dorlas, T. C. and Dukes, W.M.B.
Date	2001
Citation	Dorlas, T. C. and Dukes, W.M.B. (2001) On the Generalised Random Energy Model. (Preprint)
URL	https://dair.dias.ie/id/eprint/441/
DOI	DIAS-STP-01-12



ON THE GENERALISED RANDOM ENERGY MODEL

T.C. DORLAS & W.M.B. DUKES

School of Theoretical Physics
Dublin Institute for Advanced Studies
10 Burlington Road, Dublin 4, Ireland

June 29, 2001

Abstract

The Generalised Random Energy Model is a generalisation of the Random Energy Model introduced by Derrida to mimic the ultrametric structure of the Parisi solution of the Sherrington-Kirkpatrick model of a spin glass. It was solved exactly in two special cases by Derrida and Gardner. A rigorous analysis by Capocaccia et al. claimed to give a complete solution for the thermodynamics of the model in the general case. Here we use Large Deviation Theory to analyse the model along the lines followed by Dorlas and Wedagedera for the Random Energy Model. The resulting variational expression for the free energy is the same as that found by Capocaccia et al. We show that it can be evaluated in a very simple way. We find that the answer given by Capocaccia et al. is incorrect.

1 Definition of the GREM

The generalised random energy model (GREM) was introduced by Derrida [3] as a generalisation of his random energy model (see [2]) of a spin glass in order to incorporate some correlations between energy levels. Whereas in the random energy model all energy levels E_i are independent random variables, and the partition function is given by

$$Z_N(\beta) = \sum_{i=1}^{2^N} e^{-\beta E_i},$$

the energy levels of the generalised model have a tree-like structure. The tree is defined by a number of levels n and for each level $k = 1, \dots, n$, a number $\alpha_k \in (1, 2)$ determining the number of branches per node. (See Figure 1.) To make the total number of highest-level branches in the tree add up to 2^N as before, we assume that $\prod_{i=1}^n \alpha_k = 2$. For each $k = 1, \dots, n$ there are $(\alpha_1 \cdots \alpha_k)^N$ independent random variables $\{E_j^{(k)}\}$, distributed according to $\rho_N^{(k)}$ with density

$$\rho_N^{(k)}(E) = \frac{1}{\sqrt{a_k \pi N J^2}} e^{-E^2/a_k N J^2}, \quad (1.1)$$

where the positive numbers a_k satisfy $\sum_{k=1}^n a_k = 1$. (Obviously, in general α_k^N is not an integer, but we can take its integer part which is very nearly the same for large N . We shall disregard the difference in the following.)

The partition function of the GREM is defined by

$$Z_N(\beta) = \sum_{i_1=1}^{\alpha_1^N} \sum_{i_2=(i_1-1)\alpha_2^N+1}^{\alpha_1\alpha_2^N} \cdots \sum_{i_n=(i_{n-1}-1)\alpha_n^N+1}^{i_{n-1}\alpha_n^N} \exp \left[-\beta \left(\sum_{k=1}^n E_{i_k}^{(k)} \right) \right]. \quad (1.2)$$




Figure 1: The tree-like structure of the GREM. The nodes on the n^{th} layer represent the configurations. The energy of any configuration is the sum of the energies on the branches up to the source node.

This formula is best understood by referring to Figure 1. As usual the free energy is defined by

$$f(\beta) = -\frac{1}{\beta} \lim_{N \rightarrow \infty} \frac{1}{N} \ln Z_N(\beta). \quad (1.3)$$

We shall prove that this limit exists almost surely w.r.t. the distribution of the energies $\{E_i^{(k)}\}$. To do this, we introduce the random distribution functions $F_N(x_1, \dots, x_n)$ and $\bar{F}_N(x_1, \dots, x_n)$ as follows.

$$\begin{aligned} F_N(x_1, \dots, x_n) &:= \frac{1}{2^N} \sum_{i_1=1}^{\alpha_1^N} \sum_{i_2=(i_1-1)\alpha_2^N+1}^{\alpha_1^N \alpha_2^N} \cdots \sum_{i_n=(i_{n-1}-1)\alpha_n^N+1}^{i_{n-1}\alpha_n^N} \mathbb{I}_{i_1}^{(1)} \mathbb{I}_{i_2}^{(2)} \cdots \mathbb{I}_{i_n}^{(n)}, \\ \bar{F}_N(x_1, \dots, x_n) &:= \frac{1}{2^N} \sum_{i_1=1}^{\alpha_1^N} \sum_{i_2=(i_1-1)\alpha_2^N+1}^{\alpha_1^N \alpha_2^N} \cdots \sum_{i_n=(i_{n-1}-1)\alpha_n^N+1}^{i_{n-1}\alpha_n^N} \mathbb{I}_{i_1}^{(1)} \mathbb{I}_{i_2}^{(2)} \cdots \mathbb{I}_{i_n}^{(n)}, \end{aligned} \quad (1.4)$$

where we use the notation $\mathbb{I}_i^{(k)} = \mathbb{1}\{E_i^{(k)} > Nx_k\}$ and $\bar{\mathbb{I}}_i^{(k)} = \mathbb{1}\{E_i^{(k)} \leq Nx_k\}$. We also define G_N and \bar{G}_N as

$$\begin{aligned} G_N(x_1, \dots, x_n) &= \int_{-\infty}^{Nx_1} \cdots \int_{-\infty}^{Nx_n} \rho_N^{(1)}(E_1) \cdots \rho_N^{(n)}(E_n) dE_n \cdots dE_1, \\ \bar{G}_N(x_1, \dots, x_n) &:= \int_{Nx_1}^{+\infty} \cdots \int_{Nx_n}^{+\infty} \rho_N^{(1)}(E_1) \cdots \rho_N^{(n)}(E_n) dE_n \cdots dE_1. \end{aligned}$$

We will abbreviate $\bar{G}_N(x_1, \dots, x_n)$ to \bar{G}_N and $\bar{F}_N(x_1, \dots, x_n)$ to \bar{F}_N . Let us also use as short-hand,

$$p_i := \mathbb{P}(E^{(i)} > Nx_i).$$

Note that $\bar{G}_N = p_1 p_2 \cdots p_n$. In the following section we prove a large deviation property (LDP) for the distribution functions F_N analogous to that of Dorlas and Wedagedera [5].

2 The Rate Function

Theorem 2.1 *The sequence of measures $\mu_N(x_1, \dots, x_n)$ with distribution function $F_N(x_1, \dots, x_n)$ satisfies a LDP with rate function $I(x_1, \dots, x_n)$ where*

$$I(x_1, \dots, x_n) = \begin{cases} \frac{1}{J^2} \sum_{1 \leq i \leq n} \frac{x_i^2}{a_i}, & \text{if } (x_1, \dots, x_n) \in \Psi(J; a_1, \dots, a_n; \alpha_1, \dots, \alpha_n), \\ +\infty, & \text{otherwise,} \end{cases}$$

where the region $\Psi(J; a_1, \dots, a_n; \alpha_1, \dots, \alpha_n)$ is given by

$$\left\{ (x_1, \dots, x_n) \in \mathbb{R}^n \mid \sum_{i=1}^k \frac{x_i^2}{a_i} \leq J^2 \sum_{i=1}^k \ln \alpha_i, \text{ for all } 1 \leq k \leq n \right\}$$

PROOF: First we do the case for $(x_1, \dots, x_n) \in \Psi(J; a_1, \dots, a_n; \alpha_1, \dots, \alpha_n)$. By Chebyshev's inequality, for all $\epsilon \in (0, 1)$,

$$\mathbb{P}(|\bar{F}_N - \bar{G}_N| > \epsilon \bar{G}_N) \leq \frac{1}{\epsilon^2 \bar{G}_N^2} \mathbb{E}(|\bar{F}_N - \bar{G}_N|^2).$$

Now $\mathbb{E}(|\bar{F}_N - \bar{G}_N|^2) = \mathbb{E}(\bar{F}_N^2) - 2\bar{G}_N \mathbb{E}(\bar{F}_N) + \bar{G}_N^2$ and

$$\begin{aligned} \mathbb{E}(\bar{F}_N) &= \frac{1}{2^N} \mathbb{E} \left(\sum_{i_1=1}^{\alpha_1^N} \sum_{i_2=(i_1-1)\alpha_2^N+1}^{i_1\alpha_2^N} \cdots \sum_{i_n=(i_{n-1}-1)\alpha_n^N+1}^{i_{n-1}\alpha_n^N} \mathbb{I}_{i_1}^{(1)} \mathbb{I}_{i_2}^{(2)} \cdots \mathbb{I}_{i_n}^{(n)} \right) \\ &= \frac{1}{2^N} \alpha_1^N p_1 \alpha_2^N p_2 \cdots \alpha_n^N p_n \\ &= p_1 p_2 \cdots p_n \\ &= \bar{G}_N. \end{aligned}$$

To obtain $\mathbb{E}(\bar{F}_N^2)$ we introduce some new notation. Let

$$\mathcal{B}_k := \mathbb{E} \left\{ \sum_{\substack{i_k=\alpha_k^N \\ j_k=\alpha_k^N \\ i_k=1 \\ j_k=1}}^{i_k=\alpha_k^N} \mathbb{I}_{i_k}^{(k)} \mathbb{I}_{j_k}^{(k)} \sum_{\substack{i_{k+1}=(i_k-1)\alpha_{k+1}^N+1 \\ j_{k+1}=(j_k-1)\alpha_{k+1}^N+1}}^{i_{k+1}=i_k\alpha_{k+1}^N} \mathbb{I}_{i_{k+1}}^{(k+1)} \mathbb{I}_{j_{k+1}}^{(k+1)} \cdots \sum_{\substack{i_n=(i_{n-1}-1)\alpha_n^N+1 \\ j_n=(j_{n-1}-1)\alpha_n^N+1}}^{i_n=i_{n-1}\alpha_n^N} \mathbb{I}_{i_n}^{(n)} \mathbb{I}_{j_n}^{(n)} \right\}.$$

Now notice that the following recursion holds:

$$\mathcal{B}_k = \alpha_k^N p_k (\mathcal{B}_{k+1} + (\alpha_k^N - 1) p_k (\alpha_{k+1}^N p_{k+1} \cdots \alpha_n^N p_n)^2),$$

for all $1 \leq k < n$. The initial value is $\mathcal{B}_n = \alpha_n^N p_n + (\alpha_n^{2N} - \alpha_n^N) p_n^2$ but this may be obtained by defining $\mathcal{B}_{n+1} := 1$ and applying the above recursion for $k = n$. Notice also that $\mathbb{E}(\bar{F}_N^2) = \frac{1}{2^{2N}} \mathcal{B}_1$.

Alongside the above recursion, let us define a sequence \mathcal{D}_k by which we upper bound \mathcal{B}_k . Let $\mathcal{D}_{n+1} := 1$ and define

$$\mathcal{D}_k = y_k (\mathcal{D}_{k+1} + y_k (y_{k+1} \cdots y_n)^2).$$

This gives rise to

$$\mathcal{D}_1 = y_1 y_2 \cdots y_n (1 + y_n + y_{n-1} y_n + \cdots + y_1 y_2 \cdots y_n).$$

If we now take $y_k = \alpha_k^N p_k$, then it is clear that $D_k \geq \mathcal{B}_k$ for all $1 \leq k \leq n$, hence the following bound:

$$\begin{aligned} \mathbb{E}(\bar{F}_N^2) &= \frac{1}{2^{2N}} \mathcal{B}_1 \\ &\leq \frac{1}{2^{2N}} \mathcal{D}_1 \\ &= \left(\prod_{1 \leq k \leq n} \alpha_k^N p_k \right) \left(1 + \sum_{k=1}^n \alpha_k^N p_k \cdots \alpha_n^N p_n \right) \\ &= 2^N \bar{G}_N \left(1 + \sum_{k=1}^n \alpha_k^N p_k \cdots \alpha_n^N p_n \right). \end{aligned}$$

Thus

$$\begin{aligned} \frac{1}{\epsilon^2 \bar{G}_N^2} \mathbb{E}(|\bar{F}_N - \bar{G}_N|^2) &\leq \frac{1}{\epsilon^2 \bar{G}_N^2} \left\{ \frac{1}{2^{2N}} 2^N \bar{G}_N \left(1 + \sum_{k=1}^n \alpha_k^N p_k \cdots \alpha_n^N p_n \right) - \bar{G}_N^2 \right\} \\ &= \frac{1}{\epsilon^2 \bar{G}_N^2} \left\{ \frac{1}{2^N} \bar{G}_N \left(1 + \sum_{k=1}^n \alpha_k^N p_k \cdots \alpha_n^N p_n \right) - \bar{G}_N^2 \right\} \\ &= \frac{1}{\epsilon^2 \bar{G}_N^2} \left\{ \frac{1}{2^N} \bar{G}_N \left(1 + 2^N \bar{G}_N + \sum_{k=2}^n \alpha_k^N p_k \cdots \alpha_n^N p_n \right) - \bar{G}_N^2 \right\} \\ &= \frac{1}{\epsilon^2 2^N \bar{G}_N} \left\{ 1 + \sum_{k=2}^n \alpha_k^N p_k \cdots \alpha_n^N p_n \right\} \\ &= \frac{1}{\epsilon^2 2^N \bar{G}_N} + \frac{1}{\epsilon^2} \sum_{k=2}^n \frac{\alpha_k^N p_k \cdots \alpha_n^N p_n}{2^N p_1 p_2 \cdots p_n} \\ &= \frac{1}{\epsilon^2 2^N \bar{G}_N} + \frac{1}{\epsilon^2} \sum_{k=2}^n \frac{1}{\alpha_1^N p_1 \cdots \alpha_{k-1}^N p_{k-1}} \\ &= \frac{1}{\epsilon^2 2^N \bar{G}_N} + \frac{1}{\epsilon^2} \sum_{k=1}^{n-1} \frac{1}{\alpha_1^N p_1 \cdots \alpha_k^N p_k} \\ &= \frac{1}{\epsilon^2} \sum_{k=1}^n \frac{1}{\alpha_1^N p_1 \cdots \alpha_k^N p_k}. \end{aligned} \tag{2.1}$$

Using the inequality $\int_a^\infty e^{-u^2/2} du > \frac{1}{a + a^{-1}} e^{-a^2/2}$ (see McKean [6]), the k^{th} term in this sum is bounded above by

$$\frac{1}{\epsilon^2} \left\{ \prod_{1 \leq i \leq k} \frac{\sqrt{\pi}(2N x_i^2 + a_i J^2)}{x_i J \sqrt{a_i N}} \right\} \exp \left[\frac{N}{J^2} \left(-J^2 \sum_{i=1}^k \ln \alpha_i + \sum_{1 \leq i \leq k} \frac{x_i^2}{a_i} \right) \right].$$

which will converge if and only if $\sum_{1 \leq i \leq k} \frac{x_i^2}{a_i} < J^2 \sum_{1 \leq i \leq k} \ln \alpha_i$. Thus it is seen that equation (2.1)

converges if all the sums of its individual terms converges. The values for which this happens are precisely those which define the region $\Psi(J; a_1, \dots, a_n; \alpha_1, \dots, \alpha_n)$ as stated in the lemma. Introducing the events

$$\mathcal{A}_N = \left\{ \{E_{i_1}^{(1)}, \dots, E_{i_n}^{(n)}\} \mid |\bar{G}_N - \bar{F}_N| > \epsilon \bar{G}_N \right\},$$

we see that $\sum_N \mathbb{P}(\mathcal{A}_N) < +\infty$. Hence by the Borel-Cantelli Lemma,

$$\mathbb{P}\left(\bigcap_{v=1}^{\infty} \bigcup_{N=v}^{\infty} \mathcal{A}_N\right) = 0.$$

This means that with probability 1,

$$\{E_{i_1}^{(1)}, \dots, E_{i_n}^{(n)}\} \in \left(\bigcap_{v=1}^{\infty} \bigcup_{N=v}^{\infty} \mathcal{A}_N\right)^C = \bigcup_{v=1}^{\infty} \bigcap_{N=v}^{\infty} \mathcal{A}_N^C$$

In other words, for almost all $\{E_{i_1}^{(1)}, \dots, E_{i_n}^{(n)}\}$ there exists a $v \in \mathbb{N}$ such that for all $N \geq v$, $\{E_{i_1}^{(1)}, \dots, E_{i_n}^{(n)}\} \in \mathcal{A}_N^C$. Hence $\overline{F}_N = \overline{G}_N$ with probability 1 for all $N \geq v$.

For the case $(x_1, \dots, x_n) \notin \Psi(J; a_1, \dots, a_n; \alpha_1, \dots, \alpha_n)$, then it must hold that

$$\sum_{1 \leq i \leq k} \frac{x_i^2}{a_i} > J^2 \sum_{1 \leq i \leq k} \ln \alpha_i$$

for some k with $1 \leq k \leq n$. We may now upper bound the function $\overline{F}_N(x_1, \dots, x_n)$ by

$$\begin{aligned} \overline{F}_N(x_1, \dots, x_n) &\leq \frac{1}{2^N} \alpha_{k+1}^N \cdots \alpha_n^N \sum_{i_1=1}^{\alpha_1^N} \sum_{i_2=(i_1-1)\alpha_2^N+1}^{i_1\alpha_2^N} \cdots \sum_{i_k=(i_{k-1}-1)\alpha_k^N+1}^{i_{k-1}\alpha_k^N} \mathbb{1}_{i_1}^{(1)} \mathbb{1}_{i_2}^{(2)} \cdots \mathbb{1}_{i_k}^{(k)} \\ &= \frac{1}{\alpha_1^N \cdots \alpha_k^N} \sum_{i_1=1}^{\alpha_1^N} \sum_{i_2=(i_1-1)\alpha_2^N+1}^{i_1\alpha_2^N} \cdots \sum_{i_k=(i_{k-1}-1)\alpha_k^N+1}^{i_{k-1}\alpha_k^N} \mathbb{1}_{i_1}^{(1)} \mathbb{1}_{i_2}^{(2)} \cdots \mathbb{1}_{i_k}^{(k)} \\ &=: H_N(x_1, \dots, x_k) \end{aligned}$$

We will show that $H_N(x_1, \dots, x_k) = 0$ with probability 1 if N is large enough. We have

$$\begin{aligned} &\left\{ \{E_{i_1}^{(1)}, \dots, E_{i_k}^{(k)}\} \mid H_N(x_1, \dots, x_k) = 0 \right\} \\ &= \left\{ \{E_{i_1}^{(1)}, \dots, E_{i_k}^{(k)}\} \mid \sum_{i_1=1}^{\alpha_1^N} \sum_{i_2=(i_1-1)\alpha_2^N+1}^{i_1\alpha_2^N} \cdots \sum_{i_k=(i_{k-1}-1)\alpha_k^N+1}^{i_{k-1}\alpha_k^N} \mathbb{1}_{i_1}^{(1)} \mathbb{1}_{i_2}^{(2)} \cdots \mathbb{1}_{i_k}^{(k)} < 1 \right\} \end{aligned}$$

By Chebyshev's inequality,

$$\begin{aligned} &\mathbb{P}\left(\sum_{i_1=1}^{\alpha_1^N} \sum_{i_2=(i_1-1)\alpha_2^N+1}^{i_1\alpha_2^N} \cdots \sum_{i_k=(i_{k-1}-1)\alpha_k^N+1}^{i_{k-1}\alpha_k^N} \mathbb{1}_{i_1}^{(1)} \mathbb{1}_{i_2}^{(2)} \cdots \mathbb{1}_{i_k}^{(k)} \geq 1\right) \\ &\leq \mathbb{E}\left(\sum_{i_1=1}^{\alpha_1^N} \sum_{i_2=(i_1-1)\alpha_2^N+1}^{i_1\alpha_2^N} \cdots \sum_{i_k=(i_{k-1}-1)\alpha_k^N+1}^{i_{k-1}\alpha_k^N} \mathbb{1}_{i_1}^{(1)} \mathbb{1}_{i_2}^{(2)} \cdots \mathbb{1}_{i_k}^{(k)}\right) \\ &= \alpha_1^N \cdots \alpha_k^N \mathbb{P}(E^{(1)} > Nx_1) \cdots \mathbb{P}(E^{(k)} > Nx_k) \\ &\leq \alpha_1^N \cdots \alpha_k^N \prod_{1 \leq i \leq k} \frac{J\sqrt{a_i}}{2x_i\sqrt{\pi N}} \exp\left(-\frac{Nx_i^2}{a_i J^2}\right) \\ &= \left(\prod_{1 \leq i \leq k} \frac{J\sqrt{a_i}}{2x_i\sqrt{\pi N}}\right) \exp\left\{N \sum_{1 \leq i \leq k} \left(\ln \alpha_i - \frac{x_i^2}{a_i J^2}\right)\right\}. \end{aligned}$$

Since

$$\sum_{1 \leq i \leq k} \frac{x_i^2}{a_i} > J^2 \sum_{1 \leq i \leq k} \ln \alpha_i,$$

the series

$$\sum_{N=1}^{\infty} \left(\prod_{1 \leq i \leq k} \frac{J\sqrt{a_i}}{2x_i\sqrt{\pi N}} \right) \exp \left\{ N \sum_{1 \leq i \leq k} \left(\ln \alpha_i - \frac{x_i^2}{a_i J^2} \right) \right\}$$

converges. Introducing the events

$$\mathcal{A}_N = \left\{ \{E_{i_1}^{(1)}, \dots, E_{i_k}^{(k)}\} \mid \sum_{i_1=1}^{\alpha_1^N} \sum_{i_2=(i_1-1)\alpha_2^N+1}^{i_1\alpha_2^N} \dots \sum_{i_k=(i_{k-1}-1)\alpha_k^N+1}^{i_{k-1}\alpha_k^N} \mathbb{1}_{i_1}^{(1)} \mathbb{1}_{i_2}^{(2)} \dots \mathbb{1}_{i_k}^{(k)} \geq 1 \right\}$$

we see again by the Borel-Cantelli Lemma, for almost all $\{E_{i_1}^{(1)}, \dots, E_{i_k}^{(k)}\}$ there exists a $v \in \mathbb{N}$ such that for all $N \geq v$, $\{E_{i_1}^{(1)}, \dots, E_{i_k}^{(k)}\} \in \mathcal{A}_N^C$ and hence $H_N(x_1, \dots, x_k) = 0$. Thus we have:

$$\begin{aligned} \limsup_N \frac{1}{N} \ln \bar{F}_N(x_1, \dots, x_n) &\leq \limsup_N \frac{1}{N} \ln H_N(x_1, \dots, x_k) \\ &= -\infty. \end{aligned}$$

□

3 The Variational Problem

We may re-write the partition function in (1.2) as

$$\mathcal{Z}_N(\beta) = 2^N \int_{\mathbb{R}^n} \exp \{-N\beta(x_1 + \dots + x_n)\} dF_N(x_1, \dots, x_n)$$

where $F_N(x_1, \dots, x_n)$ is given in (1.4). Using Varadhan's Lemma, we may evaluate $-\beta f(\beta)$ as follows:

$$\begin{aligned} -\beta f(\beta) &= \lim_{N \rightarrow \infty} \frac{1}{N} \{\ln \mathcal{Z}_N(\beta)\} \\ &= \ln 2 + \sup_{(x_1, \dots, x_n) \in \mathbb{R}^n} \{-\beta(x_1 + \dots + x_n) - I(x_1, \dots, x_n)\} \\ &= \ln 2 - \inf_{\vec{x} \in \Psi} \left\{ \sum_{i=1}^n \frac{x_i^2}{a_i J^2} + \beta x_i \right\} \\ &= \ln 2 + \frac{1}{4} \beta^2 J^2 - \frac{1}{J^2} \inf_{\vec{x} \in \Psi} \left\{ \sum_{i=1}^n \frac{1}{a_i} (x_i + \frac{1}{2} a_i \beta J^2)^2 \right\}. \end{aligned}$$

Performing the change of variables: $x_i = J y_i \sqrt{a_i}$, $\beta' = \frac{1}{2} \beta J$ and $\gamma_i = \ln \alpha_i$, the above expression becomes

$$= \ln 2 + \frac{1}{4} \beta^2 J^2 - \inf_{\vec{y} \in \Psi'} \left\{ \sum_{i=1}^n (y_i - \sqrt{a_i} \beta')^2 \right\}$$

where

$$\Psi' = \left\{ (y_1, \dots, y_n) \in \mathbb{R}^n \mid \sum_{i=1}^k y_i^2 \leq \sum_{i=1}^k \gamma_i, \text{ for all } 1 \leq k \leq n \right\}.$$

3.1 Evaluation of the infimum in \mathbb{R}^n

Define the numbers $B(j, k)$ for all $1 \leq j \leq k \leq n$:

$$B(j, k) := \sqrt{\frac{\gamma_j + \dots + \gamma_k}{a_j + \dots + a_k}}.$$

Let $m_0 := 0$ and recursively define the numbers m_i as

$$m_i := \inf \{m > m_{i-1} | B(m_{i-1} + 1, m) \leq B(m_{i-1} + 1, l), \text{ for all } m_{i-1} + 1 \leq l \leq n\}$$

terminating at the value K such that $m_K = n$. A crucial property of rational expressions like $B(j, k)$ is the following: if a, b, c and d are positive reals, then $\frac{a}{b} < \frac{c}{d}$ if and only if $\frac{a}{b} < \frac{a+c}{b+d}$. Define the sequence of inverse temperatures β_i ($i = 0, \dots, K+1$) by

$$\beta_i := B(m_{i-1} + 1, m_i), \quad i = 1, \dots, K.$$

and $\beta_0 := 0, \beta_{K+1} := +\infty$. Note that this sequence is increasing by the above property.

Lemma 3.1 *If $\beta_j < \beta' < \beta_{j+1}$ for some $0 \leq j \leq K$, then the infimum is attained at \vec{x} given by*

$$x_i = \begin{cases} \beta_l \sqrt{a_i}, & \text{if } i \in [m_{l-1} + 1, \dots, m_l] \text{ for some } 1 \leq l \leq j, \\ \beta' \sqrt{a_i}, & \text{if } i \in [m_j + 1, \dots, n], \end{cases}$$

for all $1 \leq i \leq n$.

PROOF: Let $p_i = \sqrt{a_i}$ for all $1 \leq i \leq n$. We will show that the point \vec{x} with coordinates given above is the point such that for all $y \in \Psi'$, $\|\vec{y} - \beta' \vec{p}\| \geq \|\vec{x} - \beta' \vec{p}\|$. First, let us note two trivial inequalities,

$$\sum_{l=1}^j \sum_{i=m_{l-1}+1}^{m_l} (y_i - \beta_l p_i)^2 \geq 0, \quad (3.1)$$

$$\sum_{i=m_j+1}^n (y_i - \beta' p_i)^2 \geq 0. \quad (3.2)$$

Note that for all $1 \leq l \leq j$, $\left(\frac{\beta'}{\beta_l} - 1\right) > 0$. By the Cauchy-Schwarz inequality we have, for all $1 \leq j' \leq j$,

$$\begin{aligned} \sum_{l=1}^{j'} \sum_{i=m_{l-1}+1}^{m_l} \beta_l p_i y_i &\leq \left(\sum_{i=1}^{m_{j'}} y_i^2 \right)^{1/2} \left(\sum_{l=1}^{j'} \sum_{i=m_{l-1}+1}^{m_l} \beta_l^2 p_i^2 \right)^{1/2} \\ &\leq \left(\sum_{i=1}^{m_{j'}} \gamma_i \right)^{1/2} \left(\sum_{l=1}^{j'} \beta_l^2 \sum_{i=m_{l-1}+1}^{m_l} p_i^2 \right)^{1/2}. \end{aligned}$$

Notice that $\sum_{l=1}^{j'} \beta_l^2 \sum_{i=m_{l-1}+1}^{m_l} p_i^2 = \sum_{l=1}^{j'} \sum_{i=m_{l-1}+1}^{m_l} \gamma_i = \sum_{i=1}^{m_{j'}} \gamma_i$ and so the above expression becomes

$$= \sum_{l=1}^{j'} \beta_l^2 \sum_{i=m_{l-1}+1}^{m_l} p_i^2.$$

Thus we have

$$\sum_{l=1}^{j'} \sum_{i=m_{l-1}+1}^{m_l} \beta_l p_i (\beta_l p_i - y_i) \geq 0$$

for all $1 \leq j' \leq j$ and $\vec{y} \in \Psi'$. Introducing the numbers $\left(\frac{\beta'}{\beta_l} - 1\right)$ into the sum, it is shown by a recursive argument (see Appendix) that

$$\sum_{l=1}^j \left(\frac{\beta'}{\beta_l} - 1\right) \sum_{i=m_{l-1}+1}^{m_l} \beta_l p_i (\beta_l p_i - y_i) \geq 0.$$

Multiplying the above inequality by 2 and re-writing we have

$$2 \sum_{l=1}^j \sum_{i=m_{l-1}+1}^{m_l} (\beta' p_i - \beta_l p_i)(\beta_l p_i - y_i) \geq 0. \quad (3.3)$$

Combining equations (3.1), (3.2) and (3.3) while noting that $\beta' p_i =: x_i$ (for $m_j + 1 \leq i \leq n$) and $\beta_l p_i =: x_i$ (for $1 \leq i \leq m_j$), we have

$$(\vec{y} - \vec{x}) \cdot (\vec{y} + \vec{x} - 2\beta' \vec{p}) \geq 0.$$

Re-writing gives

$$\|\vec{x} - \beta' \vec{p}\| \geq \|\vec{y} - \beta' \vec{p}\|,$$

for all $\vec{y} \in \Psi'$.

□

3.2 Expression for the Free energy

Applying the coordinates of our point of infimum to the expression for the free energy gives the required expressions. Recalling $\beta' := \frac{1}{2}\beta J$, $p_i = \sqrt{a_i}$ and $\gamma_i = \ln \alpha_i$ gives

Corollary 3.2 *The free energy is given by*

$$-\beta f(\beta) = \begin{cases} \ln 2 + \frac{1}{4}\beta^2 J^2, & \text{if } \beta < \frac{2}{J}\beta_1 \\ \sum_{i=m_j+1}^n \left(\ln \alpha_i + \frac{1}{4}\beta^2 J^2 a_i \right) + \beta J \sum_{l=1}^j \sqrt{\left(\sum_{i=m_{l-1}+1}^{m_l} a_i \right) \left(\sum_{i=m_{l-1}+1}^{m_l} \ln \alpha_i \right)}, & \text{if } \frac{2}{J}\beta_j < \beta < \frac{2}{J}\beta_{j+1} \\ \beta J \sum_{l=1}^K \sqrt{\left(\sum_{i=m_{l-1}+1}^{m_l} a_i \right) \left(\sum_{i=m_{l-1}+1}^{m_l} \ln \alpha_i \right)}, & \text{if } \frac{2}{J}\beta_K < \beta. \end{cases}$$

Applying $n = 2$ to the above expression yields the same answer as Derrida [3]. In this case the answer depends on whether $a_1/\ln \alpha_1 \geq a_2/\ln \alpha_2$. If $a_1/\ln \alpha_1 > a_2/\ln \alpha_2$, then

$$-\beta f(\beta) = \begin{cases} \ln 2 + \frac{J^2 \beta^2}{4}, & \text{if } \beta < \frac{2}{J} \sqrt{\frac{\ln \alpha_1}{a_1}} \\ \ln \alpha_2 + \frac{1}{4} a_2 \beta^2 J^2 + \beta J \sqrt{a_1 \ln \alpha_1}, & \text{if } \frac{2}{J} \sqrt{\frac{\ln \alpha_1}{a_1}} < \beta < \frac{2}{J} \sqrt{\frac{\ln \alpha_2}{a_2}} \\ \beta J \sqrt{a_1 \ln \alpha_1} + \beta J \sqrt{a_2 \ln \alpha_2}, & \text{if } \frac{2}{J} \sqrt{\frac{\ln \alpha_2}{a_2}} < \beta. \end{cases}$$

Otherwise,

$$-\beta f(\beta) = \begin{cases} \ln 2 + \frac{J^2 \beta^2}{4}, & \text{if } \beta < \frac{2\sqrt{\ln 2}}{J} \\ \beta J \sqrt{\ln 2}, & \text{if } \beta > \frac{2\sqrt{\ln 2}}{J}. \end{cases}$$

It is an easy exercise to see the solutions also concur for cases A and B in Derrida and Gardner [4]. Capocaccia et. al. [1] approach to the variational problem contains a few minor flaws which are easily seen by setting $n = 2$ in their final expression for the free energy. In this case, their result does not distinguish between the above two cases and their critical temperature is incorrect.

Appendix

Lemma A *Let $x_1, x_2, \dots, x_n > 0$ and $\{y_i\}_{i=1}^n$ be a sequence of reals. Let $G_m := \sum_{i=1}^m y_i$ be such that $G_m \geq 0$ for all $1 \leq m \leq n$. Then*

$$F(n) := \sum_{i=1}^n x_i y_i \geq 0.$$

PROOF: Let us define $G_0 = 0$. Notice that $y_i = G_i - G_{i-1}$ for all $1 \leq i \leq n$. Then

$$\begin{aligned} F(n) &= \sum_{i=1}^n x_i (G_i - G_{i-1}) \\ &= x_n G(n) + \sum_{i=1}^n (x_i - x_{i-1}) G(i) \\ &\geq 0, \end{aligned}$$

since $x_i - x_{i-1} > 0$ for all i .

□

References

- [1] M. Cassandro D. Capocaccia and P. Picco. On the Existence of Thermodynamics for the Generalized Random Energy Model. *J. Stat. Phys.*, 46(3/4):493–505, 1987.
- [2] B. Derrida. *Phys. Rev. Lett.*, 45:79, 1980.
- [3] B. Derrida. A generalization of the Random Energy Model which includes correlations between energies. *J. Physique Lett.*, 46:401–407, 1985.
- [4] B. Derrida and E. Gardner. Solution of the generalized random energy model. *J. Phys. C: Solid State Phys.*, 19:2253–2274, 1986.
- [5] T.C. Dorlas and J.R. Wedagedera. Large deviation and the random energy model. *International Journal of Modern Physics B*, 15(1):1–15, 2001.
- [6] H. McKean. *Stochastic Integrals*. Acad. Press, 1969.