

Title	Quantization of Flag Manifolds and their Supersymmetric Extensions
Creators	Murray, Seán and Sämann, Christian
Date	2006
Citation	Murray, Seán and Sämann, Christian (2006) Quantization of Flag Manifolds and their Supersymmetric Extensions. (Preprint)
URL	https://dair.dias.ie/id/eprint/489/
DOI	DIAS-STP-06-21

Quantization of Flag Manifolds and their Supersymmetric Extensions

Seán Murray^{1,2} and Christian Sämann¹

¹ *School of Theoretical Physics
Dublin Institute for Advanced Studies
10 Burlington Road, Dublin 4, Ireland*

² *Department of Mathematical Physics
NUI Maynooth
Maynooth, Co. Kildare, Ireland*

Email: smury, csamann@stp.dias.ie

Abstract

We first review the description of flag manifolds in terms of Plücker coordinates and coherent states. Using this description, we construct fuzzy versions of the algebra of functions on these spaces in both operatorial and star product language. Our main focus is here on flag manifolds appearing in the double fibrations underlying the most common twistor correspondences. After extending the Plücker description to certain supersymmetric cases, we also obtain the appropriate deformed algebra of functions on a number of fuzzy flag supermanifolds. In particular, fuzzy versions of Calabi-Yau supermanifolds are found.

Contents

1. Introduction and results	2
2. Plücker coordinates and the geometry of flag manifolds	4
2.1. Flag manifolds of $U(4)$	4
2.2. Description of $\mathbb{C}P^3$	5
2.3. Plücker coordinates and projectors describing irreducible flag manifolds	6
2.4. The description of reducible flag manifolds	9
2.5. Geometric structures on the flag manifolds	11
2.6. Spherical functions on the flag manifolds	14
3. Flag manifolds and coherent states	15
3.1. Representations of $SU(n)$ and coherent states	15
3.2. Examples for $SU(3)$	17
3.3. The flags in \mathbb{C}^4	18
4. Fuzzification of the flag manifolds	19
4.1. Fuzzification	19
4.2. The fuzzy complex projective space $\mathbb{C}P_F^3$	20
4.3. The fuzzy Grassmannian $G_{2;4}^F$	23
4.4. The fuzzy dual complex projective space $F_{3;4}^F$	25
4.5. The fuzzy reducible flag manifold $F_{12;4}^F$	27
4.6. The fuzzy reducible flag manifolds $F_{23;4}^F$, $F_{13;4}^F$ and $F_{123;4}^F$	28
4.7. Continuous limits of the fuzzy flag manifolds	29
5. Super Plücker embeddings and flag supermanifolds	30
5.1. Flag supermanifolds of $U(4 n)$	30
5.2. The Plücker and super Plücker embeddings	32
5.3. Plücker coordinates and projector description of irreducible flag supermanifolds	34
5.4. The reducible flag supermanifolds	37
5.5. Geometric structures on the flag supermanifolds	37
6. Fuzzy flag supermanifolds	38
6.1. Supercoherent states	38
6.2. The fuzzy complex projective superspace $\mathbb{C}P_F^{3 4}$	39
6.3. The remaining fuzzy Grassmannian supermanifolds	41
6.4. The fuzzy reducible flag supermanifolds	41
6.5. Fuzzy Calabi-Yau supermanifolds	42
Acknowledgements	42
Appendices	
A. Supermathematics, conventions and definitions	42
B. Representations of $\mathfrak{su}(4)$ and $\mathfrak{u}(4 4)$	45

1. Introduction and results

Quite often in physics, approximation methods like perturbation theory are necessary for explicit computations. In particular, non-perturbative methods, which typically involve the reduction of the field theory to a model with a finite number of degrees of freedom, are required to access the physics of field theories in the strong coupling regime. The standard method of this type is lattice field theory. It has been very successful in the study of confinement in quantum chromodynamics and for non-perturbative regularization of quantum field theories.

Lattice discretizations do have some disadvantages, however. They do not retain the symmetries of the exact theory except in some rough sense. By limiting the couplings to nearest neighbor, the topology and differential geometry of the underlying manifolds are treated only indirectly. Furthermore, the description of fermions in this context leads to the well-known fermion doubling problem.

Fortunately, the lattice is not the only method of reducing a field theory to a finite number of degrees of freedom. An alternative is what has become known as the fuzzy approach [1]-[10], see [11] for a detailed review. The basic idea is here to take a classical phase space of finite volume, quantize it and thus obtain a space carrying a function algebra with a finite number of degrees of freedom. There are certain limitations to this approach, such as the even dimensionality of the parent manifold, that can be avoided when the phase space is a co-adjoint orbit of a Lie group. The functions on the resultant fuzzy spaces are described by linear operators on irreducible representations of the group. The simplest such example is the two sphere S^2 , with the resulting phase space known as the fuzzy sphere [1]. Field theory models on the fuzzy sphere then possess only a finite number of modes. The simplest such field theory with ϕ^4 interaction was proposed in [3].

There are other reasons to consider fuzzy spaces. They lead to matrix models, which have seen much interest by string theorists especially in describing D-branes: When considering D-branes on group manifolds [12], turning on background fields can render the target space geometry fuzzy [13]. Similarly, a system of D0-branes in a nontrivial background can form the fuzzy sphere [14]; see also [15].

The spaces we chose for deformation play a prominent rôle in various geometrical areas. Flag manifolds, i.e. the spaces of sequences of nested subvector spaces in a given vector space, are generalizations of Grassmannians (and thus of complex projective spaces) and serve as non-trivial examples in algebraic geometry. They are special cases of coset spaces, and in particular coset superspaces received growing attention recently [16, 17]. Moreover, flag manifolds arise naturally in the theory of characteristic classes of vector bundles, in representation theory, in mirror symmetry and in twistor theory. It is therefore clear that studying fuzzy versions of flag manifolds may lead to a deeper understanding of both differential and algebraic geometry on fuzzy spaces. Although the results presented in this paper generalize to arbitrary flag manifolds, we will restrict our attention to those which appear naturally in the double fibrations of twistor theory described e.g. in [18]. A few examples of quantized flag manifolds which have a noncommutative algebra of functions with finitely many degrees of freedom have already been studied in a different form in [15, 19].

We start our discussion by giving a detailed description of flag manifolds in terms of

Plücker coordinates and the geometric structures on these spaces. The latter is induced from a canonical embedding of the flag manifolds into Euclidean space. We continue with the description of the correspondence between flag manifolds and coherent states in various representations of the Lie group $SU(n)$. In particular, a relationship between the patches covering a flag manifold and dominant weight states in the corresponding representation is established.

With the appropriate representations found in the coherent state picture together with the Plücker description, the discussion of fuzzy flag manifolds is rather straightforward. We present the matrix algebras corresponding to the algebra of functions on these spaces together with the equivalent star product picture. The latter is used to translate derivatives, which contain information about the geometry of the flag manifolds, into the operator language. In particular, the Laplacian turns into the second order Casimir operator in the considered representation. It is also shown that the constructed matrix algebras converge towards the algebra of continuous functions on flag manifolds in the limit of infinite-dimensional representations.

To prepare the fuzzification of flag supermanifolds, we develop the superanalogue to the Plücker embedding, which is novel, as far as we know. Also, the embedding of these flag supermanifolds into Euclidean superspaces is discussed. A relation between supercoherent states and points on flag supermanifolds is found, which is closely related to the corresponding picture in the case of ordinary flag manifolds. The fuzzification can then be obtained in a rather straightforward way. We give a series of matrix algebras, which approximate functions on the flag supermanifolds and present the equivalent star product formulation. All derivatives can again be translated into the operator language and encode geometric information about the spaces.

The results we obtain may find several applications. First, it is desirable to see whether the Penrose-Ward transform (see e.g. [20] for a review) can be carried over to an analogous construction built on a double fibration of fuzzy spaces. This, however, would demand a clearer understanding of the various gauge theories (i.e. holomorphic Chern-Simons and Yang-Mills theory) on the involved fuzzy geometries together with an explicit notion of holomorphic vector bundles over fuzzy spaces, see [21] for progress in this direction. Second, the Graßmannian $G_{2,4}$ is the conformal compactification of complex Minkowski space and after imposing reality conditions, one arrives at the compactified form of four-dimensional space-times with all possible signatures. Fuzzy versions of these spaces would certainly be very useful; unfortunately, it is not clear, how to impose the corresponding reality conditions in the fuzzy case. The main purpose of constructing fuzzy flag manifolds and in particular their supersymmetric counterparts is, however, to have at hand fuzzy versions of Calabi-Yau supermanifolds. These spaces, as e.g. the fuzzy version of the complex projective superspace $\mathbb{C}P^{3|4}$ discussed in this paper, might be used for the construction of first examples of interacting supersymmetric field theories on fuzzy spaces that can be simulated numerically. Furthermore, there is a conjectured mirror symmetry [22] between two of the flag supermanifolds we describe in this paper ($\mathbb{C}P^{3|4}$ and $F_{(1|0)(3|3);4|3}$), and trying to understand this mirror symmetry in terms of fuzzy spaces seems very promising.

2. Plücker coordinates and the geometry of flag manifolds

2.1. Flag manifolds of $U(4)$

Consider the vector space \mathbb{C}^n . A *flag* $f_{k_1 \dots k_r; n}$ in \mathbb{C}^n is a sequence of nested vector subspaces $V_{k_1} \subsetneq \dots \subsetneq V_{k_r} \subset \mathbb{C}^n$ such that $\dim_{\mathbb{C}} V_j = j$. A *flag manifold* $F_{k_1 \dots k_r; n}$ is the set of all flags $f_{k_1 \dots k_r; n}$.

The simplest example of a flag manifold is $F_{1; n}$, which is the complex projective space $\mathbb{C}P^{n-1}$. Furthermore, the Grassmannian $G_{k; n}$, the space of k -dimensional vector subspaces of \mathbb{C}^n , is the flag manifold $F_{k; n}$. A flag $f_{k; n} = V_k$ is obviously invariant under the subgroup $H = U(n-k) \times U(k) \subset U(n)$, as the elements of $U(n-k)$ do not change vectors in V_k , while the elements of $U(k)$ are just the unitary maps $V_k \rightarrow V_k$. Therefore, the group H defines (maximal) equivalence classes of flags in $U(n)$ and we can write $F_{k; n} = U(n)/H$. This can be generalized to

$$\begin{aligned} F_{k_1 \dots k_r; n} &= U(n)/(U(n-k_r) \times U(k_r - k_{r-1}) \dots \times U(k_1)) \\ &= SU(n)/S(U(n-k_r) \times U(k_r - k_{r-1}) \dots \times U(k_1)) , \end{aligned} \quad (2.1)$$

and thus the dimension of this flag manifold is $n^2 - (n-k_r)^2 - (k_r - k_{r-1})^2 - \dots - (k_1)^2$. Note that the above equation cannot be used as a defining relation, as the embedding of the subgroup factored out is not specified. The flag manifolds of $SU(4)$ can also be obtained as coset spaces of $SL(4, \mathbb{C})$, the complexification of $SU(4)$, see e.g. [23]. Here, one factors out the group of certain upper block triangular matrices and from this complexified description it follows that flag manifolds are complex manifolds. They are in fact Kähler manifolds and we will construct their Kähler structure explicitly later on. We will also see that flag manifolds are adjoint orbits $\{g\mathcal{P}g^{-1} | g \in SU(4)\}$ of certain projectors \mathcal{P} and therefore carry a natural symplectic structure. Furthermore, a flag manifold is a *homogeneous space*.

The flag manifolds of $U(n)$ split naturally into *irreducible* and *reducible* ones, where the irreducible flag manifolds are the Grassmannians $F_{k_1; n} = G_{k_1; n}$. In their case, the compact subgroup H consists of two factors. These flag manifolds form *hermitian symmetric spaces*, i.e. the commutators of two elements of $\mathfrak{u}(n)/(\mathfrak{u}(n-k_1) \times \mathfrak{u}(k_1))$ is an element of $\mathfrak{u}(n-k_1) \times \mathfrak{u}(k_1)$.

In the following, we will be exclusively interested¹ in flag manifolds of $U(4)$, which naturally appear in the double fibrations underlying the most important twistor correspondences, see e.g. [18]. These fibrations are obtained by truncating the flags in an obvious manner, e.g. there is a projection $F_{12;4} \rightarrow F_{2;4}$. All the twistor double fibrations are included in the following diagram:

$$\begin{array}{ccccc} & & F_{23;4} & & \\ & \swarrow & \uparrow & \searrow & \\ F_{2;4} & & & & F_{3;4} \\ & \swarrow & \uparrow & \searrow & \\ & & F_{123;4} & & \\ & \swarrow & \downarrow & \searrow & \\ F_{12;4} & & & & F_{13;4} \\ & \swarrow & \downarrow & \searrow & \\ & & F_{1;4} & & \end{array} \quad (2.2)$$

¹Nevertheless, all of our discussion trivially translates into the case of flag manifolds of $U(n)$.

In the twistor context, the space $F_{2;4}$ is the conformal compactification of complexified Minkowski space \hat{M} and the spaces $F_{1;4}$, $F_{3;4}$, $F_{13;4}$ are the spaces of self-dual null planes in \hat{M} (twistor space), anti-self-dual null planes in \hat{M} (dual twistor space) and null geodesics in \hat{M} (a thickening of which is the ambitwistor space), respectively. One can also consider affine (non-compact) subspaces of all the above spaces and the corresponding double fibrations. For more details on this point, see [20].

The dimensions of the involved spaces are easily calculated from the formula given below the defining equation (2.1). The minimal number of patches in a covering of the flag manifolds can be calculated inductively in the following way. The minimal number of patches covering all of $G_{k;n}$ is $\binom{n}{k}$; in particular, we have n as the minimal number of patches for $\mathbb{C}P^{n-1}$. The number of patches needed for a flag manifold is then obtained by multiplying the patches of the contained subflags. For example, to cover $F_{12;4}$, one needs at least 6 for $F_{2;4}$ times 2 for $F_{1;2}$ equals 12 patches. We summarize the results of these calculations in the following table:

Flag manifold	$F_{1;4}$	$F_{2;4}$	$F_{3;4}$	$F_{12;4}$	$F_{13;4}$	$F_{23;4}$	$F_{123;4}$
complex dimension	3	4	3	5	5	5	6
minimal # patches	4	6	4	12	12	12	24

2.2. Description of $\mathbb{C}P^3$

There are various aspects of the classical description of flag manifolds, that we will use for their fuzzification. In particular, we need a description in terms of homogeneous coordinates, a description in terms of projectors and the link between both of them. We will first discuss the simple example of $F_{1;4} = \mathbb{C}P^3 = \text{U}(4)/(\text{U}(3) \times \text{U}(1))$ in detail before going over to the more complicated spaces.

A normalized vector in \mathbb{C}^4 clearly spans a one-dimensional vector subspace of \mathbb{C}^4 and thus corresponds to a flag $f_{1;4}$. There is, however, a redundancy in the total phase of the vector, which needs to be factored out. One is thus naturally led to consider the generalized Hopf fibration defined by the short exact sequence

$$1 \longrightarrow \text{U}(1) \longrightarrow S^{2n-1} \longrightarrow \mathbb{C}P^{n-1} \longrightarrow 1 \quad (2.3)$$

for the case $n = 4$. In coordinates a^i on \mathbb{C}^4 , the projection down to S^7 amounts to imposing the condition $a^i \bar{a}^j \delta_{ij} = 1$ and the subsequent projection down to $\mathbb{C}P^3$ is performed by considering the auxiliary coordinates

$$x_{1;4}^a := \bar{a}^i \lambda_{ij}^a a^j, \quad (2.4)$$

where λ_{ij}^a , $a = 1, \dots, 15$ are the Gell-Mann matrices² of $\text{SU}(4)$. These coordinates describe an embedding of $\mathbb{C}P^3$ in \mathbb{R}^{15} . Note that we factored out only the $\text{U}(1)$ (internal) part from the invariance group of the flag $f_{1;4}$ acting non-trivially on the one-dimensional subspace of

²We shall adopt the following convention throughout:

$$\text{tr}(\lambda^a \lambda^b) = \delta^{ab}, \quad [\lambda^a, \lambda^b] = \sqrt{2} i f^{ab}{}_c \lambda^c.$$

\mathbb{C}^4 spanned by the vector a . The remaining $U(3)$ (external) part acts orthogonally to this vector and therefore leaves it invariant. In an equivalent construction [21], the action of this external group appears more explicitly.

The homogeneous coordinates a^i are a special case of the so-called Plücker coordinates, which we will discuss in the next section. Before, however, let us give a second description of $\mathbb{C}P^3$ in terms of projectors, see e.g. [6].

A projector \mathcal{P} is a hermitian 4×4 matrix satisfying $\mathcal{P}^2 = \mathcal{P}$. The *rank* of the projector \mathcal{P} , $\text{tr}(\mathcal{P})$, is equal to the dimension of the subspace it projects onto. It is therefore evident that every point on an irreducible flag manifold $F_{k;4}$ corresponds to a rank k projector $\mathcal{P}_{k;4}(x)$; in particular, $\mathbb{C}P^3$ is isomorphic to the space of rank one projectors $\mathcal{P}_{1;4}(x)$.

The space of projectors acting on \mathbb{C}^4 is spanned by the Gell-Mann matrices of $SU(4)$ and the identity. We can write

$$\mathcal{P} = x^{\hat{a}} \lambda_{\hat{a}} = x^0 \lambda_0 + x^a \lambda_a, \quad (2.5)$$

where $\hat{a} = 0, \dots, 15$ and $a = 1, \dots, 15$. We use $\lambda_0 = \mathbb{1}/\sqrt{4}$, which implies that $x^0 = \text{tr}(\mathcal{P})/\sqrt{4}$ and

$$\lambda_a \lambda_b = \frac{\delta_{ab}}{\sqrt{4}} \lambda_0 + \frac{1}{\sqrt{2}} (d_{ab}{}^c + i f_{ab}{}^c) \lambda_c, \quad (2.6)$$

where $d_{ab}{}^c$ and $f_{ab}{}^c$ are the (traceless) symmetric invariant tensor and the structure constants of $SU(4)$, respectively. Recall that the Lie algebra indices are raised and lowered with the Killing metric δ_{ab} .

As stated above, the irreducible flag manifolds $F_{k;4}$ correspond to the space of projector $\mathcal{P}_{k;4}$ of rank k and the condition $(\mathcal{P}_{k;4})^2 = \mathcal{P}_{k;4}$ defines a set of quadratic constraints, embedding the flag manifolds in \mathbb{R}^{16} (or \mathbb{R}^{15} , if one considers $x_{k;4}^0$ already fixed by the condition on the trace of $\mathcal{P}_{k;4}$). Explicitly, they read as

$$x_{k;4}^a x_{k;4}^a = \frac{4k - k^2}{4} \quad \text{and} \quad x_{k;4}^a x_{k;4}^b d_{ab}{}^c = \sqrt{2} \frac{4 - 2k}{4} x_{k;4}^c. \quad (2.7)$$

Given a projector $\mathcal{P}_{k;4}^0$ of rank k , all of the space $F_{k;4}$ is obtained by its orbit $g \mathcal{P}_{k;4}^0 g^{-1}$, $g \in U(4)$. However, two elements g and g' related by $g = g' h$, where $h \in H = U(4-k) \times U(k)$, will rotate to the same element $g \mathcal{P}_{k;4}^0 g^{-1} = g' \mathcal{P}_{k;4}^0 g'^{-1}$. This simply reflects the definition (2.1) of $F_{k;4}$ as a coset space.

There is evidently a relation between $F_{1;4}$ and $F_{3;4}$ since the coordinates $x_{1;4}^a$ of a projector $\mathcal{P}_{1;4}$ yield the coordinates of a projector $\mathcal{P}_{3;4}$ by $x_{3;4}^a = -x_{1;4}^a$, as one easily checks using (2.7). Furthermore, the coordinates $x_{2;4}^a$ of a projector $\mathcal{P}_{2;4}$ yield a second projector $\check{\mathcal{P}}_{2;4}$ with coordinates $\check{x}_{2;4}^a = -x_{2;4}^a$. The meaning of these dualities will become clear in the next section.

For $\mathbb{C}P^3 = F_{1;4}$, the projector is obtained by extending the definition (2.4) of the auxiliary coordinates to $x_{1;4}^{\hat{a}} = \bar{a}^i \lambda_{ij}^{\hat{a}} a^j$, which satisfy the constraints (2.7). Due to $\lambda_{ij}^{\hat{a}} \lambda_{kl}^{\hat{a}} = \delta_{il} \delta_{kj}$, the resulting projector is then explicitly given by the matrix $\mathcal{P}_{1;4} = a \bar{a}^T$ and one easily verifies $(\mathcal{P}_{1;4})^2 = \mathcal{P}_{1;4}$.

2.3. Plücker coordinates and projectors describing irreducible flag manifolds

To define a two-plane in \mathbb{C}^4 , we can use two normalized vectors $a, b \in \mathbb{C}^4$ antisymmetrized to $A_2 := a \wedge b = \frac{1}{2} (a \otimes b - b \otimes a) = (A_2^{ij}) := a^{[i} b^{j]}$. As one easily observes, the antisym-

metrization projects on the mutually orthogonal components of a and b . The A_2^{ij} are so-called *Plücker coordinates* on $F_{2;4} = \text{U}(4)/(\text{U}(2) \times \text{U}(2))$ and satisfy by construction the identity

$$\varepsilon_{ijkl} A_2^{ij} A_2^{kl} = 0. \quad (2.8)$$

As there are six projective Plücker coordinates, we learn that the Grassmannian $F_{2;4}$ is a quadric in $\mathbb{C}P^5$, the so-called *Klein quadric*, see e.g. [24]. Equation (2.8) is an example of the *Plücker relations*, which describe an embedding of a Grassmannian $G_{k;n}$ in $\mathbb{P}(\Lambda^k \mathbb{C}^n)$. Although the Plücker relation is straightforward in the present case of $F_{2;4} = G_{2;4}$, we will need more nontrivial such relations when discussing flag supermanifold and we will present a more explicit discussion in section 5.2. There are certainly other approaches to coordinatizing flag manifolds; see e.g. [23] for “Bruhat coordinates” and more background material on flag manifolds.

Let us now consider the space of hyperplanes in \mathbb{C}^4 , i.e. $F_{3;4} = \text{U}(4)/(\text{U}(1) \times \text{U}(3))$. Analogous to the case of the two-plane, a three-plane is spanned by three antisymmetrized vectors $a \wedge b \wedge c$, which are naturally dual to a single vector $\vec{d} = (\vec{d}_i) = (\varepsilon_{ijkl} a^j b^k c^l)$, which in turn spans the orthogonal complement to the hyperplane. However, the non-dualized picture will be useful later on and therefore let us also introduce the Plücker coordinates $A_3^{ijk} = a^{[i} b^j c^k]$.

We can contract these new Plücker coordinates by tensor products of the Gell-Mann matrices, which yields auxiliary coordinates describing an embedding of the Grassmannians in Euclidean space. In the case of $F_{2;4}$, we have

$$x_{2;4}^{\hat{a}\hat{b}} = \bar{A}_2^{i_1 i_2} (\lambda^{\hat{a}} \wedge \lambda^{\hat{b}})_{i_1 i_2, j_1 j_2} A_2^{j_1 j_2} \quad (2.9)$$

with the antisymmetrized tensor product \wedge defined in components as

$$(A \wedge B)_{ij;kl} = \frac{1}{4} (A_{ik} B_{jl} - A_{jk} B_{il} - A_{il} B_{jk} + A_{jl} B_{ik}). \quad (2.10)$$

The choice of this contraction, which again factors out a phase, will become obvious after discussing the description of $F_{2;4}$ in terms of projectors. Note that $x_{2;4}^{\hat{a}\hat{b}}$ is *symmetric* in its indices. The above contraction is in agreement with the generalized Hopf fibration³

$$1 \longrightarrow \text{U}(2) \longrightarrow S^7 \times S^5 \longrightarrow F_{2;4} = G_{2;4} \longrightarrow 1. \quad (2.11)$$

As before, a normalized complex vector in \mathbb{C}^4 defines a point on S^7 , and we choose a to be this point. In the combination $A_2 = a \wedge b$, the component of b parallel to a vanishes trivially, and thus the relevant component of b is a point on S^5 . Factoring out the internal $\text{U}(2)$ which describes rotations in the plane $a \wedge b$, one obtains $G_{2;4}$. The other $\text{U}(2)$ factor is again trivially factored out, since it does not affect A_2^{ij} . To see that the contraction (2.9) indeed factors out an $\text{U}(2)$, note that the action

$$\begin{pmatrix} a^1 & \dots & a^4 \\ b^1 & \dots & b^4 \end{pmatrix} \mapsto g \begin{pmatrix} a^1 & \dots & a^4 \\ b^1 & \dots & b^4 \end{pmatrix} \quad \text{with } g \in \text{U}(2), \quad (2.12)$$

³After imposing a certain reality condition, this fibration reduces naturally to

$$1 \rightarrow S^1 \times S^1 \rightarrow S^3 \times S^3 \rightarrow S^2 \times S^2 \cong G_{2;4}^{\mathbb{R}} \rightarrow 1.$$

leaves invariant both \bar{A}_2^{ij} and A_2^{kl} up to a phase, and therefore (2.9) is indeed invariant. Correspondingly, one can discuss the isotropy groups for all the flag manifolds we construct in the following. We refrain from doing this, but present a more detailed discussion in the quantized picture.

In the case of $F_{3;4}$, we choose the auxiliary coordinates

$$x_{3;4}^{\hat{a}\hat{b}\hat{c}} = \bar{A}_3^{i_1 i_2 i_3} (\lambda^{\hat{a}} \wedge \lambda^{\hat{b}} \wedge \lambda^{\hat{c}})_{i_1 i_2 i_3, j_1 j_2 j_3} A_3^{j_1 j_2 j_3} . \quad (2.13)$$

In the dual picture, this corresponds to

$$\tilde{x}_{3;4}^{\hat{a}} = \bar{d}^{\tilde{k}} \tilde{\lambda}_{kl}^{\hat{a}} \tilde{d}^{\tilde{l}} \quad \text{with} \quad \tilde{\lambda}_{kl}^{\hat{a}} \sim \varepsilon_{k i_1 i_2 i_3} \varepsilon_{l j_1 j_2 j_3} (\lambda^{\hat{b}} \wedge \lambda^{\hat{c}} \wedge \lambda^{\hat{d}})_{i_1 i_2 i_3, j_1 j_2 j_3} , \quad (2.14)$$

and the implied map of the Lie algebra indices $(\hat{b}\hat{c}\hat{d}) \mapsto \hat{a}$ can easily be calculated. This contraction corresponds to the generalized Hopf fibration

$$1 \longrightarrow \mathrm{U}(3) \longrightarrow S^7 \times S^5 \times S^3 \longrightarrow F_{3;4} \longrightarrow 1 . \quad (2.15)$$

Although we already gave a description of the Graßmannians $F_{k;4}$ in terms of projectors in the previous section, it will be more convenient to switch to certain rank 1 projectors $\mathcal{P}_{k;4}$ acting on the representation spaces of the $\mathbf{6}$ and $\bar{\mathbf{4}}$ of $\mathfrak{u}(4)$ in the cases $F_{2;4}$ and $F_{3;4}$, respectively. This can be done in three equivalent ways. In the first one, one chooses two or three rank one projectors and antisymmetrizes them

$$\mathcal{P}_{2;4} = 2\mathcal{P}^1 \wedge \mathcal{P}^2 \quad \text{and} \quad \mathcal{P}_{3;4} = 3\mathcal{P}^1 \wedge \mathcal{P}^2 \wedge \mathcal{P}^3 , \quad (2.16)$$

where $\mathcal{P}^r = x_r^{\hat{a}} \lambda_{\hat{a}}$ are some rank one projectors. Besides the usual conditions (2.7) on rank one projectors, additional conditions between the coordinate vectors x_r and x_s arise to guarantee that $\mathcal{P}_{2;4}$ and $\mathcal{P}_{3;4}$ are projectors. These conditions state, e.g. for $\mathcal{P}_{2;4}$ that $\mathcal{P}^1 + \mathcal{P}^2$ is again a projector, which amounts to $\mathcal{P}^1 \mathcal{P}^2 + \mathcal{P}^2 \mathcal{P}^1 = 0$. In terms of coordinates, the first projector is constructed from a complex vector a^i by $x_{1;4}^a = \bar{a}^i \lambda_{ij}^a a^j$, while the second one is constructed from an orthonormalized vector b_{\perp}^i with $b_{\perp}^i \sim b^i - (\bar{a}^j b^j) a^i$ by $x_{2;4}^a = \bar{b}_{\perp}^i \lambda_{ij}^a b_{\perp}^j$. The sum of these two rank one projectors will automatically yield a rank two projector.

Alternatively, we can also antisymmetrize our previous rank two and rank three projectors $\mathcal{P}_{2;4}$ and $\mathcal{P}_{3;4}$

$$\mathcal{P}'_{2;4} = \mathcal{P}_{2;4} \wedge \mathcal{P}_{2;4} \quad \text{and} \quad \mathcal{P}'_{3;4} = \mathcal{P}_{3;4} \wedge \mathcal{P}_{3;4} \wedge \mathcal{P}_{3;4} , \quad (2.17)$$

as discussed in [8]. Note that both approaches are equivalent, the latter, however, is slightly more economical in the use of parameters. Furthermore, due to the formula

$$\mathrm{tr}(A \wedge B) = \frac{1}{2}(\mathrm{tr}(A) \mathrm{tr}(B) - \mathrm{tr}(AB)) , \quad (2.18)$$

which is easily verified using (2.10), and a similar one for the antisymmetrization of three projectors, all of the above projectors have unit trace and therefore indeed rank 1.

Here, we choose to work in the first approach, embedding the Graßmannians in the space of symmetrized products of vectors in \mathbb{R}^{16} . This will eventually lead to simpler expressions

⁴Recall that these representations carry two and three antisymmetrized indices of the fundamental of $\mathfrak{u}(4)$, respectively.

for the star product on all the Graßmannians. It is also linked to the contractions we obtained from the various Hopf fibrations in the previous section. Let us first introduce the shorthand notation

$$\lambda^{\hat{a}_1 \dots \hat{a}_n} = \lambda^{\hat{a}_1} \wedge \dots \wedge \lambda^{\hat{a}_n} . \quad (2.19)$$

Recall that $\lambda^{\hat{a}_1 \dots \hat{a}_n}$ turns out to be totally symmetric in its indices. Using this notation, we can easily define the appropriate projectors in the $\mathbf{6}$ and $\bar{\mathbf{4}}$ of $\mathfrak{u}(4)$ as

$$\begin{aligned} \mathcal{P}_{2;4} &= 2x_{2;4}^{(\hat{a}\hat{b})} \lambda^{\hat{a}\hat{b}} \quad \text{with} \quad x_{2;4}^{(\hat{a}\hat{b})} = \bar{a}^{i_1} \bar{b}^{i_2} (\lambda^{\hat{a}\hat{b}})_{i_1 i_2, j_1 j_2} a^{j_1} b^{j_2} , \\ \mathcal{P}_{3;4} &= 3x_{3;4}^{(\hat{a}\hat{b}\hat{c})} \lambda^{\hat{a}\hat{b}\hat{c}} \quad \text{with} \quad x_{3;4}^{(\hat{a}\hat{b}\hat{c})} = \bar{a}^{i_1} \bar{b}^{i_2} \bar{c}^{i_3} (\lambda^{\hat{a}\hat{b}\hat{c}})_{i_1 i_2 i_3, j_1 j_2 j_3} a^{j_1} b^{j_2} c^{j_3} . \end{aligned} \quad (2.20)$$

Here, the subspaces are spanned by complex vectors a, b and a, b, c , respectively, and $x_{2;4}^{(\hat{a}\hat{b})}$ and $x_{3;4}^{(\hat{a}\hat{b}\hat{c})}$ describe embeddings of $F_{2;4}$ and $F_{3;4}$ in $\mathbb{R}^{16 \cdot 17/2-1}$ and $\mathbb{R}^{16 \cdot 17 \cdot 18/(2 \cdot 3)-1}$, respectively. Note that the coordinates $x_{2;4}^{00}$ and $x_{3;4}^{000}$ are fixed by the ranks of the projectors $\mathcal{P}_{2;4}$ and $\mathcal{P}_{3;4}$.

To check that these operators are indeed projectors, one uses identities like

$$(A \wedge B)(C \wedge D) = \frac{1}{2}((AC \wedge BD) + (AD \wedge BC)) \quad (2.21)$$

yielding the Fierz identities discussed in appendix B. For example, $\mathcal{P}_{2;4}$ can be shown to read as

$$(\mathcal{P}_{2;4})_{ij;kl} = a^{[i} b^{j]} \bar{a}^{[k} \bar{b}^{l]} , \quad (2.22)$$

where we have chosen a and b orthogonal to each other. It then follows immediately that

$$(\mathcal{P}_{2;4})_{ij;kl} (\mathcal{P}_{2;4})_{kl;mn} = (\mathcal{P}_{2;4})_{ij;mn} . \quad (2.23)$$

Note that the naïve contraction to obtain the auxiliary coordinates for $F_{2;4}$

$$x_{2;4}^{\hat{a}} = \bar{A}_2^{ij} (\lambda^{\hat{a}} \wedge \mathbb{1})_{ij,kl} A_2^{kl} \quad (2.24)$$

does not yield a projector since $x_{2;4}^{\hat{a}} (\lambda^{\hat{a}} \wedge \mathbb{1})_{ij,kl}$ is not idempquadratic.

2.4. The description of reducible flag manifolds

The construction of the Plücker coordinates for the reducible flag manifolds is performed in successive steps. For the *complete flag manifold*⁵ $F_{123;4} = \mathbf{U}(4)/(\mathbf{U}(1))^4$, we start from the Plücker coordinates for a line in \mathbb{C}^4 , a^i , and add a plane containing this line, $A_2^{ij} = a^{[i} b^{j]}$ as well as a hyperplane containing this plane, $A_3^{ijk} = a^{[i} b^{j} c^{k]}$. We arrive at the set of coordinates

$$a^{[i} b^{j} c^{k]} , \quad a^{[i} b^{j]} , \quad a^i , \quad (2.25)$$

from which we can construct the auxiliary coordinates

$$\begin{aligned} x_{123;4}^{\hat{w}_1 \dots \hat{w}_6} &= \bar{a}^{[i_1} \bar{b}^{i_2} \bar{c}^{i_3]} (\lambda^{\hat{w}_1 \hat{w}_2 \hat{w}_3})_{i_1 i_2 i_3, j_1 j_2 j_3} a^{[j_1} b^{j_2} c^{j_3]} \times \\ &\times \bar{a}^{[i_4} \bar{b}^{i_5]} (\lambda^{\hat{w}_4 \hat{w}_5})_{i_4 i_5, j_4 j_5} a^{[j_4} b^{j_5]} \bar{a}^{i_6} (\lambda^{\hat{w}_6})_{i_6 j_6} a^{j_6} . \end{aligned} \quad (2.26)$$

⁵The manifolds consisting of non-maximal flags are called *partial flag manifolds*.

The Hopf fibration underlying this contraction reads as

$$1 \longrightarrow \mathrm{U}(1) \times \mathrm{U}(1) \times \mathrm{U}(1) \longrightarrow S^7 \times S^5 \times S^3 \longrightarrow F_{123;4} \longrightarrow 1, \quad (2.27)$$

and the three $\mathrm{U}(1)$ factors leave invariant the three factors in $x^{\hat{w}_1 \dots \hat{w}_6}$.

On the remaining flag manifolds, the Plücker coordinates are given by subsets of the coordinates for $F_{123;4}$. For example, on $F_{12;4} = \mathrm{U}(4)/(\mathrm{U}(2) \times \mathrm{U}(1) \times \mathrm{U}(1))$, we have the Plücker coordinates

$$a^{[i}b^j], \quad a^i \quad (2.28)$$

with obvious auxiliary coordinates. The Hopf fibration reads as

$$1 \longrightarrow \mathrm{U}(1) \times \mathrm{U}(1) \longrightarrow S^7 \times S^5 \longrightarrow F_{12;4} \longrightarrow 1. \quad (2.29a)$$

This fibration is a reduction of the Hopf fibration for $F_{2;4}$ to the case in which the (internal) isotropy group of the flags is merely $\mathrm{U}(1) \times \mathrm{U}(1)$.

The construction of $F_{13;4} = \mathrm{U}(4)/(\mathrm{U}(1) \times \mathrm{U}(2) \times \mathrm{U}(1))$ and $F_{23;4} = \mathrm{U}(4)/(\mathrm{U}(1) \times \mathrm{U}(1) \times \mathrm{U}(2))$ follows the same line of argument, and the two Hopf fibrations read as

$$1 \longrightarrow \mathrm{U}(1) \times \mathrm{U}(2) \longrightarrow S^7 \times S^5 \times S^3 \longrightarrow F_{13;4} \longrightarrow 1, \quad (2.29b)$$

$$1 \longrightarrow \mathrm{U}(2) \times \mathrm{U}(1) \longrightarrow S^7 \times S^5 \times S^3 \longrightarrow F_{23;4} \longrightarrow 1, \quad (2.29c)$$

respectively. In particular, the flag manifold $F_{13;4}$ is described by the set of Plücker coordinates $(a^{[i}b^j c^k], a^i)$. In twistor theory, the common description of this space is in terms of a quadric in the space $\mathbb{C}P^3 \times \mathbb{C}P^3_*$ with coordinates a^i and a_i^* . The quadric condition reads as $a^i a_i^* = 0$ and with the identification $a_i^* = \varepsilon_{ijkl} a^j b^k c^l$, its relation to the Plücker description becomes clear.

Also the reducible flag manifolds can be mapped to the space of certain tensor products of projectors. For example in the case $F_{12;4}$, we combine $\mathcal{P}_{2;4} = \mathcal{P}_{1;4}^1 \wedge \mathcal{P}_{1;4}^2$ with an additional rank one projector $\mathcal{P}_{1;4}^3$ given by a linear combination of $\mathcal{P}_{1;4}^1$ and $\mathcal{P}_{1;4}^2$:

$$\mathcal{P}_{1;4}^3 = \alpha \mathcal{P}_{1;4}^1 + \beta \mathcal{P}_{1;4}^2, \quad \alpha^2 + \beta^2 = 1. \quad (2.30)$$

Thus, $\mathcal{P}_{1;4}^3$ projects onto a one-dimensional subspace of the plane which $\mathcal{P}_{2;4}$ projects onto. The definition of $\mathcal{P}_{1;4}^3$ implies that the coordinates are linear combinations:

$$x_3^{\hat{a}} = \alpha x_1^{\hat{a}} + \beta x_2^{\hat{a}}, \quad (2.31)$$

and together with the constraints on the projectors and the antisymmetrization of $\mathcal{P}_{1;4}^1$ and $\mathcal{P}_{1;4}^2$ in $\mathcal{P}_{2;4}$, this equation describes an embedding of the flag manifold $F_{12;4}$ in Euclidean space.

For the complete flag manifold $F_{123;4}$, we use altogether six rank one projectors, combined in $\mathcal{P}_{3;4}^{123}$, $\mathcal{P}_{2;4}^{45}$ and $\mathcal{P}_{1;4}^6$, each satisfying equation (2.7) and furthermore fulfilling conditions corresponding to (2.31). The coordinates $x_{123}^{\hat{a}\hat{b}\hat{c}}$, $x_{45}^{\hat{a}\hat{b}}$ and $x_6^{\hat{a}}$ form an over-complete set of coordinates on $F_{123;4}$, and the restrictions we impose are an embedding of $F_{123;4}$ in Euclidean space.

All of the reducible flag manifolds can again be described in terms of rank one projectors \mathcal{P} . Explicitly, these projectors read as:

$$\begin{aligned}
\mathcal{P}_{12;4} &= x^{\hat{w}_1 \hat{w}_2 \hat{w}_3} \lambda^{\hat{w}_1 \hat{w}_2} \otimes \lambda^{\hat{w}_3} , \\
\mathcal{P}_{13;4} &= x^{\hat{w}_1 \dots \hat{w}_4} \lambda^{\hat{w}_1 \dots \hat{w}_3} \otimes \lambda^{\hat{w}_4} , \\
\mathcal{P}_{23;4} &= x^{\hat{w}_1 \dots \hat{w}_5} \lambda^{\hat{w}_1 \dots \hat{w}_3} \otimes \lambda^{\hat{w}_4 \hat{w}_5} , \\
\mathcal{P}_{123;4} &= x^{\hat{w}_1 \dots \hat{w}_6} \lambda^{\hat{w}_1 \dots \hat{w}_3} \otimes \lambda^{\hat{w}_4 \hat{w}_5} \otimes \lambda^{\hat{w}_6} ,
\end{aligned} \tag{2.32}$$

where the $x^{\hat{w}_1 \dots \hat{w}_k}$ are the auxiliary coordinates constructed from the (independent) Plücker coordinates on the various flag manifolds. Note that in all cases the number of generators in the projector corresponds to the sum of the dimensions of the nested vector spaces in the flag. Even though this description contains a vast redundancy, it turns out to be rather convenient for describing the geometric structures on the flag manifolds inherited from the embedding in Euclidean space, which is the purpose of the next section.

2.5. Geometric structures on the flag manifolds

In this section, we will develop expressions for the complex structure, the metric and the symplectic structure on the flag manifolds introduced above, following closely [6]. Given a projector \mathcal{P}^0 , which describes a point on the flag manifold $M = F_{i_1 \dots i_k; 4}$, all of M is obtained by an appropriate action of $U(4)$ on \mathcal{P}^0 . That is, the tangent directions are given by infinitesimal actions of $U(4)$ and thus the space of tangent vectors is

$$T_{\mathcal{P}^0} M = \{R(\Lambda) \mathcal{P}^0 | \Lambda \in \mathfrak{su}(4)\} . \tag{2.33}$$

Here, we have to distinguish the different representations R of Λ for the different projectors \mathcal{P}^0 used for the various flag manifolds. For projectors consisting of k -fold antisymmetric combinations of rank 1 projectors, $R(\Lambda)$ is the sum of a k -fold tensor product with all entries 1 but one, which is $\text{ad}_\Lambda := i[\Lambda, \cdot]$. In particular, we have

$$\begin{aligned}
R(\Lambda) &:= \text{ad}_\Lambda \quad \text{for } \mathcal{P}_{1;4} , \\
R(\Lambda) &:= \text{ad}_\Lambda \otimes \mathbb{1} + \mathbb{1} \otimes \text{ad}_\Lambda \quad \text{for } \mathcal{P}_{2;4} , \\
R(\Lambda) &:= \text{ad}_\Lambda \otimes \mathbb{1} \otimes \mathbb{1} + \mathbb{1} \otimes \text{ad}_\Lambda \otimes \mathbb{1} + \mathbb{1} \otimes \mathbb{1} \otimes \text{ad}_\Lambda \quad \text{for } \mathcal{P}_{3;4} .
\end{aligned} \tag{2.34}$$

for the irreducible flag manifolds. The representations in the case of reducible flag manifolds are constructed in an obvious manner, and one has e.g.

$$\begin{aligned}
R(\Lambda) &:= \text{ad}_\Lambda \otimes \mathbb{1} \otimes \text{ad}_\Lambda + \mathbb{1} \otimes \text{ad}_\Lambda \otimes \mathbb{1} \quad \text{for } \mathcal{P}_{12;4} , \\
R(\Lambda) &:= \text{ad}_\Lambda \otimes \mathbb{1} \otimes \mathbb{1} \otimes \text{ad}_\Lambda + \mathbb{1} \otimes \text{ad}_\Lambda \otimes \mathbb{1} \otimes \mathbb{1} + \mathbb{1} \otimes \mathbb{1} \otimes \text{ad}_\Lambda \otimes \mathbb{1} \quad \text{for } \mathcal{P}_{13;4} .
\end{aligned} \tag{2.35}$$

If and only if Λ is a generator of H , $R(\Lambda) \mathcal{P}^0$ vanishes, and therefore $T_{\mathcal{P}^0} M$ is of the same dimension as M . By construction, we have for an element $V \in T_{\mathcal{P}^0} M$:

$$V^\dagger = V , \quad \{\mathcal{P}^0, V\} = V , \quad \text{tr} V = 0 . \tag{2.36}$$

The orthogonal complement of $T_{\mathcal{P}^0} M$ in the embedding space is spanned by all other actions of $U(4)$ onto \mathcal{P}^0 . In particular for $\mathbb{C}P^3$, the generators κ^a of the stabilizer subgroup

$H = \mathbf{U}(3) \times \mathbf{U}(1)$ of \mathcal{P}^0 span the orthogonal complement of $T_{\mathcal{P}^0}M$ in the embedding space \mathbb{R}^{16} as they satisfy by definition $[\mathcal{P}^0, \kappa^a] = 0$ and therefore they are orthogonal to any element of $T_{\mathcal{P}^0}M$: $\text{i tr}(\kappa^a[\Lambda, \mathcal{P}^0]) = 0$.

To define a complex structure I , we start from such a structure on the embedding space, which in turn induces a complex structure on the tangent space at \mathcal{P}^0 . Consider the generators $\lambda^{\hat{a}}$ of $\mathfrak{u}(4)$. We can pair them into $(\lambda^{2p}, \lambda^{2p+1})$, $p = 0 \dots 7$ and define $I(\lambda^{2p}, \lambda^{2p+1}) = (-\lambda^{2p+1}, \lambda^{2p})$, which amounts to the canonical complex structure on \mathbb{R}^{16} . This translates into a complex structure on any general embedding space and the pairing together with the projection onto $T_{\mathcal{P}^0}M$ is performed by taking the commutator with \mathcal{P}^0 :

$$IV = -\text{i}[\mathcal{P}^0, V] \quad \text{with} \quad V \in T_{\mathcal{P}^0}M . \quad (2.37)$$

One easily checks that $I^2 = -1$ and $IW = 0$ for $W \in T_{\mathcal{P}^0}^\perp M$. This definition extends from $T_{\mathcal{P}^0}M$ to the full tangent bundle and yields an almost complex structure on M , which turns out to be integrable.

Also the metric is induced from the one on the embedding space, which is the Euclidean (Killing) metric and, after translation into matrices, simply given by the trace. To incorporate the projection onto $T_{\mathcal{P}^0}M$, we can multiply each vector in $T_{\mathcal{P}^0}M$ by the complex structure before taking the trace which yields the hermitian metric

$$g(V_1, V_2) = \text{tr}(IV_1 IV_2) = -\text{tr}([\mathcal{P}^0, V_1][\mathcal{P}^0, V_2]) , \quad (2.38)$$

as for elements $V_1, V_2 \in T_{\mathcal{P}^0}M$, we have $g(V_1, V_2) = g(IV_1, IV_2)$. The continuation of this metric to all of M is evident. There is furthermore a symplectic structure defined as

$$\Omega(V_1, V_2) := g(IV_1, V_2) , \quad (2.39)$$

which we can combine as usually with the metric into the Kähler structure J defined as

$$J(V_1, V_2) := \frac{1}{2}(g(V_1, V_2) + \text{i}\Omega(V_1, V_2)) = \text{tr}(\mathcal{P}^0 V_1 (\mathbb{1} - \mathcal{P}^0) V_2) , \quad (2.40)$$

which also extends globally.

Note that the projectors we use in the description of flag manifolds are of rank one, and therefore, the above formula simplifies to

$$J(V_1, V_2) = \text{tr}(\mathcal{P}^0 V_1 V_2) - \text{tr}(\mathcal{P}^0 V_1) \text{tr}(\mathcal{P}^0 V_2) . \quad (2.41)$$

On $\mathbb{C}P^3$, Λ is a generator of the fundamental representation of $\mathbf{SU}(4)$ and we can introduce the components $\Omega_{1;4}^{\hat{a}\hat{b}} = \Omega(\lambda^{\hat{a}}, \lambda^{\hat{b}})$ as well as $J_{1;4}^{\hat{a}\hat{b}} = J(\lambda^{\hat{a}}, \lambda^{\hat{b}})$. For these, we have the useful identities

$$\Omega_{1;4}^{\hat{a}\hat{b}} = \sqrt{2} f^{ab}{}_c x^c \quad (2.42)$$

and

$$\begin{aligned} J_{1;4}^{\hat{a}\hat{b}} \lambda_{\hat{b}} &= \text{tr}(\mathcal{P}^0 \lambda^{\hat{a}} (\mathbb{1} - \mathcal{P}^0) \lambda^{\hat{b}}) \lambda_{\hat{b}} = \mathcal{P}^0 \lambda^{\hat{a}} (\mathbb{1} - \mathcal{P}^0) , \\ \lambda_{\hat{a}} J_{1;4}^{\hat{a}\hat{b}} &= \lambda_{\hat{a}} \text{tr}(\mathcal{P}^0 \lambda^{\hat{a}} (\mathbb{1} - \mathcal{P}^0) \lambda^{\hat{b}}) = (\mathbb{1} - \mathcal{P}^0) \lambda^{\hat{b}} \mathcal{P}^0 . \end{aligned} \quad (2.43)$$

In deriving these identities, one needs the relation

$$\text{tr}(\lambda^{\hat{b}} \lambda^{\hat{a}}) \lambda_{\hat{a}} = \lambda^{\hat{b}} \quad \text{for} \quad \lambda^{\hat{a}}, \lambda^{\hat{b}} \in \mathfrak{u}(4) , \quad (2.44)$$

cf. formulæ (B.23). Due to $(\mathcal{P}^0)^2 = \mathcal{P}^0$, the relations (2.43) remain valid after omitting the hats over the indices.

Let us briefly comment on the explicit form of the structures obtained in the above discussion for the various flag manifolds. We start with the Graßmannians. These spaces are described by rank one projectors in the representation R , which is here the previously defined k -fold \wedge -product of the fundamental one. The tangent directions are given by infinitesimal actions of elements of $U(4)$:

$$\begin{aligned} T_{\mathcal{P}_{2;4}^0} F_{2;4} &= \{(\text{ad}_\Lambda \otimes \mathbb{1} + \mathbb{1} \otimes \text{ad}_\Lambda) \mathcal{P}_{2;4}^0 | \Lambda \in \mathfrak{su}(4)\} , \\ T_{\mathcal{P}_{3;4}^0} F_{3;4} &= \{(\text{ad}_\Lambda \otimes \mathbb{1} \otimes \mathbb{1} + \mathbb{1} \otimes \text{ad}_\Lambda \otimes \mathbb{1} + \mathbb{1} \otimes \mathbb{1} \otimes \text{ad}_\Lambda) \mathcal{P}_{3;4}^0 | \Lambda \in \mathfrak{su}(4)\} . \end{aligned} \quad (2.45)$$

Note that the most general action e.g. on the projector $\mathcal{P}_{2;4}^0$ is given by $\text{Ad}_{g_1} \otimes \text{Ad}_{g_2}$. Since $\mathcal{P}_{2;4}^0$ is the sum of antisymmetrized tensor products of the form $\lambda^{\hat{a}} \wedge \lambda^{\hat{b}}$, only the *symmetrized form* of $\text{Ad}_{g_1} \otimes \text{Ad}_{g_2}$ is relevant, which is $\frac{1}{2}((\text{Ad}_{g_1} + \text{Ad}_{g_2}) \otimes (\text{Ad}_{g_1} + \text{Ad}_{g_2}))$. At infinitesimal level, this yields the action $\text{ad}_\Lambda \otimes \mathbb{1} + \mathbb{1} \otimes \text{ad}_\Lambda$.

All the properties (2.36) are easily verified to hold also for the tangent vectors of all the Graßmannians. Furthermore, the definitions of the complex structure, the metric and the Kähler structure is done in a straightforward manner, since the only essential aspect in their definition on $\mathbb{C}P^3$ was that \mathcal{P}^0 is a projector.

Using again the shorthand notation (2.19), the appropriate components of the symplectic and the Kähler structure are given by

$$\begin{aligned} \Omega_{2;4}^{\hat{a}\hat{b},\hat{c}\hat{d}} &= \Omega(\lambda^{\hat{a}\hat{b}}, \lambda^{\hat{c}\hat{d}}) \quad \text{and} \quad \Omega_{3;4}^{\hat{a}\hat{b}\hat{c},\hat{d}\hat{e}\hat{f}} = \Omega(\lambda^{\hat{a}\hat{b}\hat{c}}, \lambda^{\hat{d}\hat{e}\hat{f}}) , \\ J_{2;4}^{\hat{a}\hat{b},\hat{c}\hat{d}} &= J(\lambda^{\hat{a}\hat{b}}, \lambda^{\hat{c}\hat{d}}) \quad \text{and} \quad J_{3;4}^{\hat{a}\hat{b}\hat{c},\hat{d}\hat{e}\hat{f}} = J(\lambda^{\hat{a}\hat{b}\hat{c}}, \lambda^{\hat{d}\hat{e}\hat{f}}) . \end{aligned} \quad (2.46)$$

The corresponding versions of the identities (2.42), (2.43) read as

$$\Omega_{2;4}^{\hat{a}\hat{0},\hat{c}\hat{d}} = {}_{(cd)}\sqrt{2} f^{acb} x^{bd} \quad \text{and} \quad \Omega_{3;4}^{\hat{a}\hat{0}\hat{0},\hat{d}\hat{e}\hat{f}} = {}_{(def)}\sqrt{2} f^{adb} x^{bef} , \quad (2.47)$$

where these relations only hold for the components symmetric in (cd) and (def) , respectively, and

$$\begin{aligned} J_{2;4}^{\hat{a}\hat{b},\hat{c}\hat{d}} \lambda^{\hat{c}\hat{d}} &= \text{tr}(\mathcal{P}^0 \lambda^{\hat{a}\hat{b}} (\mathbb{1} \otimes \mathbb{1} - \mathcal{P}^0) \lambda^{\hat{c}\hat{d}}) \lambda^{\hat{c}\hat{d}} = \mathcal{P}^0 \lambda^{\hat{a}\hat{b}} (\mathbb{1} \otimes \mathbb{1} - \mathcal{P}^0) , \\ J_{3;4}^{\hat{a}\hat{b}\hat{c},\hat{d}\hat{e}\hat{f}} \lambda^{\hat{d}\hat{e}\hat{f}} &= \text{tr}(\mathcal{P}^0 \lambda^{\hat{a}\hat{b}\hat{c}} (\mathbb{1} \otimes \mathbb{1} \otimes \mathbb{1} - \mathcal{P}^0) \lambda^{\hat{d}\hat{e}\hat{f}}) \lambda^{\hat{d}\hat{e}\hat{f}} = \mathcal{P}^0 \lambda^{\hat{a}\hat{b}\hat{c}} (\mathbb{1} \otimes \mathbb{1} \otimes \mathbb{1} - \mathcal{P}^0) , \end{aligned} \quad (2.48)$$

which follow from the relations in (B.23).

To describe the tangent space to the reducible flag manifolds $M = F_{k_1 k_2;4}$ at a point $\mathcal{P}_{k_1 k_2;4}^0 = \mathcal{P}_{k_2}^0 \otimes \mathcal{P}_{k_1(k_2);4}^0$, one proceeds completely analogously to above and defines

$$T_{\mathcal{P}_{k_1 k_2;4}^0} M := \{(\text{ad}_\Lambda \otimes \mathbb{1} \otimes \dots + \mathbb{1} \otimes \text{ad}_\Lambda \otimes \dots + \dots \otimes \mathbb{1} \otimes \text{ad}_\Lambda) (\mathcal{P}_{k_2;4}^0 \otimes \mathcal{P}_{k_1(k_2);4}^0) | \Lambda \in \mathfrak{su}(4)\} . \quad (2.49)$$

It immediately follows that the elements of $T_{\mathcal{P}_{k_1 k_2;4}^0} M$ all satisfy (2.36). The stabilizer subgroup H of $\mathcal{P}_{k_1 k_2;4}^0$ is indeed $U(4 - k_2) \times U(k_2 - k_1) \times U(k_1)$.

The definition of the complex structure, the metric, the symplectic and the Kähler structure are again straightforward. For the latter, we introduce components, e.g. for $F_{12;4}$:

$$J_{12;4}^{\hat{a}\hat{b},\hat{c}\hat{d};\hat{e}\hat{f}} = J_{2;4}^{\hat{a}\hat{b},\hat{c}\hat{d}} \otimes J_{1;4}^{\hat{e}\hat{f}} , \quad (2.50)$$

and one has again obvious identities corresponding to (2.43).

2.6. Spherical functions on the flag manifolds

Before discussing the fuzzification of functions on the flag manifolds, let us briefly review some aspects of harmonic analysis on these spaces. That is, we want to describe the construction of spherical functions on flag manifolds, which form a complete orthonormal basis on these spaces and are simultaneously eigenfunctions of the Laplace operator. For this, we will extend the standard discussion of spherical functions using the various generalized Hopf fibrations described above. (A detailed discussion for $\mathbb{C}P^n = F_{1;n}$ is found e.g. in [25].)

The spherical functions on the sphere S^m are simply the restrictions of the homogeneous harmonic polynomials on \mathbb{R}^{m+1} to S^m [26] and the dimension of the eigenspaces H^j , $j \in \mathbb{N}$, corresponding to the eigenvalue $j(j+m-1)$ are $\binom{m+j}{m}$. For $m = 2n-1$, the eigenspaces H^j are spanned by homogeneous polynomials in the complex coordinates a^i on \mathbb{C}^n plus their complex conjugate minus all terms containing contractions $a^i \bar{a}^i$, as these terms belong to spaces H^k with $k < j$.

The complex projective space $\mathbb{C}P^3$ is obtained from the generalized Hopf fibration $U(1) \rightarrow S^7 \rightarrow \mathbb{C}P^3$. The eigenfunctions of the Laplace operator on this space are the subset of the corresponding eigenfunctions on S^7 which are invariant under $U(1)$. These functions are obviously the product of a homogeneous polynomial of order k in a^i and another such polynomial in \bar{a}^j minus all the possible contractions. We thus get the ‘‘hyperspherical harmonics’’

$$Y_{1;4}^k{}^{i_1 \dots i_k j_1 \dots j_k} := a^{i_1} \dots a^{i_k} \bar{a}^{j_1} \dots \bar{a}^{j_k} - \text{contraction terms} , \quad (2.51)$$

which have eigenvalues $\lambda_k = k(k+3)$ and their eigenspaces $H_{1;4}^k$ have dimensions

$$\dim H_{1;4}^k = \binom{k+3}{k}^2 - \binom{k+2}{k-1}^2 , \quad (2.52)$$

where the last term subtracts the dimensions of the contraction terms.

The Grassmannian $G_{2;4}$ is obtained from the Hopf fibration $U(2) \rightarrow S^7 \times S^5 \rightarrow G_{2;4}$. The spherical functions on $S^7 \times S^5$ are constructed from homogeneous polynomials in A^{ij} and \bar{A}^{kl} , where we add again the complex conjugate polynomial to render the expression real and finally subtract all terms containing contractions. The subspace of $U(2)$ -invariant functions is now spanned by those polynomials, which have an equal number of A^{ij} s and \bar{A}^{kl} s, and we thus have:

$$Y_{2;4}^k{}^{i_1 j_1 \dots i_k j_k l_1 m_1 \dots l_k m_k} := A^{i_1 j_1} \dots A^{i_k j_k} \bar{A}^{l_1 m_1} \dots \bar{A}^{l_k m_k} - \text{contraction terms} . \quad (2.53)$$

From the generalized Hopf fibrations discussed in the preceding sections, the construction of the remaining flag manifolds is obvious. For our purposes, it is more important to note that these spherical functions on G/H are in one-to-one correspondence with so-called *spherical representations* of G/H , i.e. (finite dimensional) representations of G with non-trivial H -invariant vectors, see e.g. [27, 28]. This allows us to associate each set of eigenfunctions of the Laplace operator on the flag manifolds with certain sets of irreducible representations of G , which we will do in the next section. Later on, we will use these representations to construct the algebra of fuzzy functions on the flag manifolds. From this construction, one can also read off the eigenvalues of the eigenspaces of the Laplace operator on the various flag manifolds.

3. Flag manifolds and coherent states

To quantize the flag manifolds, we would like to connect every point on these spaces to a state in a Hilbert space. A function then automatically becomes an operator on this space. To establish this connection, recall that every point on a flag manifold which is a coset space of $G = \text{U}(n)$ is in one-to-one correspondence to a generalized coherent state in a specific representation of G . We will review this relation in the next section and partly follow the discussion of Perelomov [29], see [30, 15] for quantization using coherent states.

3.1. Representations of $\text{SU}(n)$ and coherent states

Consider the Dynkin diagram of $\text{SU}(n)$ for simple roots $\alpha_1, \dots, \alpha_{n-1}$

$$\begin{array}{ccccccc} a_1 & & a_2 & & & & a_{n-1} \\ \circ & \text{---} & \circ & \text{---} & \dots & \text{---} & \circ \\ \alpha_1 & & \alpha_2 & & & & \alpha_{n-1} \end{array} \quad (3.1)$$

An irreducible representation T^Λ with highest weight Λ can be labeled by the Dynkin indices⁶ a_1, \dots, a_{n-1} . In the representation (Hilbert) space \mathcal{H}^Λ , there is a corresponding highest weight vector $|\Lambda\rangle$ and \mathcal{H}^Λ has a basis of weight vectors⁷ $\{|\mu\rangle\}$ i.e. $H_j|\mu\rangle = \mu_j|\mu\rangle$.

The isotropy subgroup H_μ for any weight vector $|\mu\rangle$ contains the Cartan subgroup H of $\text{SU}(n)$, which is isomorphic to the maximal torus $T^{n-1} = \text{U}(1)^{\times n-1} = \text{U}(1) \times \dots \times \text{U}(1)$ and for general weight vectors the subgroup H_μ coincides with T^{n-1} . For degenerate representation, where the highest weight Λ is orthogonal to some simple root α_i , i.e.

$$(\Lambda, \alpha_i) = 0 = a_i, \quad (3.2)$$

the isotropy subgroup H_μ may be larger than T^{n-1} for some weight vectors. This is evident since, as explained in appendix B, the a_i indicate the highest power of $E_{-\alpha}$, whose action is still nontrivial on $|\Lambda\rangle$.

A helpful picture arises, when enlarging $\text{SU}(n)$ to $\text{U}(n)$. The Cartan subalgebra consists now of n factors of $\text{U}(1)$, which one can imagine sitting around the a_i . Every a_i which is zero combines the $\text{U}(m_1)$ left of it with the $\text{U}(m_2)$ right to it to a $\text{U}(m_1 + m_2)$. This allows us to construct all the isotropy groups one encounters in flag manifolds, and we will be more explicit in the next section.

To construct a coherent state system, one has to choose an initial vector $|\psi_0\rangle$ in \mathcal{H}^Λ . Then the system of states $\{|\psi_g\rangle = T^\Lambda(g)|\psi_0\rangle\}$ is called the coherent state system $\{T^\Lambda, |\psi_0\rangle\}$. Let H_0 be the isotropy subgroup for the state $|\psi_0\rangle$. Then a coherent state $|\psi_g\rangle$ is determined by a point $x = x(g)$ in the coset space G/H_0 , corresponding to the element g by $|\psi_g\rangle = \exp(i\alpha)|x\rangle$ up to a phase, $|\psi_0\rangle = |0\rangle$. The isotropy subgroup for a linear combination of weight vectors is, in general, a subgroup of the Cartan subgroup. Therefore it is convenient to choose a weight vector $|\mu\rangle$ as an initial state. For non-degenerate representations, the isotropy subgroup H_μ is isomorphic to the Cartan subgroup H , and the coherent state $|x\rangle$ is characterized by a point of G/H , or equivalently by a point of the orbit of the adjoint representation

$$H'_j|x\rangle = T(g)H_jT^{-1}(g)|x\rangle, \quad |x\rangle = T(g)|\mu\rangle. \quad (3.3)$$

⁶See appendix B for more details.

⁷which can be chosen to be orthonormal

In a representation with some Dynkin labels vanishing, the isotropy subgroup H_μ is larger than H for some weight vectors $|\mu\rangle$, in which case the orbit may be degenerate.

There is considerable choice in the selection of the initial state $|0\rangle$, even on restriction to weight vectors. Perelomov [31] has shown that the state $|0\rangle$ must be $|\mu\rangle$, where μ is a dominant weight, if it is to be closest to classical. That is, $|\mu\rangle$ is obtained from the highest weight by the Weyl reflection group. Then the coherent states minimize the invariant uncertainty relation

$$\Delta C_2 = \min , \quad (3.4)$$

where

$$C_2 = \sum_j (H_j)^2 + \sum_{\alpha \in \Sigma_+} (E_\alpha E_{-\alpha} + E_{-\alpha} E_\alpha) \quad (3.5)$$

is the quadratic Casimir operator and

$$\Delta C_2 = \langle C_2 \rangle - \left(\sum_j \langle H_j \rangle^2 + 2 \sum_{\alpha \in \Sigma_+} \langle E_\alpha \rangle \langle E_{-\alpha} \rangle \right) . \quad (3.6)$$

Let us take the initial state to be the highest weight vector $|\Lambda\rangle$. The coherent state system is then more explicitly defined by

$$|a\rangle = N T^\Lambda(g) |\Lambda\rangle , \quad N^{-2} = \langle \Lambda | T^\Lambda(g) | \Lambda \rangle , \quad (3.7)$$

$$\begin{aligned} |a\rangle &= N \exp \left(\sum_{\alpha \in \Sigma_+} a_\alpha^- E_{-\alpha} \right) \exp (h_i H_i) \exp \left(\sum_{\alpha \in \Sigma_+} a_\alpha^+ E_\alpha \right) |\Lambda\rangle \\ &= N \exp \left(\sum_{\alpha \in \Sigma_+} a_\alpha^- E_{-\alpha} \right) |\Lambda\rangle , \end{aligned} \quad (3.8)$$

where we have restricted ourselves to elements of G which have a Gaußian decomposition. Note that if some of the Dynkin labels a_i vanish (degenerate representations), then the corresponding coordinates a_α^- (and possibly some others, see the appendix) are no longer independent and can be eliminated from the definition of the group element g and thus the coherent states correspond to points on various flag manifolds. The number of independent coordinates a_α^- gives the (complex) dimension of the flag manifold.

Note that our construction of coherent states yields merely one patch of the covering of the flag manifolds. Starting from a different dominant weight state corresponds to working on a different patch, as the state $|w(\Lambda)\rangle$, where w is an element of the Weyl group W , is not contained in the set of the coherent states $|a\rangle$ constructed from $|\Lambda\rangle$. This is because we have restricted ourselves to group elements that have a Gaußian decomposition. We can assume all the weight states to be orthogonal. In particular, two states $|\Lambda\rangle$ and $|w(\Lambda)\rangle$ have no overlap, i.e. $\langle \Lambda | w(\Lambda) \rangle = 0$. Consider now the coherent state $|a\rangle$ as constructed above,

$$|a\rangle = N \left(1 + \sum_{n=1}^{\infty} \frac{1}{n!} \left(\sum_{\alpha \in \Sigma_+} a_\alpha^- E_{-\alpha} \right)^n \right) |\Lambda\rangle . \quad (3.9)$$

As $E_{-\alpha}$ contains only lowering operators, we have $\langle \Lambda | E_{-\alpha} | \Lambda \rangle = 0$, which implies

$$\langle \Lambda | a^- \rangle = \langle \Lambda | N | \Lambda \rangle + 0 = N . \quad (3.10)$$

It is thus clear that all $|a\rangle$ have a component parallel to $|\Lambda\rangle$ and therefore $|a\rangle$ never equals $|w(\Lambda)\rangle$, which we wanted to prove.

The number of such dominant weight states from which we can start and thus the number of patches is evidently given by the dimension of $\{w(\Lambda), w \in W\}$ or equivalently the number of corners of the convex hull of the states in the weight diagram of the representation. This number is just the rank of the Weyl group modulo group elements acting trivially in a certain representation. As the Weyl group for $SU(n)$ is essentially the permutation group of n elements with rank $n!$, the minimal number of patches covering the complete flag manifold $F_{123\dots n-1;n}$ is given by $n!$, while the number of patches for all other flag manifolds of $SU(n)$ is smaller. The reason for this is simply that for other flag manifolds, certain Dynkin labels a_i are zero, which implies that the corresponding Weyl reflections

$$|S_{\alpha_i}\Lambda\rangle = |\Lambda - \frac{2(\alpha_i, \Lambda)}{(\alpha_i, \alpha_i)}\alpha_i\rangle, \quad (3.11)$$

act trivially on the highest weight state.

To clarify the above construction, we will first discuss the simple case of flags in $SU(3)$, whose weight diagrams are two-dimensional, before presenting the construction for the flag manifolds of $SU(4)$.

3.2. Examples for $SU(3)$


There are two flag manifolds which arise as coset spaces of $SU(3)$: The complex projective space $\mathbb{C}P^2 = F_{1;3} \cong F_{2;3} = \mathbb{C}P^2_* \cong \frac{SU(3)}{U(2)}$ and the reducible flag manifold $F_{12;3} = \frac{SU(3)}{U(1) \times U(1)}$. The representations **3**, **$\bar{3}$** and **6** of $SU(3)$, corresponding to the diagrams

$$\begin{array}{c} 1 \\ \circ \end{array} \text{---} \begin{array}{c} 0 \\ \circ \end{array} = \square, \quad \begin{array}{c} 0 \\ \circ \end{array} \text{---} \begin{array}{c} 1 \\ \circ \end{array} = \begin{array}{|c|} \hline \square \\ \hline \end{array}, \quad \begin{array}{c} 2 \\ \circ \end{array} \text{---} \begin{array}{c} 0 \\ \circ \end{array} = \square\square, \quad (3.12)$$

all have dominant weight states with isotropy group $U(2)$. Thus, the coherent states constructed from these representations are in one-to-one correspondence with points on $\mathbb{C}P^2$. The adjoint representation **8** as well as the **27** corresponding to the diagrams

$$\begin{array}{c} 1 \\ \circ \end{array} \text{---} \begin{array}{c} 1 \\ \circ \end{array} = \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \quad \begin{array}{c} 2 \\ \circ \end{array} \text{---} \begin{array}{c} 1 \\ \circ \end{array} = \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array} \quad (3.13)$$

have dominant weight states, whose isotropy group is only the maximal torus $U(1) \times U(1)$, and therefore the derived coherent states are in one-to-one correspondence with points on the flag manifold $F_{12;3}$. Recall that the number of patches is found to be the number of corners in (the convex hulls of) the weight diagrams. Consider e.g. the diagrams




corresponding to the representations with Dynkin labels $(1, 0)$, $(1, 1)$ and $(2, 1)$, respectively. We see that the complex projective space $\mathbb{C}P^2$ is covered by at least three patches, while the flag manifold $F_{12;3}$ requires a minimum of six patches.

3.3. The flags in \mathbb{C}^4

Let us now come to the coherent states which correspond to points on flag manifolds of $SU(4)$. For these, the choice of representations as well as the Dynkin diagrams are given in Table 1.

Note that the representations for the reducible flag manifolds are not unique at level L . One can choose any representation $(a_1(L), a_2(L), a_3(L))$; however, for considering the limit $L \rightarrow \infty$, the functions a_i should be polynomials of the same order in L .

The minimal numbers of patches for the various flag manifolds are again the numbers of corners in the weight diagrams for the various representations. For example, consider the weight diagrams of the representations $(1, 0, 0)$, $(1, 0, 1)$, $(1, 1, 1)$,



which yield coherent states corresponding to $\mathbb{C}P^3$, $F_{13;4}$ and $F_{123;4}$ with minimal coverings of 4, 12 and 24 patches, respectively.

	$\dim_{\mathbb{C}}$	patches	isotropy group in $U(4)$	Dynkin labels	Young diagrams
$F_{1;4}$	3	4	$U(1) \times U(3)$	$(L, 0, 0)$	$\overbrace{\begin{array}{ c c c } \hline \square & \square & \square \\ \hline \end{array}}^L$
$F_{2;4}$	4	6	$U(2) \times U(2)$	$(0, L, 0)$	$\overbrace{\begin{array}{ c c } \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}}^L$
$F_{3;4}$	3	4	$U(3) \times U(1)$	$(0, 0, L)$	$\overbrace{\begin{array}{ c c c } \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}}^L$
$F_{12;4}$	5	12	$U(1) \times U(1) \times U(2)$	$(L, L, 0)$	$\overbrace{\begin{array}{ c c c c } \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}}^{L+L}$
$F_{13;4}$	5	12	$U(1) \times U(2) \times U(1)$	$(L, 0, L)$	$\overbrace{\begin{array}{ c c c c } \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}}^{L+L}$
$F_{23;4}$	5	12	$U(2) \times U(1) \times U(1)$	$(0, L, L)$	$\overbrace{\begin{array}{ c c c c } \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}}^{L+L}$
$F_{123;4}$	6	24	$U(1) \times U(1) \times U(1) \times U(1)$	(L, L, L)	$\overbrace{\begin{array}{ c c c c c } \hline \square & \square & \square & \square & \square \\ \hline \square & \square & \square & \square & \square \\ \hline \end{array}}^{L+L+L}$

Table 1: Representations of $SU(4)$ related to the flag manifolds in \mathbb{C}^4 .

4. Fuzzification of the flag manifolds

Combining the description of the flag manifolds in terms of Plücker coordinates which we developed in section 2 with the correspondence to coherent states in the previous section, we have an obvious way in which one can truncate the algebra of functions on these spaces to obtain the latter's fuzzy versions. Before we describe the construction of the fuzzy flag manifolds in detail, let us briefly recall the underlying principles.

4.1. Fuzzification

By *fuzzy geometry*, we mean a truncation of the algebra of functions on a compact space such that the coordinates become noncommutative while all isometries are manifestly preserved. Given a compact Riemannian manifold M without boundary, the spectrum of the Laplace operator is discrete and the eigenfunctions form an orthogonal basis B of $L^2(M)$. A naïve guess for a discretization would be to truncate the expansion of a function by using only a finite subset B^L of elements in B . Multiplication of functions, however, clearly necessitates a subsequent projection back on to B^L , which in turn will render the product non-associative in general.

If the manifold $M = G/H$ is a coadjoint orbit of a Lie group G , we can easily circumvent this problem: we can map functions to operators acting as automorphisms⁸ on the representation space of some representation R of G which admits singlets under H and replace the product between functions by the operator product. In the previous section, we described which representations R are suitable for the various flag manifolds of $SU(4)$. We will see that the choice of R corresponds to a choice of the truncation and the closure of multiplication is trivially given.

Using a projector $\rho_R(x) = |x\rangle\langle x|$ which corresponds to a point $x \in G/H$ and acts on the representation space of R , we can establish a map between operators and functions on the coset space by the formula

$$f_R(x) = \text{tr} \left(\rho_R(x) \hat{f} \right) . \quad (4.1)$$

The operator product then induces a star product via

$$(f_R \star g_R)(x) = \text{tr} \left(\rho_R(x) \hat{f} \hat{g} \right) . \quad (4.2)$$

In general, there is an infinite sequence of suitable representations R_i for any coset space and for each of these representations, the star product is different. Choosing higher-dimensional representations amounts to a better approximation of the functions by operators, and there is usually a well-defined limit, in which the complete set of functions on the coset together with the ordinary product is reproduced.

Let us return to equation (4.1). To each operator \hat{f} representing a function on M_R , assign a corresponding symbol $\tilde{f}(g)$, $g \in G$ by [8]

$$\tilde{f}(g) = n_R \text{tr} \left(D^R(g^{-1}) \hat{f} \right) , \quad (4.3)$$

⁸That is, they can be represented by square matrices.

where $D^R(g)$ is the group element g acting in the representation R . The normalization constant n_R is defined by

$$\int d\mu(g) D^R(g^{-1})_{ij} D^R(g)_{kl} = \frac{1}{n_R} \delta_{il} \delta_{jk} \quad (4.4)$$

with $d\mu(g)$ being the Haar measure on G . Inversely, the operator \hat{f} corresponding to the symbol \tilde{f} is therefore obtained from

$$\hat{f} = \int d\mu(g) \tilde{f}(g) D^R(g) . \quad (4.5)$$

From the symbol \tilde{f} of an operator \hat{f} , we can easily calculate the function defined in (4.1) using

$$f_R(x) = \int d\mu(g) \omega_R(x, g) \tilde{f}(g) \quad \text{with} \quad \omega_R(x, g) := \text{tr} (\rho_R(x) D^R(g)) . \quad (4.6)$$

In the definition of the star product (4.2), this translates into

$$(f_R \star g_R)(x) = \int d\mu(g) \int d\mu(g') \omega_R(x, gg') \tilde{f}(g) \tilde{g}(g') , \quad (4.7)$$

and it is for this formula that we will find explicit expressions for all the flag manifolds later on.

In the discussion of fuzzy flag manifolds using star products, we can use both of the two equivalent descriptions: either real coordinates describing an embedding of the coset space into flat Euclidean space or the complex homogeneous or Plücker coordinates. In the latter coordinates the star product can be shown to simplify considerably. Moreover, they allow for a direct translation to the operator picture.

Note that so far, we only arrived at an algebra of functions on a topological space. The explicit geometry of this space, i.e. its metric structure, has not been described yet. In noncommutative geometry, this information is encoded in a Dirac operator, or – in a slightly weaker way – in a Laplacian. Using the above mentioned embedding, we obtain a canonical metric on the coset space and can show that the Laplace operator naturally translates into the second order Casimir in the representation R .

For more details on the principle underlying fuzzification, see also [6].

4.2. The fuzzy complex projective space $\mathbb{C}P_F^3$

The fuzzification of $F_{1,4}$ (and therefore also that of its dual $F_{3,4}$) is well-known [6], and we follow the usual discussion of the procedure for $\mathbb{C}P^N$. That is, we promote the vector a and its complex conjugate \bar{a} to a four-tuple of annihilation and creation operators satisfying the algebra $[\hat{a}^i, \hat{a}^{j\dagger}] = \delta^{ij}$. The auxiliary coordinates defined in (2.4) also become operators

$$\hat{x}^{\hat{a}} := \frac{1}{\hat{a}^{k\dagger} \hat{a}^l \delta_{kl}} \hat{a}^{i\dagger} \lambda_{ij}^{\hat{a}} \hat{a}^j , \quad (4.8)$$

which evidently commute with the number operator $\hat{N} = \hat{a}^{k\dagger} \hat{a}^l \delta_{kl}$. Therefore, we can restrict the algebra of functions to the subspace of the Fock space, on which $\hat{N} = L$. This subspace is spanned by

$$\frac{1}{C} \hat{a}^{i_1\dagger} \dots \hat{a}^{i_L\dagger} |0\rangle \quad \text{with} \quad C = \sqrt{n_1! n_2! n_3! n_4!} , \quad (4.9)$$

where the n_i are the number of indices being i . The truncated algebra of functions \mathcal{A}_L on this space is the algebra of operators with basis

$$\hat{a}^{i_1\dagger} \dots \hat{a}^{i_L\dagger} |0\rangle \langle 0| \hat{a}^{j_1} \dots \hat{a}^{j_L} . \quad (4.10)$$

It is immediately obvious that these operators will commute with the number operator \hat{N} , which amounts to factoring out a $U(1)$ as implied in the definition of any flag manifold. The coefficients of the expansion of an operator in terms of the basis (4.10) form square matrices of dimension $\left(\frac{(3+L)!}{3!L!}\right)^2$ [6], and in terms of Young diagrams of $SU(4)$, we have

$$\overbrace{\square \square \square \square}^L \otimes \overbrace{\square \square \square \square}^L = \overbrace{\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}}^L \otimes \overbrace{\square \square \square \square}^L . \quad (4.11)$$

To expose the underlying $SU(4)$ structure and to construct the polarization tensors, we can contract indices from the creation operators with indices from the annihilation operators using the 15 generators λ_{ij}^a of $SU(4)$. A contraction with $\lambda_{ij}^0 \sim \delta_{ij}$ yields the embedded subalgebra truncated at level $L - 1$, since the trace over a fundamental and an antifundamental index corresponds to the determinant over four indices in either the fundamental or antifundamental representation and thus to a column of 4 boxes in a Young diagram, which is cancelled. The tensor product expansion looks as

$$\overbrace{\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}}^L \otimes \overbrace{\square \square \square \square}^L = \mathbf{1} \oplus \overbrace{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}} \oplus \overbrace{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} \oplus \dots \oplus \overbrace{\begin{array}{|c|c|c|c|c|c|} \hline \square & \square & \square & \square & \square & \square \\ \hline \square & \square & \square & \square & \square & \square \\ \hline \square & \square & \square & \square & \square & \square \\ \hline \square & \square & \square & \square & \square & \square \\ \hline \end{array}}^{2L} . \quad (4.12)$$

A representation of the Lie algebra of $SU(4)$ is given by the Schwinger construction and we can write

$$\hat{L}^{\hat{a}} = \hat{a}^{i\dagger} \lambda_{ij}^{\hat{a}} \hat{a}^j . \quad (4.13)$$

One can easily verify the algebra $[\hat{L}^a, \hat{L}^b] = i\sqrt{2} f^{ab}{}_c \hat{L}^c$ using $[\hat{a}^i, \hat{a}^{j\dagger}] = \delta^{ij}$.

In the representations $R = R(L)$ introduced above, the projector $\mathcal{P}_{1;4} = \mathcal{P}(x_{1;4})$, which describes the embedding of $\mathbb{C}P^3$ in \mathbb{R}^{15} , is simply the L -fold symmetrized tensor product $\rho_{1;4}^L = \rho^L(x_{1;4}) = \mathcal{P}(x_{1;4}) \otimes \dots \otimes \mathcal{P}(x_{1;4})$, and we can map any operator \hat{f} in the algebra \mathcal{A}_L to a corresponding function f_L on the embedding of $\mathbb{C}P^3$ in \mathbb{R}^{15} by

$$f_L(x_{1;4}) = \text{tr}(\rho^L(x_{1;4}) \hat{f}) . \quad (4.14)$$

Furthermore, this map induces a star product on $\mathbb{C}P^3$ defined as

$$(f_L \star g_L)(x_{1;4}) = f_L(x_{1;4}) \star g_L(x_{1;4}) = \text{tr}(\rho^L(x_{1;4}) \hat{f} \hat{g}) , \quad (4.15)$$

where f_L and g_L are the functions corresponding to the operators \hat{f} and \hat{g} , respectively.

To make the star product more explicit, we calculate $\omega^L(x_{1;4}, gg')$ for the fundamental representation $L = 1$:

$$\omega^L(x_{1;4}, gg') = \text{tr}(\mathcal{P}_{1;4} gg') = \text{tr}(\mathcal{P}_{1;4} g \mathcal{P}_{1;4} g') + \text{tr}(\mathcal{P}_{1;4} g (\mathbb{1} - \mathcal{P}_{1;4}) g') . \quad (4.16)$$

Since $\mathcal{P}_{1;4}$ is a rank one projector, we have

$$\text{tr}(\mathcal{P}_{1;4} g \mathcal{P}_{1;4} g') = \text{tr}(\mathcal{P}_{1;4} g) \text{tr}(\mathcal{P}_{1;4} g') = \omega^L(x_{1;4}, g) \omega^L(x_{1;4}, g') \quad (4.17)$$

and with the identities (2.43), it follows immediately that

$$\begin{aligned} \text{tr}(\mathcal{P}_{1;4}g(\mathbb{1} - \mathcal{P}_{1;4})g') &= \text{tr}(\lambda^a g) \text{tr}(\mathcal{P}_{1;4}\lambda^a(\mathbb{1} - \mathcal{P}_{1;4})g') \\ &= \left(\frac{\partial}{\partial x^a} \omega(x_{1;4}, g) \right) J^{ab} \left(\frac{\partial}{\partial x^b} \omega(x_{1;4}, g') \right) . \end{aligned} \quad (4.18)$$

For the representations with $L > 1$, we can simply take the L -fold tensor product of $\omega^1(x_{1;4}, g)$ [8]:

$$\omega^L(x_{1;4}, g) = (\omega^1(x_{1;4}, g))^{\otimes L} , \quad (4.19)$$

and the total star product reads as [6]

$$(f_L \star g_L)(x_{1;4}) = \sum_{l=0}^L \frac{(L-l)!}{L!l!} (\partial_{a_1 \dots a_l} f_L(x_{1;4})) J_{1;4}^{a_1 b_1} \dots J_{1;4}^{a_l b_l} (\partial_{b_1 \dots b_l} g_L(x_{1;4})) . \quad (4.20)$$

In the homogeneous coordinates a^i, \bar{a}^i on $\mathbb{C}P^3$, the space of functions is spanned by homogeneous polynomials of the form

$$\bar{a}^{i_1} \dots \bar{a}^{i_L} a^{j_1} \dots a^{j_L} , \quad (4.21)$$

which correspond to the operators (4.10) under the map (4.14). In these coordinates, the star product simplifies to [32]

$$(f \star g) = \mu \left[\frac{1}{L!} \frac{\partial}{\partial a^{\alpha_1}} \dots \frac{\partial}{\partial a^{\alpha_L}} \otimes \frac{1}{L!} \frac{\partial}{\partial \bar{a}^{\alpha_1}} \dots \frac{\partial}{\partial \bar{a}^{\alpha_L}} (f \otimes g) \right] , \quad (4.22)$$

where $\mu(a \otimes b) = a \cdot b$. Note that this construction of a star product generalizes in a rather straightforward way to other spaces, as soon as we have a suitable projector $\rho^L(x)$ at hand.

To relate the given matrix algebra to the space $\mathbb{C}P^3$, we need some additional structure to encode the geometry of this space. For this, consider the vector fields on $\mathbb{C}P^3$ from the perspective of the embedding space \mathbb{R}^{16} :

$$\mathcal{L}_a = -\sqrt{2} f_{ab}{}^c x^b \frac{\partial}{\partial x^c} , \quad [\mathcal{L}_a, \mathcal{L}_b] = i\sqrt{2} f_{ab}{}^c \mathcal{L}_c , \quad (4.23)$$

where $f_{ab}{}^c$ are the structure constants of $\text{SU}(4)$. Note here that in the limit $L \rightarrow \infty$, the fuzzy derivatives approach the ones from the continuum in an obvious way. It is now rather straightforward to show that [6]

$$\mathcal{L}^a f_L(x) = \sqrt{2} \Omega^{ab} \frac{\partial}{\partial x^b} f_L(x) = \frac{L}{\sqrt{2}} (x^a \star f_L(x) - f_L(x) \star x^a) = \text{tr} \left(\rho^L(x_{1;4}) [\hat{L}^a, \hat{f}] \right) , \quad (4.24)$$

where $[\hat{L}^a, \cdot]$ are the generators of $\text{SU}(4)$ in the representation $(L, 0, 0) \otimes (0, 0, L)$. It is therefore also clear, that the Laplace operator on $\mathbb{C}P^3$, $\Delta = \mathcal{L}^a \mathcal{L}^b \delta_{ab}$, is mapped to the second order Casimir in the adjoint representation $(L, 0, 0) \otimes (0, 0, L)$:

$$\hat{\Delta}_{1;4} \hat{f} = [\hat{L}^a, [\hat{L}^b, \hat{f}]] \delta_{ab} . \quad (4.25)$$

4.3. The fuzzy Graßmannian $G_{2;4}^F$

We proceed analogously to the case of $\mathbb{C}P^3$, which leads to the results presented in [8] in a somewhat simpler form. That is, we take the Plücker description discussed in section 2.3 and promote the vector components to creation and annihilation operators $\hat{a}^i, \hat{a}^{i\dagger}, \hat{b}^i, \hat{b}^{i\dagger}$. We thus arrive at the algebra

$$[\hat{a}^i, \hat{a}^{j\dagger}] = [\hat{b}^i, \hat{b}^{j\dagger}] = \delta^{ij}, \quad (4.26)$$

and all other commutators vanish. From these operators, we construct the composite creation and annihilation operators

$$\hat{A}_2^{ij} = \hat{a}^{[i}\hat{b}^{j]} \quad \text{and} \quad \hat{A}_2^{i\dagger j\dagger} = \hat{a}^{[i\dagger}\hat{b}^{j\dagger]}, \quad (4.27)$$

which satisfy

$$[\hat{A}_2^{ij}, \hat{A}_2^{kl\dagger}] = \left(\delta^{ik}\delta^{jl} + \delta^{jl}\hat{a}^{k\dagger}\hat{a}^i + \delta^{ik}\hat{b}^{l\dagger}\hat{b}^j \right)_{[ij][kl]}, \quad (4.28)$$

where $(\cdot)_{[ij][kl]}$ denotes antisymmetrization of the enclosed components, as well as

$$[[\hat{A}_2^{ij}, \hat{A}_2^{kl\dagger}], \hat{A}_2^{mn\dagger}] = \left(2\delta^{jl}\delta^{im}\hat{A}_2^{kn\dagger} \right)_{[ij][kl][mn]}. \quad (4.29)$$

We can now use $\hat{A}_2^{mn\dagger}$ to build an L -particle⁹ Hilbert space $\mathcal{H}_{1;4}^L$. This space is spanned by

$$\frac{1}{C}\hat{A}_2^{i_1 j_1 \dagger} \dots \hat{A}_2^{i_L j_L \dagger} |0\rangle, \quad (4.30)$$

where C is the norm of the state. Acting with \hat{A}_2^{mn} on such a state yields a state in $\mathcal{H}_{2;4}^{L-1}$ due to (4.28) and (4.29). Recall that in the Plücker description of $G_{2;4}$, we constructed the plane by antisymmetrizing two vectors, which could be chosen orthogonal i.e. $\bar{a}^i b^i = 0$. On the operator level, this translates into

$$[\hat{a}^{i\dagger}\hat{b}^i, \hat{A}_2^{kl\dagger}] = 0, \quad [\hat{a}^i\hat{b}^{i\dagger}, \hat{A}_2^{kl\dagger}] = 0, \quad [\hat{a}^{i\dagger}\hat{b}^i, \hat{A}_2^{kl}] = 0 \quad \text{and} \quad [\hat{a}^i\hat{b}^{i\dagger}, \hat{A}_2^{kl}] = 0, \quad (4.31)$$

and therefore the action of $\hat{a}^{i\dagger}\hat{b}^i$ on any state in $\mathcal{H}_{2;4}^L$ vanishes. This implies that we can introduce the number operator

$$\hat{N} = \hat{a}^{i\dagger}\hat{a}^i \delta_{ij} = \hat{b}^{i\dagger}\hat{b}^i \delta_{ij} = L \quad (4.32)$$

and

$$\hat{A}_2^{i\dagger j\dagger} \hat{A}_2^{kl} \delta_{ik} \delta_{jl} = 2\hat{N}(\hat{N} - 1) = 2L(L - 1), \quad (4.33)$$

where the equalities hold only after restriction to $\mathcal{H}_{2;4}^L$. Thus, $\mathcal{H}_{2;4}^L$ is indeed an L -particle Hilbert space. We already know from the discussion in the previous section, that the states (4.30) are invariant under $S(U(2) \times U(2))$. Let us nevertheless be more explicit on the action of the internal $SU(2)$, which acts nontrivially on both \hat{a}^i and \hat{b}^i . Its generators L_{int}^r act according to

$$L_{\text{int}}^r = \text{ad} \left(\hat{a}_p^{i\dagger} \lambda_{pq}^r \hat{a}_q^i \right), \quad (4.34)$$

⁹Note that a particle is here a composite object consisting of two excitations.

where $\hat{a}_1^i = \hat{a}^i$, $\hat{a}_2^i = \hat{b}^i$ and λ_{pq}^r , $p, q = 1, 2$, $r = 1, 2, 3$ are the Gell-Mann matrices of $SU(2)$. The combinations A_2^{ij} and $A_2^{ij\dagger}$ are now invariant under this action due to the general formula

$$[a_p^{i\dagger} \lambda_{pq}^r a_q^i, a_1^{[j_1\dagger} \dots a_k^{j_k]^\dagger}] = 0, \quad (4.35)$$

where $p = 1, \dots, k$ and $i, j_p = 1, \dots, n$; see appendix B for a proof.

The truncated algebra of functions \mathcal{A}_L is the algebra of operators spanned by

$$\hat{A}_2^{i_1 j_1 \dagger} \dots \hat{A}_2^{i_L j_L \dagger} |0\rangle \langle 0| \hat{A}_2^{k_1 l_1} \dots \hat{A}_2^{k_L l_L}, \quad (4.36)$$

and the coefficients in an expansion in terms of these operators are square matrices of size $\frac{(3+L)!(2+L)!}{3!L!2!(L+1)!}$. Note that these operators again commute with the number operator \hat{N} and in terms of Young diagrams, we have here

$$\begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline & & \\ \hline \end{array} \otimes \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline & & \\ \hline \end{array} = \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline & & \\ \hline \end{array} \otimes \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline & & \\ \hline \end{array}. \quad (4.37)$$

Again, each of the tensor product decompositions at level L contains the tensor product decomposition at lower levels:

$$\begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline & & \\ \hline \end{array} \otimes \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline & & \\ \hline \end{array} = \mathbf{1} \oplus \begin{array}{|c|} \hline \\ \hline \\ \hline \end{array} \oplus \begin{array}{|c|c|} \hline & \\ \hline & \\ \hline & \\ \hline \end{array} \oplus \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline & & \\ \hline \end{array} \oplus \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array} \oplus \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array} \oplus \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array} \oplus \dots \quad (4.38)$$

Contrary to the case of $\mathbb{C}P^3$, where increasing the level L by one yielded precisely one new type of Young tableau in the sum, we here obtain $L + 1$ new diagrams in each step, which consist of three rows with $a + b + a$, $a + b$ and a boxes respectively. The new diagrams at level L are the ones for which $a = n$ and $b = 2L - 2n$ for $n = 1 \dots L$.

All these notions readily translate for arbitrary Grassmannians.

To find a representation of the Lie algebra of $SU(4)$, we use a generalized Schwinger construction

$$\hat{L}^a = \frac{1}{\hat{N}} \hat{A}_2^{ij\dagger} \Lambda_{ijkl}^a \hat{A}_2^{kl} \quad \text{with} \quad \Lambda_{ijkl}^a = (\lambda^a \wedge \mathbb{1})_{ijkl}. \quad (4.39)$$

It is important to stress that the \hat{L}^a by themselves do not form a representation of $SU(4)$, but again after having them act on a state in $\mathcal{H}_{2,4}^L$, they do. This is simply due to the fact that because of (4.31), \hat{L}^a reduces when acting on a state in $\mathcal{H}_{2,4}^L$ to

$$\hat{L}^a = \hat{a}^{i\dagger} \lambda_{ij}^a \hat{a}^j + \hat{b}^{i\dagger} \lambda_{ij}^a \hat{b}^j. \quad (4.40)$$

The projector yielding a star product on this space is the symmetrized L -fold tensor product $\rho_{2,4}^L = \rho^L(x_{2,4}) = \mathcal{P}_{2,4} \otimes \dots \otimes \mathcal{P}_{2,4}$. Proceeding precisely along the lines of the discussion of the star product on $\mathbb{C}P^3$, we find that

$$\omega^1(x^{\hat{a}\hat{b}}, gg') = \omega^1(x, g) \left(1 + \overleftarrow{\partial}_{\hat{a}\hat{b}} J^{\hat{a}\hat{b}, \hat{c}\hat{d}} \overrightarrow{\partial}_{\hat{c}\hat{d}} \right) \omega^1(x, g'), \quad (4.41)$$

where at least one of the indices in each $\hat{a}\hat{b}$ or $\hat{c}\hat{d}$ is nonzero. That is, the component x^{00} plays a similar rôle to the component x^0 in the case of $\mathbb{C}P^3$. Furthermore, one should stress that as usual for the derivatives on spaces with symmetrized tensors as coordinates, one has

$$\partial_{\hat{a}\hat{b}} = \begin{cases} \frac{\partial}{\partial x^{\hat{a}\hat{b}}} & \text{for } \hat{a} = \hat{b} \neq 0 \\ \frac{1}{2} \frac{\partial}{\partial x^{\hat{a}\hat{b}}} & \text{for } \hat{a} \neq \hat{b} \end{cases}. \quad (4.42)$$

It is then quite obvious that the star product is given by

$$(f_L \star g_L)(x_{2;4}) = \sum_{l=0}^L \frac{(L-l)!}{L!l!} \left(\partial_{(\hat{a}_1 \hat{b}_1) \dots (\hat{a}_l \hat{b}_l)} f_L(x_{2;4}) \right) J_{2;4}^{\hat{a}_1 \hat{b}_1; \hat{c}_1 \hat{d}_1} \dots \left(\partial_{(\hat{c}_1 \hat{d}_1) \dots (\hat{c}_l \hat{d}_l)} g_L(x_{2;4}) \right), \quad (4.43)$$

where $\partial_{(\hat{a}_1 \hat{b}_1) \dots (\hat{a}_l \hat{b}_l)} := \partial_{\hat{a}_1 \hat{b}_1} \dots \partial_{\hat{a}_l \hat{b}_l}$. Note that our choice of embedding $F_{2;4}$ in $\mathbb{R}^{16 \cdot 17/2 - 1}$ yielded a slightly simpler expression for the star product on this space than the one in [8], which used an embedding in \mathbb{R}^{15} .

The expression for the star product further simplifies, if we switch again to complex coordinates $a^{[i} b^{j]}$ on $\mathbb{C}^4 \wedge \mathbb{C}^4$. The functions corresponding to the operators (4.36) read as

$$\bar{a}^{[i_1 \bar{j}^1]} \dots \bar{a}^{[i_L \bar{j}^L]} a^{[k_1 b^1]} \dots a^{[k_L b^L]} \quad (4.44)$$

and the star product is here defined as

$$(f \star g) = \mu \left[\frac{1}{L!L!} \frac{\partial}{\partial a^{[i_1}} \frac{\partial}{\partial b^{j_1]} } \dots \frac{\partial}{\partial a^{[i_L}} \frac{\partial}{\partial b^{j_L]} } \otimes \frac{1}{L!L!} \frac{\partial}{\partial \bar{a}^{[i_1}} \frac{\partial}{\partial \bar{b}^{j_1]} } \dots \frac{\partial}{\partial \bar{a}^{[i_L}} \frac{\partial}{\partial \bar{b}^{j_L]} } (f \otimes g) \right].$$

The natural Laplacian on $\mathcal{H}_{2;4}^L$ encoding the geometry of $G_{2;4}$ is derived from the embedding of $G_{2;4}$ in $\mathbb{R}^{16 \cdot 17/2 - 1}$. In terms of Plücker coordinates, the generators read as

$$\mathcal{L}^{\hat{a}} = \bar{a}^i \lambda_{ij}^{\hat{a}} \frac{\partial}{\partial \bar{a}^j} + \bar{b}^i \lambda_{ij}^{\hat{a}} \frac{\partial}{\partial \bar{b}^j} - a^j \lambda_{ij}^{\hat{a}} \frac{\partial}{\partial a^i} - b^j \lambda_{ij}^{\hat{a}} \frac{\partial}{\partial b^i}. \quad (4.45)$$

From this expression and equation (2.9), we obtain the following expression in terms of the embedding coordinates:

$$\mathcal{L}_a = -i\sqrt{2} f_{ab}{}^c x^{bd} \frac{\partial}{\partial x^{cd}}, \quad (4.46)$$

which satisfies the algebra $[\mathcal{L}^a, \mathcal{L}^b] = i\sqrt{2} f^{ab}{}_c \mathcal{L}^c$, as is easily verified. Using the first relation in (2.47), we can write

$$\mathcal{L}^a f_L(x) = \sqrt{2} I^{a0,cd} \frac{\partial}{\partial x^{cd}} f_L(x) = x^{a0} \star f_L - f_L \star x^{a0} = -i \operatorname{tr}(\rho^L(x_{2;4}) [\hat{L}^a, \hat{f}_L]). \quad (4.47)$$

We thus see again that all the derivatives are mapped to the generators of $\mathrm{SU}(4)$ acting in the adjoint, and therefore the Laplacian is given by the second order Casimir of this representation of $\mathrm{SU}(4)$,

$$\hat{\Delta}_{2;4} = \operatorname{ad}_{\hat{L}^a} \operatorname{ad}_{\hat{L}^b} \delta_{ab}. \quad (4.48)$$

This observation from the cases $\mathbb{C}P^3$ and $G_{2;4}$ translates to all flag manifolds and we suppress this calculation in the remaining cases. For the discussion of the eigenvalues of these Casimirs, see appendix B.

4.4. The fuzzy dual complex projective space $F_{3;4}^F$

One can infer the fuzzification of $F_{3;4}$ in a straightforward manner from the ones of $F_{1;4}$ and $F_{2;4}$. We start from the Plücker description and promote the vectors a, b, c to a triple

of four-tuples of oscillators with creation and annihilation operators $\hat{a}^i, \hat{a}^{i\dagger}, \hat{b}^i, \hat{b}^{i\dagger}, \hat{c}^i, \hat{c}^{i\dagger}$. We furthermore introduce the composite operators

$$\hat{A}_3^{ijk} = \hat{a}^{[i} \hat{b}^j \hat{c}^{k]} \quad \text{and} \quad \hat{A}_3^{ijk\dagger} = \hat{a}^{[i\dagger} \hat{b}^{j\dagger} \hat{c}^{k\dagger]}, \quad (4.49)$$

satisfying the commutation relations

$$\begin{aligned} [\tilde{d}_i, \tilde{d}_m^\dagger] &:= \varepsilon_{ijkl} \varepsilon_{mnrs} [\hat{A}_3^{jkl}, \hat{A}_3^{nrs}] \\ &= \delta_{im} + \varepsilon_{ijkl} \varepsilon_{mnrs} \left(\hat{a}^{n\dagger} \hat{a}^j \delta^{rk} \delta^{sl} + \delta^{nj} \hat{b}^{r\dagger} \hat{b}^k \delta^{sl} + \delta^{nj} \delta^{rk} \hat{c}^{s\dagger} \hat{c}^l + \right. \\ &\quad \left. \hat{a}^{n\dagger} \hat{a}^j \hat{b}^{r\dagger} \hat{b}^k \delta^{sl} + \hat{a}^{n\dagger} \hat{a}^j \delta^{rk} \hat{c}^{s\dagger} \hat{c}^l + \delta^{jn} \hat{b}^{r\dagger} \hat{b}^k \hat{c}^{s\dagger} \hat{c}^l \right). \end{aligned} \quad (4.50)$$

The expression for $[[[\tilde{d}_i, \tilde{d}_j^\dagger] \tilde{d}_k^\dagger] \tilde{d}_l^\dagger]$ contains only the combination $\tilde{d}_m^\dagger = \varepsilon_{mnrs} \hat{A}_3^{nrs\dagger}$.

The L -particle Fock space $\mathcal{H}_{3;4}^L$ is evidently spanned by the states

$$\hat{A}_3^{i_1 j_1 k_1 \dagger} \dots \hat{A}_3^{i_L j_L k_L \dagger} |0\rangle, \quad (4.51)$$

and this space forms the representation $(0, 0, L)$ of $\text{SU}(4)$. The isotropy subgroup of any state in this representation is thus $\text{S}(\text{U}(3) \times \text{U}(1))$, and the internal $\text{SU}(3)$ action, affecting all the elementary oscillators $\hat{a}, \hat{b}, \hat{c}$ is given by

$$\hat{L}_{\text{int}}^r = \text{ad} \left(\hat{a}_p^{i\dagger} \lambda_{pq}^r \hat{a}_q^i \right), \quad (4.52)$$

where $\hat{a}_1^i, \hat{a}_2^i, \hat{a}_3^i$ stand for $\hat{a}^i, \hat{b}^i, \hat{c}^i$, respectively, and λ^r are the Gell-Mann matrices of $\text{SU}(3)$. Invariance of the operators \hat{A}_3^{ijk} and $\hat{A}_3^{ijk\dagger}$ follows from equation (4.35). It is this representation which underlies the construction of vector bundles over $\mathbb{C}P_F^3$ [21].

The algebra of functions truncated at level L is again constructed from two copies of the Fock space and their $\text{SU}(4)$ transformation property is captured by the diagrams

$$\begin{array}{c} L \\ \hline \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline & & \\ \hline \end{array} \end{array} \otimes \begin{array}{c} L \\ \hline \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline & & \\ \hline \end{array} \end{array} = \begin{array}{c} L \\ \hline \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline & & \\ \hline \end{array} \end{array} \otimes \begin{array}{c} L \\ \hline \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline & & \\ \hline \end{array} \end{array}. \quad (4.53)$$

We clearly see that this algebra is dual to (4.11), i.e. that of $F_{1;4}^F$. For this reason, we will not go into any further details.

The definition of a star product is obvious. The projector in the symmetrized L -fold tensor product reads as $\rho_{3;4}^L(x_{3;4}) = \mathcal{P}_{3;4} \otimes \dots \otimes \mathcal{P}_{3;4}$ and yields

$$\begin{aligned} (f_L \star g_L)(x_{3;4}) &= \sum_{l=0}^L \frac{(L-l)!}{L!l!} \left(\partial_{(\hat{a}_1 \hat{b}_1 \hat{c}_1) \dots (\hat{a}_l \hat{b}_l \hat{c}_l)} f_L(x_{3;4}) \right) J_{3;4}^{\hat{a}_1 \hat{b}_1 \hat{c}_1; \hat{d}_1 \hat{e}_1 \hat{f}_1} \dots \\ &\quad J_{3;4}^{\hat{a}_l \hat{b}_l \hat{c}_l; \hat{d}_l \hat{e}_l \hat{f}_l} \left(\partial_{(\hat{d}_1 \hat{e}_1 \hat{f}_1) \dots (\hat{d}_l \hat{e}_l \hat{f}_l)} g_L(x_{3;4}) \right), \end{aligned} \quad (4.54)$$

where again the components x^{000} are dropped in the formula and

$$\partial_{abc} = \begin{cases} \frac{\partial}{\partial x^{abc}} & \text{for } \hat{a} = \hat{b} = \hat{c} \neq 0 \\ \frac{1}{2} \frac{\partial}{\partial x^{\hat{a}\hat{b}\hat{c}}} & \text{for } \hat{a} = \hat{b} \neq \hat{c} \text{ etc.} \\ \frac{1}{6} \frac{\partial}{\partial x^{\hat{a}\hat{b}\hat{c}}} & \text{for } \hat{a} \neq \hat{b} \neq \hat{c} \neq \hat{a} \end{cases}. \quad (4.55)$$

The discussion of the star product formalism in the complex coordinates $a^{[ij]c^k}$ on $\mathbb{C}^4 \wedge \mathbb{C}^4 \wedge \mathbb{C}^4$ is trivially deduced from the cases $\mathbb{C}P_F^3$ and G_2^F . The star product here reads as

$$(f \star g) = \mu \left[\frac{1}{L!L!L!} \frac{\partial}{\partial a^{[i_1}} \frac{\partial}{\partial b^{j_1}} \frac{\partial}{\partial c^{k_1}} \cdots \frac{\partial}{\partial a^{[i_L}} \frac{\partial}{\partial b^{j_L}} \frac{\partial}{\partial c^{k_L}} \right] \otimes \left[\frac{1}{L!L!L!} \frac{\partial}{\partial \bar{a}^{[i_1}} \frac{\partial}{\partial \bar{b}^{j_1}} \frac{\partial}{\partial \bar{c}^{k_1}} \cdots \frac{\partial}{\partial \bar{a}^{[i_L}} \frac{\partial}{\partial \bar{b}^{j_L}} \frac{\partial}{\partial \bar{c}^{k_L}} \right] (f \otimes g) \Big].$$

4.5. The fuzzy reducible flag manifold $F_{12;4}^F$

In the case of the reducible flag manifolds, one needs a set of composite creation and annihilation operators. These sets are in one-to-one correspondence with the Plücker coordinates. For $F_{12;4}^F$ we thus have

$$\hat{A}_2^{ij}, \hat{A}_2^{ij\dagger} \quad \text{and} \quad \hat{a}^i, \hat{a}^{i\dagger}. \quad (4.56)$$

The missing commutation relations are easily found from (4.28) and (4.29), e.g.

$$[\hat{a}^i, \hat{A}_2^{jk\dagger}] = \delta^{i[j} \hat{b}^{k]\dagger} \quad (4.57)$$

and the (L_1, L_2) -particle Hilbert spaces are now constructed as

$$\hat{A}_2^{i_1 j_1 \dagger} \dots \hat{A}_2^{i_{L_1} j_{L_1} \dagger} \hat{a}^{k_1 \dagger} \dots \hat{a}^{k_{L_2} \dagger} |0\rangle. \quad (4.58)$$

The corresponding operators

$$\hat{A}_2^{i_1 j_1 \dagger} \dots \hat{A}_2^{i_{L_1} j_{L_1} \dagger} \hat{a}^{k_1 \dagger} \dots \hat{a}^{k_{L_2} \dagger} |0\rangle \langle 0| \hat{A}_2^{m_1 n_1} \dots \hat{A}_2^{m_{L_1} n_{L_1}} \hat{a}^{l_1} \dots \hat{a}^{l_{L_2}} \quad (4.59)$$

evidently form a closed algebra and act on the representation space of the representation given in terms of Young diagrams by

$$\begin{array}{c} \overbrace{\text{[Young Diagram 1]} \\ L_1+L_2 \end{array} \otimes \begin{array}{c} \overbrace{\text{[Young Diagram 2]} \\ L_1+L_2 \end{array} = \begin{array}{c} \overbrace{\text{[Young Diagram 3]} \\ L_2+L_1 \end{array} \otimes \begin{array}{c} \overbrace{\text{[Young Diagram 4]} \\ L_1+L_2 \end{array} \quad (4.60)$$

The internal isotropy subgroup here is $U(1) \times U(1)$, which follows from the discussion of the underlying coherent states. The explicit action of this subgroup on the elementary oscillators is given by the number operators for \hat{a}^i and \hat{b}^i .

To define a star product on this space, recall that we could describe the flag manifold $F_{12;4}$ in terms of two rank one projectors \mathcal{P}_2 and \mathcal{P}_1 satisfying $\mathcal{P}_2 \mathcal{P}_1 \mathcal{P}_2 = \mathcal{P}_1$. Furthermore, note that we can split every operator \hat{f} in this representation as

$$\hat{f} = \tilde{f}_{IJ} \hat{h}_2^I \otimes \hat{h}_1^J \quad \text{with} \quad \hat{h}_2^I \in \begin{array}{c} \overbrace{\text{[Young Diagram 5]} \\ L_1 \end{array} \otimes \begin{array}{c} \overbrace{\text{[Young Diagram 6]} \\ L_1 \end{array} \quad \text{and} \quad \hat{h}_1^J \in \begin{array}{c} \overbrace{\text{[Young Diagram 7]} \\ L_2 \end{array} \otimes \begin{array}{c} \overbrace{\text{[Young Diagram 8]} \\ L_2 \end{array}, \quad (4.61)$$

where I and J are multi-indices. To such an operator, a truncated function is assigned by

$$f(x_{2;4}, x_{1(2);4}) = \tilde{f}_{IJ} \text{tr}(\rho^L(x_{2;4}) \hat{h}_2^I) \text{tr}(\rho^L(x_{1(2);4}) \hat{h}_1^J), \quad (4.62)$$

where $\rho^L(x_{2;4})$ and $\rho^L(x_{1(2);4})$ are the projectors $\mathcal{P}(x_{2;4})$ and $\mathcal{P}(x_{1(2);4})$ in the same representations as $\hat{h}_{1,2}^I$ and $x_{1(2);4}$. Furthermore, $x_{2;4}, x_{1(2);4}$ are the coordinates on $F_{12;4}$ embedded

in Euclidean space, of the plane and the included line, respectively. On this embedding, the star product of two operators is defined as

$$(f \star g)(x_{2;4}, x_{1(2);4}) = \tilde{f}_{IJ} \tilde{g}_{MN} \text{tr}(\rho^L(x_{2;4}) \hat{h}_2^I \hat{h}_2^M) \text{tr}(\rho^L(x_{1(2);4}) \hat{h}_1^J \hat{h}_1^N). \quad (4.63)$$

Evidently, the star product between $x_{2;4}$ and $x_{1(2);4}$ is simply the ordinary product. Altogether, the star product on $F_{12;4}$ can be derived from the ones on $\mathbb{C}P^3$ and $G_{2;4}$, and we have

$$\begin{aligned} (\hat{f}_L \star g_L)(x_{12;4}) &= \sum_{l=0}^L \frac{(L-l)!}{L!l!} \sum_{k=0}^L \frac{(L-k)!}{L!k!} \left(\partial_{(\hat{a}_1 \hat{b}_1) \dots (\hat{a}_l \hat{b}_l)} \partial_{\hat{e}_1 \dots \hat{e}_k} \hat{f}_L(x_{12;4}) \right) \times \\ &\times J_{12;4}^{(\hat{a}_1 \hat{b}_1), (\hat{c}_1 \hat{d}_1); \hat{e}_1 \hat{f}_1} \dots J_{12;4}^{(\hat{a}_l \hat{b}_l), (\hat{c}_l \hat{d}_l); \hat{e}_k \hat{f}_k} \left(\partial_{(\hat{c}_1 \hat{d}_1) \dots (\hat{c}_l \hat{d}_l)} \partial_{\hat{f}_1 \dots \hat{f}_k} g_L(x_{12;4}) \right). \end{aligned} \quad (4.64)$$

4.6. The fuzzy reducible flag manifolds $F_{23;4}^F$, $F_{13;4}^F$ and $F_{123;4}^F$

After the discussion of the fuzzy version of $F_{12;4}$, the corresponding constructions for the remaining reducible flag manifolds are quite straightforward. We first choose sets of oscillators, which in turn yield the generators for the algebra of functions:

$$\begin{aligned} F_{13;4} &: \hat{A}_3^{ijk}, \quad \hat{A}_3^{ijk\dagger}, \quad \hat{a}^i, \quad \hat{a}^{i\dagger}, \\ F_{23;4} &: \hat{A}_3^{ijk}, \quad \hat{A}_3^{ijk\dagger}, \quad \hat{A}_2^{ij}, \quad \hat{A}_2^{ij\dagger}, \\ F_{123;4} &: \hat{A}_3^{ijk}, \quad \hat{A}_3^{ijk\dagger}, \quad \hat{A}_2^{ij}, \quad \hat{A}_2^{ij\dagger}, \quad \hat{a}^i, \quad \hat{a}^{i\dagger}. \end{aligned} \quad (4.65)$$

The underlying representations, on which these operators act are given by the Young diagrams

$$\begin{aligned} F_{13;4} &: \begin{array}{c} \overbrace{\square \square \square \square \square}^{L_1+L_2} \\ \square \square \square \square \square \\ \square \square \square \square \square \end{array} \otimes \begin{array}{c} \overbrace{\square \square \square \square \square}^{L_1+L_2} \\ \square \square \square \square \square \\ \square \square \square \square \square \end{array} = \begin{array}{c} \overbrace{\square \square \square \square \square}^{L_2+L_1} \\ \square \square \square \square \square \\ \square \square \square \square \square \end{array} \otimes \begin{array}{c} \overbrace{\square \square \square \square \square}^{L_1+L_2} \\ \square \square \square \square \square \\ \square \square \square \square \square \end{array}, \\ F_{23;4} &: \begin{array}{c} \overbrace{\square \square \square \square \square}^{L_1+L_2} \\ \square \square \square \square \square \\ \square \square \square \square \square \end{array} \otimes \begin{array}{c} \overbrace{\square \square \square \square \square}^{L_1+L_2} \\ \square \square \square \square \square \\ \square \square \square \square \square \end{array} = \begin{array}{c} \overbrace{\square \square \square \square \square}^{L_2+L_1} \\ \square \square \square \square \square \\ \square \square \square \square \square \end{array} \otimes \begin{array}{c} \overbrace{\square \square \square \square \square}^{L_1+L_2} \\ \square \square \square \square \square \\ \square \square \square \square \square \end{array}, \\ F_{123;4} &: \begin{array}{c} \overbrace{\square \square \square \square \square \square}^{L_1+L_2+L_3} \\ \square \square \square \square \square \square \\ \square \square \square \square \square \square \end{array} \otimes \begin{array}{c} \overbrace{\square \square \square \square \square \square}^{L_1+L_2+L_3} \\ \square \square \square \square \square \square \\ \square \square \square \square \square \square \end{array} = \begin{array}{c} \overbrace{\square \square \square \square \square \square}^{L_3+L_2+L_1} \\ \square \square \square \square \square \square \\ \square \square \square \square \square \square \end{array} \otimes \begin{array}{c} \overbrace{\square \square \square \square \square \square}^{L_1+L_2+L_3} \\ \square \square \square \square \square \square \\ \square \square \square \square \square \square \end{array}. \end{aligned} \quad (4.66)$$

From these diagrams, it is evident that $F_{23;4}$ is dual to $F_{12;4}$, and that the spaces $F_{13;4}$ and $F_{123;4}$ are dual to themselves.

Furthermore, the explicit action of the internal isotropy subgroups is easily constructed. For example, in the case $F_{23;4}$, the group acting nontrivial on the elementary oscillators but leaving invariant the states of the Fock space is $SU(2) \times U(1)$. Its action reads as

$$L_{\text{int}}^r = \text{ad} \left(\hat{a}_p^{i\dagger} \lambda_{pq}^r \hat{a}_q^i \right) \quad \text{and} \quad L_{\text{int}}^0 = \text{ad} \left(\hat{c}^{i\dagger} \hat{c}^i \right), \quad (4.67)$$

where as before $\hat{a}_1^i = \hat{a}^i$, $\hat{a}_2^i = \hat{b}^i$ and λ_{pq}^r are the Gell-Mann matrices of $SU(2)$. Evidently, the composite operators A_2^{ij} and $A_2^{ij\dagger}$ and A_3^{ijk} and $A_3^{ijk\dagger}$ are invariant under this combination.

The definition of a star product is performed analogously to the case of $F_{12;4}$. We decompose an operator \hat{f} representing a function on a fuzzy flag manifold into

$$\hat{f} = \tilde{f}_{IJ} \hat{h}_{k_1}^I \otimes \hat{h}_{k_2}^J \quad \text{and} \quad \hat{f} = \tilde{f}_{IJK} \hat{h}_3^I \otimes \hat{h}_2^J \otimes \hat{h}_1^J \quad (4.68)$$

for $F_{k_1 k_2;4}$ and $F_{123;4}$, respectively. The map from operators to functions reads as

$$f(x_{k_2;4}, x_{k_1(k_2);4}) = \tilde{f}_{IJ} \text{tr}(\rho^{L_1}(x_{k_2;4}) \hat{h}_{k_2}^I) \text{tr}(\rho^{L_2}(x_{k_1(k_2);4}) \hat{h}_{k_1}^J) \quad (4.69)$$

and

$$f(x_{3;4}, x_{2(3);4}, x_{1(23);4}) = \tilde{f}_{IJK} \text{tr}(\rho^{L_1}(x_{3;4}) \hat{h}_3^I) \text{tr}(\rho^{L_2}(x_{2(3);4}) \hat{h}_2^J) \text{tr}(\rho^{L_3}(x_{1(23);4}) \hat{h}_1^J), \quad (4.70)$$

which naturally induces a star product via the usual formula. The explicit form of the star products are then obtained from the obvious sums over the differential operators

$$\begin{aligned} F_{13;4} : & \overleftarrow{\partial}_{(\hat{a}_1 \hat{b}_1 \hat{c}_1) \dots (\hat{a}_l \hat{b}_l \hat{c}_l)} \overleftarrow{\partial}_{\hat{g}_1 \dots \hat{g}_k} J_{13;4}^{(\hat{a}_1 \hat{b}_1 \hat{c}_1), (\hat{d}_1 \hat{e}_1 \hat{f}_1); \hat{g}_1 \hat{h}_1} \dots \\ & J_{13;4}^{(\hat{a}_l \hat{b}_l \hat{c}_l), (\hat{d}_l \hat{e}_l \hat{f}_l); \hat{g}_k \hat{h}_k} \overrightarrow{\partial}_{(\hat{d}_1 \hat{e}_1 \hat{f}_1) \dots (\hat{d}_l \hat{e}_l \hat{f}_l)} \overrightarrow{\partial}_{\hat{h}_1 \dots \hat{h}_k}, \\ F_{23;4} : & \overleftarrow{\partial}_{(\hat{a}_1 \hat{b}_1 \hat{c}_1) \dots (\hat{a}_l \hat{b}_l \hat{c}_l)} \overleftarrow{\partial}_{(\hat{g}_1 \hat{h}_1) \dots (\hat{g}_k \hat{h}_k)} J_{23;4}^{(\hat{a}_1 \hat{b}_1 \hat{c}_1), (\hat{d}_1 \hat{e}_1 \hat{f}_1); (\hat{g}_1 \hat{h}_1), (\hat{m}_1 \hat{n}_1)} \dots \\ & J_{23;4}^{(\hat{a}_l \hat{b}_l \hat{c}_l), (\hat{d}_l \hat{e}_l \hat{f}_l); (\hat{g}_k \hat{h}_k), (\hat{m}_k \hat{n}_k)} \overrightarrow{\partial}_{(\hat{d}_1 \hat{e}_1 \hat{f}_1) \dots (\hat{d}_l \hat{e}_l \hat{f}_l)} \overrightarrow{\partial}_{(\hat{m}_1 \hat{n}_1) \dots (\hat{m}_k \hat{n}_k)}, \\ F_{123;4} : & \overleftarrow{\partial}_{(\hat{a}_1 \hat{b}_1 \hat{c}_1) \dots (\hat{a}_l \hat{b}_l \hat{c}_l)} \overleftarrow{\partial}_{(\hat{g}_1 \hat{h}_1) \dots (\hat{g}_k \hat{h}_k)} \overleftarrow{\partial}_{\hat{p}_1 \dots \hat{p}_r} J_{123;4}^{(\hat{a}_1 \hat{b}_1 \hat{c}_1), (\hat{d}_1 \hat{e}_1 \hat{f}_1); (\hat{g}_1 \hat{h}_1), (\hat{m}_1 \hat{n}_1); \hat{p}_1 \hat{q}_1} \dots \\ & J_{123;4}^{(\hat{a}_l \hat{b}_l \hat{c}_l), (\hat{d}_l \hat{e}_l \hat{f}_l); (\hat{g}_k \hat{h}_k), (\hat{m}_k \hat{n}_k); \hat{p}_r \hat{q}_r} \overrightarrow{\partial}_{(\hat{d}_1 \hat{e}_1 \hat{f}_1) \dots (\hat{d}_l \hat{e}_l \hat{f}_l)} \overrightarrow{\partial}_{(\hat{m}_1 \hat{n}_1) \dots (\hat{m}_k \hat{n}_k)} \overrightarrow{\partial}_{\hat{q}_1 \dots \hat{q}_r}. \end{aligned}$$

4.7. Continuous limits of the fuzzy flag manifolds

As a consistency check, one can calculate the dimension of the continuous flag manifolds from considering the $L \rightarrow \infty$ limit of the various representations used in describing the fuzzy algebra of functions on them. The underlying idea is simply that given a cutoff L , the number of eigenvalues should be proportional to L^d on a d -dimensional manifold. The number of degrees of freedom in the matrix algebras was the square of the dimension $d(a_1(L), a_2(L), a_3(L))$ of the representation $(a_1(L), a_2(L), a_3(L))$, and we can thus deduce that

$$d = \lim_{L \rightarrow \infty} \frac{\ln(d(a_1(L), a_2(L), a_3(L)))}{\ln L}. \quad (4.71)$$

A trivial calculation shows that the representations and the matrix algebras we have chosen in the previous sections indeed reproduce the right dimensions for the various flag manifolds:

Flag manifold	$F_{1;4}$	$F_{2;4}$	$F_{3;4}$	$F_{12;4}$	$F_{13;4}$	$F_{23;4}$	$F_{123;4}$
representations	$(L, 0, 0)$	$(0, L, 0)$	$(0, 0, L)$	$(L, L, 0)$	$(L, 0, L)$	$(0, L, L)$	(L, L, L)
real dimension	6	8	6	10	10	10	12

Considering the expressions for the star products on the various flag manifolds found in the previous sections, it is also evident that the star or operator product will go over to the commutative product in the limit $L \rightarrow \infty$. As discussed in [32], this limit is, however, not clearly observable in the simplified formulæ using the complex coordinates a^i, b^i, c^i . Furthermore, the derivatives in the fuzzy case containing the geometric information are evidently approaching the derivatives in the continuum for $L \rightarrow \infty$.

5. Super Plücker embeddings and flag supermanifolds

5.1. Flag supermanifolds of $U(4|n)$

Flag supermanifolds can be defined analogously to bosonic flag manifolds by considering the supervector space $\mathbb{C}^{m|n}$ [18]; see also appendix A for more details. A *superflag* is a sequence of superspaces $V_{D_1} \subsetneq \dots \subsetneq V_{D_k} \subset \mathbb{C}^{m|n}$ such that $\dim_{\mathbb{C}} V_D = D = d|\delta$. Note that inclusion requires that $d_i \leq d_j$ and $\delta_i \leq \delta_j$ for $i < j$ with at least one inequality being strict. A *flag supermanifold* $F_{D_1 \dots D_k; m|n}$ is correspondingly the set of all superflags $f_{D_1 \dots D_k; m|n}$. One can again write a flag supermanifold as a coset space:

$$F_{D_1 \dots D_k; m|n} = U(m|n) / (U(m - d_k|n - \delta_k) \times \dots \times U(d_1|\delta_1)). \quad (5.1)$$

The special unitary supergroups $SU(m|n)$ are not useful here, as for $m = n$, one has to exclude the identity matrix, which is a central element in this case, from the set of generators of $\mathfrak{su}(m|n)$. See appendix A for more details on this point.

As before, we will be interested in the flag manifolds arising naturally in the double fibration underlying well-known supertwistor correspondences. These spaces are flags in the superspace $\mathbb{C}^{4|n}$, and because there are again natural projections, they fit into the following diagram:

$$\begin{array}{ccccc}
 & & F_{(2|0)(2|n)(3|n);(4|n)} & & \\
 & \swarrow & \uparrow & \searrow & \\
 F_{(2|0)(2|n);(4|n)} & & & & F_{(3|n);(4|n)} \\
 \uparrow & \swarrow & F_{(1|0)(2|0)(2|n)(3|n);(4|n)} & \searrow & \uparrow \\
 F_{(1|0)(2|0)(2|n);(4|n)} & & & & F_{(1|0)(3|n);(4|n)} \\
 & \swarrow & \downarrow & \searrow & \\
 & & F_{(1|0);(4|n)} & &
 \end{array} \quad (5.2)$$

Here, $F_{(1|0);(4|n)}$ is the superspace $\mathbb{C}P^{3|n}$ and $F_{(2|0)(2|n);(4|n)}$ is the conformal compactification of super Minkowski space with $n = \mathcal{N}$ being the number of supersymmetries. Note that $F_{(2|0)(2|n);(4|n)}$ contains the left chiral superspace $F_{(2|0);(4|n)}$ as well as the right chiral superspace $F_{(2|n);(4|n)}$ [18]. Since in the twistor correspondences involving $F_{(1|0);(4|n)}$ and $F_{(3|n);(4|n)}$, only these chiral subspaces play a rôle, we will also restrict the correspondence spaces and only consider $F_{(1|0)(2|0);(4|n)}$ and $F_{(2|n)(3|n);(4|n)}$ instead of $F_{(1|0)(2|0)(2|n);(4|n)}$ and $F_{(2|0)(2|n)(3|n);(4|n)}$.

To get a reliable handle on the geometry of the flag supermanifolds, it is useful to introduce local coordinates. For simplicity, we will first consider the ordinary Graßmannians and then discuss the super case. The extension to reducible flag supermanifolds will be straightforward. On a Graßmannian $G_{k;n}$, a patch corresponds to a subset $I \subset \{1, \dots, n\}$ with k elements, which selects k columns of a $k \times n$ matrix Z . We identify these columns with the columns of a $k \times k$ unit matrix; they fix parts of the vectors spanning the k -dimensional vector subspaces. The remaining columns are filled by the local coordinates on the patch I . As an example, consider $\mathbb{C}P^2 = G_{1;3}$, where there are three patches

$$Z_1 = (1 \ z_1^1 \ z_1^2), \quad Z_2 = (z_2^1 \ 1 \ z_2^2) \quad \text{and} \quad Z_3 = (z_3^1 \ z_3^2 \ 1). \quad (5.3)$$

Transition functions are elements of a finite subgroup of $GL(n, \mathbb{C})$, permuting the $k \times k$ unit

matrix to different columns. We also see that on every patch we have the $U(k) \times U(n-k) \subset U(n)$ invariance manifest.

For a super Graßmannian $G_{k|\kappa;n|\nu}$, we consider a $(k|\kappa) \times (k+(n-k)|\kappa+(\nu-\kappa))$ -dimensional supermatrix Z_I , into which we insert the columns of the $k|\kappa \times k|\kappa$ -dimensional unit matrix, preserving the grading of the matrix. The set I of the columns which we selected corresponds again to a patch. The discussion of the transition functions is the same as in the bosonic case and the symmetries factored out are again manifest.

A first observation is that for the super Graßmannians we are interested in, the fermionic dimension of the subspace is always either maximal or minimal. This leads e.g. to matrices Z_I

$$\begin{aligned} G_{2|0;4|4} : \quad Z_{34|\cdot} &= \left(\begin{array}{cc|c} x_{2 \times 2} & \mathbb{1}_{2 \times 2} & \xi_{2 \times 4} \end{array} \right), \\ G_{2|0;2|4} : \quad Z_{12|\cdot} &= \left(\begin{array}{c|c} \mathbb{1}_{2 \times 2} & \xi_{2 \times 4} \end{array} \right), \\ G_{2|4;4|4} : \quad Z_{34|1234} &= \left(\begin{array}{cc|cc} x_{2 \times 2} & \mathbb{1}_{2 \times 2} & 0_{2 \times 4} & \\ \xi_{4 \times 2} & 0_{4 \times 2} & \mathbb{1}_{4 \times 4} & \end{array} \right), \end{aligned} \quad (5.4)$$

where the lines indicate the boundaries of the four canonical blocks in the supermatrix Z . The first and the third space are the compactified, complexified chiral and anti-chiral $\mathcal{N} = 4$ superspaces, respectively. It is easy to convince oneself that (the bosonic part of) the transition functions are the same as in the purely bosonic case. Therefore, the super Graßmannian (and also the flag supermanifolds) we are dealing with are simply certain fermionic vector bundles over their bodies, i.e. the embedded ordinary flag manifolds. As an explicit example, consider the space $\mathbb{C}P^{1|2}$. Following our discussion of local coordinates, we introduce the two patches U_{\pm} , corresponding to the matrices

$$Z_+ = (z_+ \ 1 \ \zeta_+^1 \ \zeta_+^2) \quad \text{and} \quad Z_- = (1 \ z_- \ \zeta_-^1 \ \zeta_-^2). \quad (5.5)$$

The transition function between both patches is evidently $f_{+-} = z_+$, and therefore we have $\zeta_+^{1,2} = z_+ \zeta_-^{1,2}$. Thus, the space $\mathbb{C}P^{1|2}$ is the total space of the rank $0|2$ vector bundle

$$\Pi\mathcal{O}(1) \oplus \Pi\mathcal{O}(1) \rightarrow \mathbb{C}P^1. \quad (5.6)$$

For more details on the definition of super Graßmannians and flag supermanifolds, see [33].

For simplicity, we restrict our attention to the case $n = \mathcal{N} = 4$ in the following. Using the description of super Graßmannians given above, it is easy to determine the dimensions of the various flag supermanifolds in (5.2) for $n = 4$. We have

$$\begin{aligned} F_{(1|0);(4|4)} &: 3|4 & F_{(2|0);(4|4)} &: 4|8 \\ F_{(2|4);(4|4)} &: 4|8 & F_{(2|0)(2|4);(4|4)} &: 4|16 \\ F_{(1|0)(2|0);(4|4)} &: 5|8 & F_{(3|4);(4|4)} &: 3|4 \\ F_{(2|4)(3|4);(4|4)} &: 5|8 & F_{(1|0)(3|4);(4|4)} &: 5|16 \\ F_{(1|0)(2|0)(2|4)(3|4);(4|4)} &: 6|16. \end{aligned} \quad (5.7)$$

The minimal numbers of patches covering the flag supermanifolds are the same as the ones for their bodies.

5.2. The Plücker and super Plücker embeddings

Before discussing the super variant of the Plücker embedding, let us briefly recall its common form; see also the review article [34]. A Grassmannian $G_{k;n}$ is the space of all k -dimensional vector subspaces in $V = \mathbb{C}^n$. Each such space is spanned by a basis f_1, \dots, f_k and we can combine this basis into an element of the k -th exterior power of V :

$$A := f_1 \wedge \dots \wedge f_k, \quad A \in \Lambda^k V. \quad (5.8)$$

Each element of this form describes a point on $G_{k;n}$, however, not every element of $\Lambda^k V$ is of this form. In particular, a sum of two different such elements will not in general be decomposable into a single wedge product of k vectors. One thus needs additional conditions to decide, whether an element of $\Lambda^k V$ is fully decomposable. Physically, this question is completely analogous to the question, whether a k -fermion state can be decomposed into a product of k single particle states.

It is well known that the necessary and sufficient condition for $A \in \Lambda^k V$ to be of the form (5.8) is

$$(B \lrcorner A) \wedge A = 0 \quad \text{for all } B \in \Lambda^{k-1} V^\vee, \quad (5.9)$$

where V^\vee denotes the space dual to V . The proof of this statement is a simplification of the one in the graded case, which we will give below. The equations arising from (5.9) are called the *Plücker relations*. In the case $k = 2$, this condition simplifies to $(B \lrcorner A) \wedge A = \frac{1}{2} B \lrcorner (A \wedge A)$ and thus to $A \wedge A = \varepsilon_{ijkl} A^{ij} A^{kl} = 0$, the condition we used in section 2.3.

Consider now a basis (e_1, \dots, e_n) of V with a dual basis $(e_\vee^1, \dots, e_\vee^n)$ of V^* , $\langle e_i, e_\vee^j \rangle = \delta_i^j$. It is obviously sufficient to consider (5.9) only for elements B of the form $e_\vee^{i_1} \wedge \dots \wedge e_\vee^{i_k}$. Writing $A = A^{i_1 \dots i_k} e_{i_1} \wedge \dots \wedge e_{i_k}$, (5.9) reduces to

$$\sum_{t=0}^k (-1)^t A^{j_1 \dots j_{k-1} i_t} A^{i_1 \dots \widehat{i_t} \dots i_{k+1}} = A^{j_1 \dots j_{k-1} [i_1} A^{i_2 \dots i_{k+1}]} = 0, \quad (5.10)$$

where $\widehat{}$ indicates an omission as usual.

Note that not all of these equations are independent, but one can easily read off the number of independent ones. As the Plücker relations are evidently projective, one can fix a nonvanishing component to unity, e.g. $A^{p_1 \dots p_k} = 1$. Then it follows that one can solve for all coordinates with $m \geq 2$ indices different from all of the p_r [34]: Consider a sequence $q_1 \dots q_k$ of indices, m of which are not in the sequence $p_1 \dots p_k$. From (5.10), we obtain for $(j_k) = (q_1 \dots \widehat{q_r} \dots q_k)$ and $(i_k) = (q_r p_1 \dots p_k)$ the following equation:

$$A^{q_1 \dots \widehat{q_r} \dots q_k q_r} A^{p_1 \dots p_k} = \sum_{t=1}^k (-1)^t A^{q_1 \dots \widehat{q_r} \dots q_k p_t} A^{p_1 \dots \widehat{p_t} \dots p_k}. \quad (5.11)$$

If p_r is contained in the sequence $q_1 \dots q_k$, the right hand side vanishes, otherwise exactly $m-1$ of the elements in the sequence $q_1 \dots \widehat{q_r} \dots q_k p_t$ are not in the sequence $p_1 \dots p_k$. Iterating this prescription, we find that all Plücker coordinates can be expressed in terms of $A^{q_1 \dots q_k}$ with at most one q_s not in the sequence $p_1 \dots p_k$. That is, of the $\binom{n}{k}$ coordinates on $\Lambda^k \mathbb{C}^n$, only $1 + k(n-k)$ are relevant, and this is the number of (homogeneous) coordinates on the Grassmannian $G_{k;n}$.

For the discussion of flag supermanifolds, we need a similar picture at hands, and we will find that for Graßmannians $G_{k|\kappa;n|\nu}$ for which $\kappa \in \{0, \nu\}$, the Plücker embedding can be straightforwardly extended. This fact is mentioned in [33], but beyond this, we are not aware of any explicit discussion of super Plücker embeddings in the literature.

The Graßmannian $G_{k|\kappa;n|\nu}$ consists of spaces spanned by k even and κ odd supervectors in $\mathbf{V} = \mathbb{C}^{n|\nu}$. Given a basis $\mathbf{e}^I = (e^i, \varepsilon^\alpha)$, we can write a point¹⁰ $\mathbf{A} \in \Lambda^{k|\kappa}\mathbb{C}^{n|\nu} \supset G_{k|\kappa;n|\nu}$ as

$$\mathbf{A} = \mathbf{A}^{I_1 \dots I_k \Upsilon_1 \dots \Upsilon_\kappa} \mathbf{e}^{I_1} \wedge \dots \wedge \mathbf{e}^{I_k} \wedge \mathbf{e}^{\Upsilon_1} \wedge \dots \wedge \mathbf{e}^{\Upsilon_\kappa}, \quad (5.12)$$

where I_r and Υ_ρ range each from 1 to $n + \nu$. The Plücker relations we are looking for are supposed to be the necessary and sufficient conditions that

$$\mathbf{A} = \mathbf{f}_1 \wedge \dots \wedge \mathbf{f}_k \wedge \phi_1 \wedge \dots \wedge \phi_\kappa, \quad (5.13)$$

where \mathbf{f}_α and ϕ_α are linearly independent¹¹ even and odd supervectors, respectively. We now claim that for $\kappa \in \{0, \nu\}$ (and these are the only cases we are interested in), the necessary and sufficient condition reads as

$$(\mathbf{B} \lrcorner \mathbf{A}) \wedge \mathbf{A} = 0 \quad \text{for any } \mathbf{B} \in \Lambda^{k-1|\kappa}\mathbf{V}^\vee. \quad (5.14)$$

First, note that an even supervector \mathbf{v} with nonvanishing body \mathbf{v}° divides an element \mathbf{A} of $\Lambda^{k-1|\kappa}\mathbf{V}^\vee$ with $\mathbf{A}^\circ \neq 0$, i.e. $\mathbf{A} = \mathbf{v} \wedge \mathbf{w}$ for some $\mathbf{w} \in \Lambda^{k-2|\kappa}$ with $\mathbf{w}^\circ \neq 0$, if and only if $\mathbf{v} \wedge \mathbf{A} = 0$. Let us assume that $\kappa = 0$. If \mathbf{A} is of the form $\mathbf{f}_1 \wedge \dots \wedge \mathbf{f}_k$, then we can complete the \mathbf{f}_α to an (orthonormal) basis of \mathbf{V} by even supervectors $\mathbf{f}_{k+1} \dots \mathbf{f}_{n+\nu}$. If \mathbf{B} is composed only of \mathbf{f}_α^\vee with $\alpha \leq k$, (5.14) is satisfied. If there is one or more of the \mathbf{f}_α with $\alpha > k$, (5.14) is also true. Since (5.14) is linear in \mathbf{B} and the span of the cases we discussed comprises all of $\Lambda^{k-1|\kappa}\mathbf{V}^\vee$, (5.14) is true in general.

To prove the remaining direction, we follow the proof in the bosonic case [34] and explicitly construct the Graßmannian $G_{k|\kappa;n|\nu}$ from the coordinates $\mathbf{A}^{I_1 \dots I_k}$. First, note that we can again fix a bosonic coordinate which has nonvanishing body (at least one such component exists, if \mathbf{A} has nonvanishing body), say $\mathbf{A}^{i_1 \dots i_k} = 1$, since the equations are projective. Furthermore, all coordinates $\mathbf{A}^{J_1 \dots J_k}$ with a sequence $J_1 \dots J_k$ of more than one index different from $i_1 \dots i_k$ can again be written in terms of the remaining coordinates; the proof is the same as in the bosonic case. We now construct k vectors spanning a k -plane in $\Lambda^{k|0}\mathbb{C}^{n|\nu}$ by putting $p_m(J) = A^{i_1 \dots i_{m-1} J i_{m+1} \dots i_k}$, $m = 1 \dots k$, $J = 1 \dots n + \nu$. This evidently yields k linearly independent vectors with nonvanishing bodies, as they differ in the k components $J = i_l$: it is $p_m(i_l) = 0$ for $m \neq l$ and $p_l(i_l) = 1$. It remains to show that the plane corresponding to these vectors is indeed compatible with all the Plücker coordinates. First, it is straightforward to see that the components $A^{i_1 \dots i_{m-1} J i_{m+1} \dots i_k}$ all are compatible with our definition. As shown above, the remaining coordinates are derived from these and thus all the Plücker coordinates are the ones corresponding to the Graßmannian we constructed. Altogether, if the equation is satisfied, then the multivector \mathbf{A} describes a Graßmannian, which completes our proof for $\kappa = 0$.

¹⁰A discussion of wedge products of supervector spaces and their duals is found in [35].

¹¹See appendix A for a discussion of linear independence of supervectors.

For $\kappa = \nu$, it is sufficient to note that the wedge product of the ν odd supervectors in \mathbf{A} either vanishes, if they are linearly dependent, or spans all of the odd subspace. Therefore, it suffices again to focus on the even supervectors in \mathbf{A} , which is done by contracting with a dual multivector $\mathbf{B} \in \Lambda^{k-1|\nu}\mathbf{V}^\vee$, and there is nothing left to prove.

5.3. Plücker coordinates and projector description of irreducible flag supermanifolds

From the discussion in section 5.1., the description of $\mathbb{C}P^{3|4}$ is evident: homogeneous coordinates on $\mathbb{C}P^{3|4}$ are provided by the components of an even supervector (normalized and in a representation of type I, i.e. in a pure even basis) $\mathbf{a}^I = (a^i, \eta^\alpha)$. From the discussion above, we know that the space $\mathbb{C}P^{3|4}$ is the fermionic rank 4 vector bundle

$$\mathbb{C}^4 \otimes \Pi\mathcal{O}(1) \rightarrow \mathbb{C}P^3, \quad (5.15)$$

and its sections are given by homogeneous polynomials of degree one in the coordinates on $\mathbb{C}P^3$. We can therefore rewrite the fermionic components of the even supervector \mathbf{a}^I as

$$\eta^\alpha = \eta_i^\alpha a^i \quad \text{with} \quad \eta_i^\alpha \in \mathbb{C}^{0|8}. \quad (5.16)$$

In the following, however, we will not be interested in sections of this bundle but rather in the algebra of functions on its total space.

We therefore continue along the lines of the bosonic case and construct a projector using the supervector \mathbf{a} according to

$$\mathcal{P}_{1|0} = \mathbf{a}\bar{\mathbf{a}}^T. \quad (5.17)$$

Note that the description of a flag manifold using projectors trivially generalizes to the supercase. First of all, the bodies of even and odd supervectors of dimension $4|4$ have non-zero components in the first and the last four components, respectively. This property is preserved by the action of $U(4|4)$. Thus, given a projector

$$\mathcal{P} = \begin{pmatrix} \mathcal{P}_A & \mathcal{P}_B \\ \mathcal{P}_C & \mathcal{P}_D \end{pmatrix}, \quad (5.18)$$

we can read off the dimension $k|\kappa$ of the subspace of $\mathbb{C}^{4|4}$ onto which it projects to be $\text{tr}(\mathcal{P}_A^\circ) | \text{tr}(\mathcal{P}_D^\circ)$, where \cdot° denotes the body of the projector. We will define the rank of such a projector to be $k|\kappa$.

Also, there is a subgroup $U(4 - k|4 - \kappa)$ of $U(4|4)$, which leaves invariant a projector of rank $k|\kappa$. This is easily seen by the usual identification of $\mathfrak{u}(4|4)$ with $\mathfrak{u}(4+4)$ after combining the odd generators with an odd parameter and the fact that the rank of a projector is left invariant by the action of $U(4|4)$.

For $\mathcal{P}_{1|0}$ to be a projector, we have to demand that

$$(\mathbf{a}, \mathbf{a}) = \bar{\mathbf{a}}\mathbf{a} = \bar{a}^i a^i + i\bar{\eta}^\alpha \eta^\alpha = 1, \quad (5.19)$$

see appendix A for a definition of the scalar product of complex supervectors. This condition can also be understood from the construction of $\mathbb{C}P^{n|n+1}$ via

$$\mathbb{C}^{n+1|n+1} \rightarrow \mathcal{S}^{2n+1|2n+2} \rightarrow \mathbb{C}P^{n|n+1}, \quad (5.20)$$

as the first projection, see also [36]. We can in fact introduce coordinates $\mathbf{x}^{\hat{A}} = \bar{\mathbf{a}}^I \boldsymbol{\lambda}_{IJ}^{\hat{A}} \mathbf{a}^J$ in the superspace $\mathbb{R}^{32|32}$, where the $\boldsymbol{\lambda}^A$ are the generators of $U(4|4)$ as described in the appendix. The coordinates $\mathbf{x}^{\hat{A}}$ describe an embedding of $\mathbb{C}P^{3|4}$ in $\mathbb{R}^{32|32}$ and due to $\boldsymbol{\lambda}_{IJ}^{\hat{A}} \boldsymbol{\lambda}_{KL}^{\hat{B}} g_{\hat{A}\hat{B}} = \delta_{IL} \delta_{JK}$, we have

$$\mathcal{P}_{1|0} = \mathbf{x}^{\hat{A}} \boldsymbol{\lambda}^{\hat{B}} g_{\hat{A}\hat{B}} = \mathbf{a} \bar{\mathbf{a}}^T . \quad (5.21)$$

Underlying this construction is again a generalized Hopf fibration

$$1 \longrightarrow U(1) \longrightarrow S^{2n+1|2n+2} \longrightarrow \mathbb{C}P^{n|n+1} \longrightarrow 1 . \quad (5.22)$$

Let us now turn to the Graßmannians, whose description is rather straightforward using the super Plücker embedding. We start with $G_{2|0;4|4}$, and the Plücker coordinates are given by

$$\mathbf{A}_{2|0}^{\{IJ\}} = \mathbf{a}^{\{I} \mathbf{b}^{J\}} = \mathbf{a}^I \mathbf{b}^J - (-1)^{\tilde{I}\tilde{J}} \mathbf{a}^J \mathbf{b}^I = \mathbf{a}^I \mathbf{b}^J - \mathbf{b}^I \mathbf{a}^J , \quad (5.23)$$

where $\mathbf{a} = (a, \eta)$ and $\mathbf{b} = (b, \zeta)$ are two even supervectors, $\{\cdot, \cdot\}$ is the supercommutator and \tilde{I} denotes the parity of the corresponding index. That is, $\tilde{i} = 0$ and $\tilde{\alpha} = 1$, plus the parity of the supervector under consideration, modulo 2. In more detail, we have

$$\mathbf{A}_{2|0}^{ij} = a^{[i} b^{j]} , \quad \mathbf{A}_{2|0}^{i\alpha} = -\mathbf{A}_{2|0}^{\alpha i} = \frac{1}{2} (a^i \zeta^\alpha - \eta^\alpha b^i) \quad \text{and} \quad \mathbf{A}_{2|0}^{\alpha\beta} = \eta^{\{\alpha} \zeta^{\beta\}} , \quad (5.24)$$

where $\{\cdot\}$ denotes symmetrization, in particular of Graßmann-odd quantities. Note that this super-antisymmetrized combination of \mathbf{a} and \mathbf{b} indeed eliminates all components of \mathbf{b} parallel to \mathbf{a} and in the following, we will assume that \mathbf{b} and \mathbf{a} are perpendicular:

$$\bar{\mathbf{a}}^I \mathbf{b}^I = \bar{a}^i b^i + i \bar{\zeta}^\alpha \eta^\alpha = 0 . \quad (5.25)$$

Together with $\mathbf{a}^\circ \neq 0 \neq \mathbf{b}^\circ$, this equation implies that both supervectors are linearly independent. The (internal) stabilizer subgroup of $U(4|4)$ leaving $\mathbf{A}_{2|0}^{IJ}$ invariant is $U(2|0)$, which rotates the supervector \mathbf{a} into \mathbf{b} and vice versa; see the discussion of the fuzzy case for more details.

We observed before that the Plücker coordinates A_2^{ij} on $G_{2;4}$ contain some redundancy: first there is a scaling, which renders them effectively coordinates on $\mathbb{C}P^5$, and second, there is the identity $\varepsilon_{ijkl} A_2^{ij} A_2^{kl} = 0$. In the present case, the redundancy is somewhat larger, but from the discussion of the super Plücker embedding, we can be specific about the number of redundant coordinates. First of all, we have 16 even and 16 odd homogeneous coordinates and thus we are using a Plücker embedding into $\mathbb{C}P^{15|16}$. Assuming that $\mathbf{A}_{2|0}^{12} = 1$ fixes the scaling, only the coordinates

$$\begin{array}{ccc} \mathbf{A}_{2|0}^{13} & \mathbf{A}_{2|0}^{14} & \mathbf{A}_{2|0}^{15} \quad \dots \quad \mathbf{A}_{2|0}^{18} \\ \mathbf{A}_{2|0}^{23} & \mathbf{A}_{2|0}^{24} & \mathbf{A}_{2|0}^{25} \quad \dots \quad \mathbf{A}_{2|0}^{28} \end{array} \quad (5.26)$$

are independent and we thus arrive at a 4|8-dimensional space, $G_{2|0;4|4}$.

From the bi-supervector $\mathbf{A}_{2|0}$, one can again construct a projector:

$$(\mathcal{P}_{2|0;4|4})^{IJ;KL} = \mathbf{A}_{2|0}^{IJ} \bar{\mathbf{A}}_{2|0}^{KL} . \quad (5.27)$$

With our choice and the orthonormality of \mathbf{a} and \mathbf{b} , $\mathcal{P}_{2|0;4|4}$ indeed satisfies $(\mathcal{P}_{2|0;4|4})^2 = \mathcal{P}_{2|0;4|4}$. Underlying the construction of this projector is again a generalized Hopf fibration, which is evidently a superextension of the one for $G_{2;4}$:

$$1 \longrightarrow \mathrm{U}(2|0) \longrightarrow S^{7|8} \times S^{5|8} \longrightarrow F_{2|0;4|4} \longrightarrow 1. \quad (5.28)$$

It is now necessary to introduce a super-antisymmetrized tensor product defined as

$$\begin{aligned} (\mathbf{A} \mathbin{\frown} \mathbf{B})_{IJ;KL} &:= \frac{1}{4} (\mathbf{A}_{IK} \mathbf{B}_{JL} - (-1)^{\tilde{I}\tilde{J}} \mathbf{A}_{JK} \mathbf{B}_{IL} - \\ &\quad (-1)^{\tilde{K}\tilde{L}} \mathbf{A}_{IL} \mathbf{B}_{JK} + (-1)^{\tilde{I}\tilde{J}} (-1)^{\tilde{K}\tilde{L}} \mathbf{A}_{JL} \mathbf{B}_{IK}). \end{aligned} \quad (5.29)$$

As in the case of the antisymmetric product of the generators of $\mathrm{U}(4)$, we also have here various Fierz identities, see appendix A. Using one of these identities, we can introduce the projector

$$\begin{aligned} (\mathcal{P}_{2|0;4|4})_{IJ;KL} &= \bar{\mathbf{A}}_{2|0}^{KL} \mathbf{A}_{2|0}^{IJ} \\ &= \bar{\mathbf{A}}_{2|0}^{K'L'} (\boldsymbol{\lambda}^{\hat{A}} \mathbin{\frown} \boldsymbol{\lambda}^{\hat{B}})_{K'L'I'J'} \mathbf{A}_{2|0}^{I'J'} (\boldsymbol{\lambda}^{\hat{C}} \mathbin{\frown} \boldsymbol{\lambda}^{\hat{D}})_{IJ;KL} g_{AC} g_{BD} \\ &=: \mathbf{x}^{\hat{A}\hat{B}} (\boldsymbol{\lambda}^{\hat{C}} \mathbin{\frown} \boldsymbol{\lambda}^{\hat{D}})_{IJ;KL} g_{AC} g_{BD}. \end{aligned} \quad (5.30)$$

The next Graßmannian to be described is the space $G_{2|4;4|4}$. This space is certainly “dual” to $G_{2|0;4|4}$, as one easily guesses from the description in terms of coordinate matrices Z_I given in the previous section. To describe this space, one needs to take two even and four odd normalized supervectors and then super-antisymmetrize them. A short remark is in order to show that super-antisymmetrizing indeed yields a projection on their mutually orthogonal components. We saw above that this is true for two even supervectors. Since the bodies of normalized even and odd supervectors are non-vanishing exactly in the even and odd indices, respectively, a pair of an even and an odd normalized supervector are always linearly independent; antisupersymmetrization just eliminates redundancies in the description of 1|1-dimensional subspaces via Plücker relations. Given two odd supervectors, super-antisymmetrization has the same effect on the even and odd components as it had on two even supervectors and thus projects out non-orthogonal components.

Altogether, we have Plücker coordinates

$$\mathbf{A}_{2|4}^{IJ\Upsilon_1 \dots \Upsilon_4} = \mathbf{a}^I \mathbf{b}^J \boldsymbol{\eta}^{\Upsilon_1} \dots \boldsymbol{\eta}^{\Upsilon_4}, \quad (5.31)$$

where \mathbf{a}^I , \mathbf{b}^I and $\boldsymbol{\eta}^\Upsilon$ are even and odd supervectors, respectively. We assume that $\mathbf{A}_{2|4}^{1245678} = 1$ to fix the scale. From the discussion of the super Plücker embedding, it then follows that the independent coordinates here are given by

$$\begin{array}{ccc} \mathbf{A}_{2|4}^{135678} & \mathbf{A}_{2|4}^{145678} & \mathbf{A}_{2|4}^{155678} \quad \dots \quad \mathbf{A}_{2|4}^{185678} \\ \mathbf{A}_{2|4}^{235678} & \mathbf{A}_{2|4}^{245678} & \mathbf{A}_{2|4}^{255678} \quad \dots \quad \mathbf{A}_{2|4}^{285678} \end{array}. \quad (5.32)$$

In the case of the dual complex projective superspace $\mathbb{C}P_*^{3|4}$, we add to this picture another even supervector \mathbf{c}^I and obtain the Plücker coordinates

$$\mathbf{A}_{2|4}^{IJK\Upsilon_1 \dots \Upsilon_4} = \mathbf{a}^I \mathbf{b}^J \mathbf{c}^K \boldsymbol{\eta}^{\Upsilon_1} \dots \boldsymbol{\eta}^{\Upsilon_4}, \quad (5.33)$$

where $\mathbf{a}^I, \mathbf{b}^I$ and $\boldsymbol{\eta}^{\Upsilon}$ are even and odd supervectors, respectively. We fix the scale by $\mathbf{A}_{3|4}^{12345678} = 1$ and the remaining independent coordinates here are

$$\mathbf{A}_{3|4}^{1245678} \quad \mathbf{A}_{3|4}^{1345678} \quad \mathbf{A}_{3|4}^{2345678} \quad \mathbf{A}_{3|4}^{155678} \quad \dots \quad \mathbf{A}_{2|4}^{185678} \quad . \quad (5.34)$$

We refrain from going into any further detail at this point and refer to the discussion of the fuzzy pictures of these super Grassmannians.

5.4. The reducible flag supermanifolds

As in the bosonic case, the description of the reducible flag supermanifolds is merely a combination of the underlying ‘‘elementary’’ Grassmannians.

The complexified, compactified super Minkowski space $F_{(2|0)(2|4);(4|4)}$ is a reducible flag supermanifold, although its body is an irreducible flag manifold. Combining the sets of Plücker coordinates on $G_{2|0;4|4}$ and $G_{2|4;4|4}$ and factoring out redundancies arising from the fact that the even supervectors in both cases are the same, one arrives at the Plücker coordinates on the space $F_{(2|0)(2|4);4|4}$, which is of superdimension $4|16$.

Similarly, the remaining flag supermanifolds are constructed by combining the coordinates of the Grassmannians corresponding to the various subflags. Again, a more detailed discussion will be presented, when we develop the fuzzy versions of these spaces.

5.5. Geometric structures on the flag supermanifolds

The description of the flag supermanifolds in terms of projectors allows us to proceed similarly to the case of ordinary flag manifolds in the description of their geometry. That is, we describe a point on a flag supermanifold \mathbf{M} again by a projector \mathcal{P}_0 , and the action of $\mathbf{G} = \mathbf{U}(4|4)$ on this point produces all of \mathbf{M} . The coordinates \mathbf{x}^A provide an embedding of \mathbf{M} in $\mathbb{R}^{m|n}$ (e.g. $\mathbb{C}P^{3|4}$ is embedded in $\mathbb{R}^{32|32}$). The tangent vectors at \mathcal{P}_0 are naturally obtained from an appropriate action $R(\Lambda)$ on \mathcal{P}_0 and we define analogously to the bosonic case

$$T_{\mathcal{P}_0}\mathbf{M} = \{R(\Lambda)\mathcal{P}_0 | \Lambda \in \mathfrak{u}(4|4)\} \quad . \quad (5.35)$$

The generators of the subgroup \mathbf{H} in $\mathbf{M} = \mathbf{G}/\mathbf{H}$ will leave \mathcal{P}_0 invariant, and thus $T_{\mathcal{P}_0}\mathbf{M}$ is of the same dimension as \mathbf{M} . In the case of $\mathbb{C}P^{3|4}$, $R(\Lambda)$ is the superadjoint action, and since \mathcal{P}_0 is an even supermatrix, it is thus clear that a vector $\mathbf{V} \in T_{\mathcal{P}_0}\mathbf{M}$ satisfies

$$\{\mathcal{P}_0, \mathbf{V}\} = \mathbf{V} \quad \text{and} \quad \text{str } \mathbf{V} = 0 \quad . \quad (5.36)$$

From this point, it is rather obvious that we can proceed with the definition of the complex structure, the metric and the Kähler structure exactly as in the bosonic case. We thus go over to an arbitrary projector \mathcal{P} describing any point on \mathbf{M} and define the complex structure as

$$I\mathbf{V} := -i\{\mathcal{P}, \mathbf{V}\} = -i[\mathcal{P}, \mathbf{V}] \quad , \quad (5.37)$$

which trivially satisfies $I^2 = -\mathbb{1}$ and the hermitian supermetric as

$$g(\mathbf{V}_1, \mathbf{V}_2) = -\text{str}(I\mathbf{V}_1 I\mathbf{V}_2) \quad , \quad (5.38)$$

which satisfies $g(I\mathbf{V}_1, I\mathbf{V}_2) = g(\mathbf{V}_1, \mathbf{V}_2)$. The latter is invariant under the action of $U(4|4)$ on the vectors \mathbf{V}_1 and \mathbf{V}_2 and defines a supersymmetric even tensor:

$$g(\mathbf{V}_1, \mathbf{V}_2) = (-1)^{\tilde{\mathbf{V}}_1 \tilde{\mathbf{V}}_2} g(\mathbf{V}_2, \mathbf{V}_1). \quad (5.39)$$

Evidently, there is the supersymplectic structure

$$\Omega(\mathbf{V}_1, \mathbf{V}_2) = g(I\mathbf{V}_1, \mathbf{V}_2), \quad (5.40)$$

satisfying $\Omega(\mathbf{V}_1, \mathbf{V}_2) = -(-1)^{\tilde{\mathbf{V}}_1 \tilde{\mathbf{V}}_2} \Omega(\mathbf{V}_2, \mathbf{V}_1)$, from which we obtain the super Kähler structure J as

$$J(\mathbf{V}_1, \mathbf{V}_2) = \frac{1}{2}(g(\mathbf{V}_1, \mathbf{V}_2) + i\Omega(\mathbf{V}_1, \mathbf{V}_2)) = \text{str}(\mathcal{P}\mathbf{V}_1(\mathbb{1} - \mathcal{P})\mathbf{V}_2). \quad (5.41)$$

Note that for any normalized supervector \mathbf{a} , there is a transformation $\mathbf{g} \in U(4|4)$ mapping it to the vector $\mathbf{a}' = (1, 0, \dots)^T$. This implies that by an appropriate action of \mathbf{g} , one can turn \mathcal{P} into $\text{diag}(1, 0, \dots)$ and therefore we have again the formula $\text{str}(\mathcal{P}\mathbf{A}\mathcal{P}\mathbf{B}) = \text{str}(\mathcal{P}\mathbf{A})\text{str}(\mathcal{P}\mathbf{B})$, which allows us to write

$$J(\mathbf{V}_1, \mathbf{V}_2) = \text{str}(\mathcal{P}\mathbf{V}_1\mathbf{V}_2) - \text{str}(\mathcal{P}\mathbf{V}_1)\text{str}(\mathcal{P}\mathbf{V}_2). \quad (5.42)$$

Furthermore, we introduce the obvious components $J^{AB} = J(\boldsymbol{\lambda}^A, \boldsymbol{\lambda}^B)$ etc., for which we have e.g. the identity

$$J^{AB}\boldsymbol{\lambda}_B = \text{str}(\mathcal{P}\boldsymbol{\lambda}^A(\mathbb{1} - \mathcal{P})\boldsymbol{\lambda}^B)\boldsymbol{\lambda}_B = \mathcal{P}\boldsymbol{\lambda}^A(\mathbb{1} - \mathcal{P}). \quad (5.43)$$

Note that a projection onto the body of all the structures introduced in this section naturally reduces them to their ordinary counterparts on bosonic flag manifolds. These geometric structures are thus (unique) supersymmetric extensions. Furthermore, the above discussion naturally extends to the case of all other flag supermanifolds involving super-antisymmetrized tensor products of the type $\boldsymbol{\lambda}^A \frown \boldsymbol{\lambda}^B \frown \dots$.

6. Fuzzy flag supermanifolds

Having a description of flag supermanifolds using the Plücker embedding, we obtain quite straightforwardly the description of fuzzy flag supermanifolds. We will be rather concise and essentially stress the differences with the bosonic case.

6.1. Supercoherent states

The discussion of supercoherent states is done in close analogy to the case of bosonic coherent states. Consider the generators of the supergroup $U(4|4)$. After taking out the matrix $\text{diag}(\mathbb{1}_4, -\mathbb{1}_4)$, we are left with 31 bosonic and 32 fermionic generators. There are seven generators of the Cartan subalgebra: the six Cartan generators of the two $SU(4)$ s contained in $U(4|4)$ together with $\mathbb{1}_8$. We pair the remaining 24 bosonic and 30 fermionic generators into raising and lowering operators. Seven of them are fundamental, the remaining ones are

generated by supercommutators of these. All of the fundamental and Cartan generators are of the form

$$H_i = \begin{pmatrix} \ddots & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & \ddots \end{pmatrix}, \quad E_i^+ = \begin{pmatrix} \ddots & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \ddots \end{pmatrix}, \quad E_i^- = \begin{pmatrix} \ddots & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & \ddots \end{pmatrix},$$

except for the following three:

$$H_4 = \left(\begin{array}{ccc|cc} \ddots & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & \ddots & \end{array} \right), \quad E_4^+ = \left(\begin{array}{ccc|cc} \ddots & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \ddots & \end{array} \right), \quad E_4^- = \left(\begin{array}{ccc|cc} \ddots & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & \ddots & \end{array} \right).$$

The lines mark the boundaries between the four blocks of the supermatrices.

Correspondingly, we can introduce a super Dynkin diagram of the form

$$\begin{array}{ccccccc} a_1 & & a_2 & & a_3 & & q_0 & & a_4 & & a_5 & & a_6 \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \otimes & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array} \quad (6.1)$$

where the Dynkin labels indicate again the number of nontrivial actions of E_i^- on a highest weight state. Setting these labels to zero, we evidently enlarge the isotropy group of the highest weight states in the same way as in the bosonic case. That is, the highest weight state in the representation $(L, 0, 0, 0, 0, 0, 0)$ has isotropy group $\text{SU}(3|4)$, while the highest weight state in the representation $(0, L, 0, 0, 0, 0, 0)$ has isotropy group $\text{SU}(2) \times \text{SU}(2|4)$. These are thus the representations, in which the coherent states correspond to points on the flag supermanifolds $F_{1|0;4|4}$ and $F_{2|0;4|4}$.

The remaining representations for the flag manifolds $F_{3|4;4|4}$ and $F_{2|4;4|4}$ are given by representations with the Dynkin labels $(0, 0, 0, 0, 0, 0, L)$ and $(0, 0, 0, 0, 0, L, 0)$. The representations corresponding to flag manifolds are derived by choosing the Dynkin labels corresponding to all the contained Graßmannians to be non-vanishing.

The further discussion of the construction of coherent states as well as the treatment of the various patches (choosing dominant weight states instead of highest weight states) is evident. Instead of going into details, we continue directly with the construction of fuzzy matrix algebras on the flag supermanifolds in the next section.

6.2. The fuzzy complex projective superspace $\mathbb{C}P_F^{3|4}$

The fuzzification of $\mathbb{C}P^{3|4}$ is obtained by promoting its homogeneous coordinates to creation and annihilation operators of bosonic and fermionic harmonic oscillators:

$$\mathbf{a}^I = (a^i, \eta^\alpha) \quad \rightarrow \quad \hat{\mathbf{a}}^I = (\hat{a}^i, \hat{\eta}^\alpha) \quad \text{with} \quad \{\mathbf{a}^I, \mathbf{a}^{J\dagger}\} = \delta^{IJ}. \quad (6.2)$$

The total number operator reads then as $\hat{\mathbf{N}} = \hat{\mathbf{a}}^{I\dagger} \hat{\mathbf{a}}^I = \hat{a}^{i\dagger} \hat{a}^i + \hat{\eta}^{\alpha\dagger} \hat{\eta}^\alpha$, and commutes with the auxiliary coordinate operator

$$\hat{\mathbf{x}}^A := \frac{1}{\hat{\mathbf{N}}} \hat{\mathbf{a}}^{I\dagger} \lambda_{IJ}^A \hat{\mathbf{a}}^J. \quad (6.3)$$

We can thus restrict the algebra of functions to the L -particle Hilbert space $\mathcal{H}_{1|0;4|4}^F$ spanned by the states

$$\hat{\mathbf{a}}^{I_1^\dagger} \dots \hat{\mathbf{a}}^{I_L^\dagger} |0\rangle \quad \text{or} \quad \hat{a}^{i_1^\dagger} \dots \hat{a}^{i_{L-k}} \hat{\eta}^{\alpha_1^\dagger} \dots \hat{\eta}^{\alpha_k^\dagger} |0\rangle , \quad (6.4)$$

where, evidently, $k \leq 4$. Using the Schwinger construction for Lie superalgebras, we can define an action of $\mathfrak{u}(4|4)$ on this space:

$$\hat{\mathbf{L}}^A = \hat{\mathbf{a}}^{J^\dagger} \lambda_{IJ}^A \hat{\mathbf{a}}^J . \quad (6.5)$$

In the standard notation¹² for Young supertableaux, the representations space $\mathcal{H}_{1|0;4|4}^F$ corresponds to

$$\overbrace{\begin{array}{|c|c|c|c|} \hline \bullet & \bullet & \bullet & \bullet \\ \hline \end{array}}^L . \quad (6.6)$$

The algebra of functions on $\mathbb{C}P^{3|4}$ is given by the matrix algebra

$$\hat{\mathbf{a}}^{I_1^\dagger} \dots \hat{\mathbf{a}}^{I_L^\dagger} |0\rangle \langle 0| \hat{\mathbf{a}}^{J_1} \dots \hat{\mathbf{a}}^{J_L} , \quad (6.7)$$

which corresponds to the tensor product

$$\overbrace{\begin{array}{|c|c|c|c|} \hline \bullet & \bullet & \bullet & \bullet \\ \hline \end{array}}^L \otimes \overbrace{\begin{array}{|c|c|c|c|} \hline / & / & / & / \\ \hline \end{array}}^L , \quad (6.8)$$

where \square stands for the dual (contragredient) representation of \square .

For simplicity, let us give the star product on this space only for the complex coordinates $\mathbf{a}^I = (a^i, \eta^\alpha)$. The algebra of functions is spanned by the monomials

$$\bar{\mathbf{a}}^{I_1} \dots \bar{\mathbf{a}}^{I_L} \mathbf{a}^{J_1} \dots \mathbf{a}^{J_L} , \quad (6.9)$$

and the star product is defined as

$$(\mathbf{f} \star \mathbf{g}) = \mu \left[\frac{1}{L!} \frac{\partial}{\partial \mathbf{a}^{I_1}} \dots \frac{\partial}{\partial \mathbf{a}^{I_L}} \otimes \frac{1}{L!} \frac{\partial}{\partial \bar{\mathbf{a}}^{I_1}} \dots \frac{\partial}{\partial \bar{\mathbf{a}}^{I_L}} (\mathbf{f} \otimes \mathbf{g}) \right] . \quad (6.10)$$

Although the second order Casimir still labels representations to some extent, the Laplacian in the continuum does not have any immediate meaning. We are thus more interested in translating all the various derivatives, written in terms of the embedding coordinates to the fuzzy picture. This is easily done using the generators \mathcal{L}_A described in appendix B. One can show in complete analogy to the case of $\mathbb{C}P^3$ that

$$\mathcal{L}_A f_L(\mathbf{x}) = \frac{L}{\sqrt{2}} \left(\mathbf{x}_A \star f_L(\mathbf{x}) - (-1)^{\tilde{A}\tilde{L}} \tilde{f}_L f_L(\mathbf{x}) \star \mathbf{x}_A \right) = \text{tr} \left(\rho^L(\mathbf{x}_{1|0;4|4}) \{ \hat{\mathcal{L}}_A, \hat{f} \} \right) . \quad (6.11)$$

Together with the Killing metric g_{AB} , this sufficiently describes the geometry on fuzzy $\mathbb{C}P^{3|4}$ embedded in $\mathbb{R}^{32|32}$.

¹²see e.g. [37, 38]

6.3. The remaining fuzzy Grassmannian supermanifolds

The next flag manifold in our list is $G_{2|0;4|4}$, which is of complex dimension $4|8$. From the super Plücker embedding of this space into $\Lambda^{2|0}\mathbb{C}^{4|4}$ (see (5.12) for a definition of this space), the fuzzification is immediately obvious. We start from two sets of supersymmetric oscillators

$$\{\hat{\mathbf{a}}^I, \hat{\mathbf{a}}^{J\dagger}\} = \delta^{IJ}, \quad \{\hat{\mathbf{b}}^I, \hat{\mathbf{b}}^{J\dagger}\} = \delta^{IJ} \quad (6.12)$$

with components $\hat{\mathbf{a}}^I = (\hat{a}^i, \hat{\eta}^\alpha)$ and $\hat{\mathbf{b}}^I = (\hat{b}^i, \hat{\theta}^\alpha)$. From these, we construct the composite annihilation and creation operators

$$\hat{\mathbf{A}}_{2|0}^{IJ} := \hat{\mathbf{a}}^{\{I} \hat{\mathbf{b}}^{J\}}, \quad \hat{\mathbf{A}}_{2|0}^{IJ\dagger} := \hat{\mathbf{a}}^{\{I\dagger} \hat{\mathbf{b}}^{J\dagger\}}, \quad (6.13)$$

or, in more detail,

$$\hat{\mathbf{A}}_{2|0}^{ij} = \hat{a}^{[i} \hat{b}^{j]}, \quad \hat{\mathbf{A}}_{2|0}^{i\alpha} = \hat{a}^i \hat{\theta}^\alpha - \hat{\eta}^\alpha \hat{b}^i, \quad \hat{\mathbf{A}}_{2|0}^{\alpha i} = -\hat{\mathbf{A}}_{2|0}^{i\alpha}, \quad \hat{\mathbf{A}}_{2|0}^{\alpha\beta} := \hat{\eta}^{\{\alpha} \hat{\theta}^{\beta\}}, \quad (6.14)$$

together with their hermitian conjugates. The construction of the usual L -particle Hilbert space is now again straightforward and this space forms a representation of $U(4|4)$. Combining this Hilbert space with a dual copy, we obtain the algebra of functions on $G_{2|0;4|4}^F$ as a matrix algebra of the form

$$\begin{array}{|c|c|c|c|} \hline & L & & \\ \hline \bullet & \bullet & \bullet & \bullet \\ \hline \bullet & \bullet & \bullet & \bullet \\ \hline \bullet & \bullet & \bullet & \bullet \\ \hline \bullet & \bullet & \bullet & \bullet \\ \hline \end{array} \otimes \begin{array}{|c|c|c|c|} \hline & L & & \\ \hline / & / & / & / \\ \hline / & / & / & / \\ \hline / & / & / & / \\ \hline / & / & / & / \\ \hline \end{array} \quad (6.15)$$

From this construction, the deformed algebra of functions on $G_{2|0;4|4}$ together with the star product are also obvious.

The construction for the remaining two Grassmannians $G_{2|4;4|4}$ and $G_{3|4;4|4} = \mathbb{C}P_*^{3|4}$ uses composite creation and annihilation operators obtained from super-antisymmetrizing 2 even and 4 odd, and 3 even and 4 odd sets of superoscillators, respectively:

$$\hat{\mathbf{A}}_{2|4}^{IJ\Upsilon_1 \dots \Upsilon_4} = \hat{\mathbf{a}}^{\{I} \hat{\mathbf{b}}^J \hat{\eta}_1^{\Upsilon_1} \dots \hat{\eta}_4^{\Upsilon_4\}} \quad \text{and} \quad \hat{\mathbf{A}}_{3|4}^{IJK\Upsilon_1 \dots \Upsilon_4} = \hat{\mathbf{a}}^{\{I} \hat{\mathbf{b}}^J \hat{\mathbf{c}}^K \hat{\eta}_1^{\Upsilon_1} \dots \hat{\eta}_4^{\Upsilon_4\}}, \quad (6.16)$$

where the $\hat{\eta}^\Upsilon = (\hat{\eta}^i, \hat{a}^\alpha)$ are sets of odd annihilation operators satisfying with their hermitian conjugate the algebra

$$\{\hat{\eta}^{\Upsilon_1}, \hat{\eta}^{\Upsilon_2\dagger}\} = \delta^{\Upsilon_1 \Upsilon_2}. \quad (6.17)$$

From the discussion above, the representations are clear.

6.4. The fuzzy reducible flag supermanifolds

The discussion of the fuzzy reducible flag supermanifolds is now completely obvious. By combining sets of oscillators from the various super Grassmannians, we obtain the appropriate sets of oscillators (and thus the relevant Fock spaces from which the algebras of functions are constructed) for the flag supermanifolds naturally projecting on these Grassmannians. Instead of repeating the discussion for all the flag supermanifolds, let us merely study the example of $F_{(1|0)(2|0);4|4}$.

For this flag supermanifold, we need the oscillators of $G_{1|0;4|4} = \mathbb{C}P^{3|4}$ together with the ones of $G_{2|0;4|4}$:

$$\hat{\mathbf{a}}^I, \quad \hat{\mathbf{a}}^{I\dagger}, \quad \hat{\mathbf{A}}_{2|0}^{IJ} = \hat{\mathbf{a}}^{\{I} \hat{\mathbf{b}}^{J\}}, \quad \hat{\mathbf{A}}_{2|0}^{IJ\dagger} = \hat{\mathbf{a}}^{\{I\dagger} \hat{\mathbf{b}}^{J\dagger\}}. \quad (6.18)$$

Using L_1 operators $\hat{\mathbf{a}}^{I\dagger}$ and L_2 operators $\hat{\mathbf{A}}_{2|0}^{JJ\dagger}$ we construct the (L_1, L_2) -particle Fock space and its dual. Tensoring them yields the algebra of functions on $F_{(1|0)(2|0);4|4}$ as the matrix algebra

$$\begin{array}{|c|c|c|c|c|c|c|c|} \hline & & & & & & & \\ \hline \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \hline \bullet & \bullet & \bullet & \bullet & & & & \\ \hline \end{array} \otimes \begin{array}{|c|c|c|c|c|c|c|c|} \hline & & & & & & & \\ \hline / & / & / & / & / & / & / & / \\ \hline / & / & / & / & / & / & / & / \\ \hline \end{array} . \quad (6.19)$$

6.5. Fuzzy Calabi-Yau supermanifolds

Calabi-Yau supermanifolds have received much attention recently in twistor string theory [39], where $\mathbb{C}P^{3|4}$ was used as a target space for the topological B-model. The interest in this particular space is due to the fact that $\mathbb{C}P^{3|4}$ is simultaneously a supertwistor space and a Calabi-Yau supermanifold. The latter spaces are defined as spaces whose canonical bundle is trivial and thus have a nowhere vanishing holomorphic volume form. It was furthermore conjectured [39, 40] that there is a mirror symmetry between $\mathbb{C}P^{3|4}$ and the superambitwistor space $\mathcal{L}^{5|6} := F_{(1|0)(3|3);4|3}$. (Note that in our above constructions, we instead considered the space $F_{(1|0)(3|4);4|4}$.) The space $\mathcal{L}^{5|6}$ is of real dimension $10|12$ and a coset space of $U(4|3)$, as defined in (5.1). The corresponding fuzzy space is obtained from merging the fuzzy versions of $\mathbb{C}P^{3|3} = F_{1|0;4|3}$ and $\mathbb{C}P_*^{3|3} = F_{3|3;4|3}$ in the same way fuzzy flag manifolds are obtained from their sub-Graßmannians. Since the construction is again rather trivial, we stop here and postpone the analysis of fuzzy mirror symmetry to future work.

Acknowledgements

We would like to thank Charles Nash and Sachin Vaidya for helpful comments. In particular, we wish to express our gratitude towards Brian Dolan, Denjoe O’Connor and Peter Prešnajder for numerous useful discussions and for valuable remarks on a draft of this paper.

Appendix

A. Supermathematics, conventions and definitions

We denote even objects by Latin letters and odd objects by Greek ones. Boldface symbols will represent superobjects. Furthermore, a tilde over an index or an object will denote the naturally assigned parity, and \cdot° attached to an object refers to its body.

Supernumbers

A *supernumber* is an element of the Graßmann algebra Λ_N , $N \in \mathbb{N} \cup \{\infty\}$ which has generators ξ^i , $i = 1, \dots, N$ satisfying $\xi^i \xi^j + \xi^j \xi^i = 0$. The Graßmann algebra decomposes into an even and an odd part, $\Lambda_{N,0} = \Lambda_{N,c}$, $\Lambda_{N,1} = \Lambda_{N,a}$, which are the subsets of supernumbers built from an even and an odd number of Graßmann generators, respectively. The *body* of a supernumber is denoted by z° and consists of the purely complex part of z containing no Graßmann generator. For *complex conjugation of Graßmann odd quantities*, there are essentially two

conventions used throughout the literature. First, and most commonly, there is

$$(\theta_1\theta_2)^* = \bar{\theta}_2\bar{\theta}_1 = -\bar{\theta}_1\bar{\theta}_2, \quad (\text{A.1})$$

which is used e.g. in [38, 41]. Second, there is

$$(\theta_1\theta_2)^* = -\bar{\theta}_2\bar{\theta}_1 = +\bar{\theta}_1\bar{\theta}_2, \quad (\text{A.2})$$

which is used in [42, 35]. The latter convention respects the sign rule that interchanging two Graßmann-odd objects in a monomial should always be accompanied by an additional sign. There is a discussion of this issue in [42]. Manin in his book [33] also discusses all of these conventions. In this paper, we use the second convention.

Supervectors

A *supervector space* is a free module over a supercommutative ring. We restrict our considerations to supervector spaces which are endowed with a so-called pure basis. In particular, consider a supervector space V with a so-called class I (even) basis (\mathbf{e}^A) of n even and ν odd elements: $(\mathbf{e}^A) = (e^1, \dots, e^n, \varepsilon^1, \dots, \varepsilon^\nu)$. The supervector space V is then said to be of dimension $n|\nu$.

A $n|\nu$ -dimensional *supervector* consists of $n + \nu$ components. If the first n of the components of a supervector are even and the remaining ν odd, the supervector is called even. If the inverse statement is true, the supervector is said to be odd:

$$\mathbf{x} = x^a e^a + \xi^\alpha \varepsilon^\alpha \Rightarrow \tilde{\mathbf{x}} = 0 \quad \text{and} \quad \mathbf{x} = \xi^a e^a + x^\alpha \varepsilon^\alpha \Rightarrow \tilde{\mathbf{x}} = 1, \quad (\text{A.3})$$

where x^i and ξ^i are complex even and odd supernumbers, respectively. If the supervector is neither odd nor even, it is of mixed parity.

We will also allow for class II bases, in which the parity of the even and odd basis elements is interchanged: $(\mathbf{e}^A) = (\varepsilon^1, \dots, \varepsilon^n, e^1, \dots, e^\nu)$. Here, the dependence of the parity of a supervector on the parity of its components is evidently inverted.

Two supervectors $(\mathbf{e}_1^A), (\mathbf{e}_2^A)$ are called *linearly independent*, if and only if

$$\alpha \mathbf{e}_1^A + \beta \mathbf{e}_2^A = 0 \Rightarrow \alpha = \beta = 0, \quad (\text{A.4})$$

where $\alpha, \beta \in \Lambda_N$. This is equivalent to their bodies $\mathbf{e}_1^\circ, \mathbf{e}_2^\circ$ being linearly independent. A *scalar product* between two complex supervectors is supposed to be graded antilinear, i.e.

$$(\mathbf{a}, \mathbf{b}) = (-1)^{\tilde{\mathbf{a}}\tilde{\mathbf{b}}} (\mathbf{b}, \mathbf{a})^*. \quad (\text{A.5})$$

For even supervectors with components $\mathbf{a} = (a^i, \eta^\alpha)$ and $\mathbf{b} = (b^i, \zeta^\alpha)$, where $a^i, b^i \in \Lambda_{N,c}$ and $\eta^\alpha, \zeta^\alpha \in \Lambda_{N,a}$, we can thus define

$$(\mathbf{a}, \mathbf{b}) := \mathbf{a}^\dagger \mathbf{b} := \bar{a}^i b^i + i \bar{\eta}^\alpha \zeta^\alpha. \quad (\text{A.6})$$

Two supervectors are perpendicular, if they have non-vanishing bodies and the scalar product between them vanishes. It follows that they are linearly independent.

A general supermatrix acting on the elements of an $n|\nu$ -dimensional supervector space is of the block form

$$\mathbf{M} = \left(\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right), \quad (\text{A.7})$$

where A is of dimension $n \times n$ and D of dimension $\nu \times \nu$. The supermatrix \mathbf{M} is called even, if A and D have only even components and B and C consist only of odd components; it thus preserves the parity of any supervector it acts on. Furthermore, if it inverts the parity of the supervector, it is called odd.

Note that the space $\mathbb{C}^{n|\nu}$ is defined in two different ways throughout the literature. Most commonly, it denotes a space described by a set of coordinates consisting of n even and complex numbers and ν complex Grassmann variables. On the other hand, it is a $n|\nu$ -dimensional supervector space over a complex supercommutative ring, as e.g. the ring of complex super-numbers. For the description of flag supermanifolds, we need the latter definition.

The supergroup $U(4|4)$

In our conventions for supergroups, we follow essentially [38], see also [43]. The Lie superalgebra $\mathfrak{u}(n|\nu)$ is given by block supermatrices of the form (A.7), where A and D are elements of $\mathfrak{u}(n)$ and $\mathfrak{u}(\nu)$, respectively, while B and C are hermitian conjugates of each other. This algebra is generated by $n^2 + \nu^2$ even generators with non-vanishing entries only in A and C , as well as $2n\nu$ odd generators with entries in B and C . Exponentiating these generators with even and odd parameters, respectively, yields the supergroup $U(n|\nu)$. To obtain the superanalogue of $\mathfrak{su}(n|\nu)$, one linearly combines the identities λ_n^0 and λ_ν^0 of $\mathfrak{u}(n)$ and $\mathfrak{u}(\nu)$ into

$$\left(\begin{array}{cc|c} \frac{1}{\sqrt{n}} \mathbb{1}_n & 0 & \\ \hline 0 & \frac{1}{\sqrt{\nu}} \mathbb{1}_\nu & \end{array} \right) \quad \text{and} \quad \left(\begin{array}{cc|c} \frac{1}{\sqrt{n}} \mathbb{1}_n & 0 & \\ \hline 0 & -\frac{1}{\sqrt{\nu}} \mathbb{1}_\nu & \end{array} \right). \quad (\text{A.8})$$

Imposing the condition $\text{str}(\cdot) = 0$ on the generators, eliminates the second generator. For $n \neq \nu$, this yields a semisimple super Lie algebra $\mathfrak{su}(n|\nu)$. For $n = \nu$, however, the first factor becomes $\frac{1}{n} \mathbb{1}_{2n}$ and generates an invariant Abelian subgroup. For this reason, one excludes this generator and arrives at $\mathfrak{psu}(n|n)$. However, the lowest dimensional representation is the adjoint, see e.g. [38, 41] for more details. To avoid these complications, we choose to work with $\mathfrak{u}(4|4)$.

The Killing form for $\mathfrak{u}(n|n)$ is easily evaluated to be

$$g_{\hat{A}\hat{B}} := \text{str}(\boldsymbol{\lambda}_{\hat{A}} \boldsymbol{\lambda}_{\hat{B}}) = \left(\begin{array}{cc|ccc} \mathbb{1}_n & 0 & & & \\ \hline 0 & \mathbb{1}_\nu & & & \\ \hline 0 & & -\sigma_2 & 0 & 0 \\ & & 0 & -\sigma_2 & 0 \\ & & 0 & 0 & \ddots \end{array} \right)_{\hat{A}\hat{B}}, \quad (\text{A.9})$$

and we also define $g^{\hat{A}\hat{B}}$ with $g_{\hat{A}\hat{B}} g^{\hat{B}\hat{C}} = \delta_{\hat{A}}^{\hat{C}}$. The Killing form is furthermore supersymmetric, i.e. $g_{\hat{A}\hat{B}} = (-1)^{\hat{A}\hat{B}} g_{\hat{B}\hat{A}}$ and non-degenerate.

Supermanifolds

We define a (complex) supermanifold [44] as a topological space X together with a sheaf \mathcal{O}_N of \mathbb{Z}_2 -graded supercommutative rings on X , satisfying the following two conditions

- (i) There is a projection on the “body” of X , which is an ordinary complex manifold of dimension m . More explicitly, consider the reduced structure sheaf $\mathcal{O}^\circ := \mathcal{O}_N/\mathcal{I}$, where \mathcal{I} is the ideal of nilpotent elements in \mathcal{O}_N . We demand that (X, \mathcal{O}°) is a complex manifold of dimension m .
- (ii) Locally, the structure sheaf is the structure sheaf of the body with values in a Grassmann algebra. That is, for every point x in X , there is an open neighborhood U such that

$$\mathcal{O}_N|_U \cong \mathcal{O}_{\text{red}}|_U(\Lambda^*\mathbb{C}^m). \quad (\text{A.10})$$

We will define the dimension of such a supermanifold to be $m|n$. For more details on supermanifolds, see [33, 41, 35].

Riemannian supergeometry

A *supermetric* on a (real) supermanifold $\mathbf{M} := (M, \mathcal{O}_N)$ is an \mathcal{O}_N -linear, even map $g : T\mathbf{M} \otimes_{\mathcal{O}_N} T\mathbf{M}$ satisfying the following properties:

- (i) g is supersymmetric: $g(X \otimes Y) = (-1)^{\tilde{X}\tilde{Y}} g(Y \otimes X)$.
- (ii) g induces a Riemannian metric on (M, \mathcal{O}°) .
- (iii) g induces a symplectic form on the fermionic tangent directions of $T\mathbf{M}$.

An *almost complex structure* on a (real) supermanifold $\mathbf{M} := (M, \mathcal{O}_N)$ is an even, smooth map $I : T\mathbf{M} \rightarrow T\mathbf{M}$, which satisfies $I^2 = -\mathbb{1}$. As in ordinary complex geometry, a real supermanifold underlying a complex supermanifold has a natural almost complex structure.

A *hermitian supermetric* on a supermanifold $\mathbf{M} := (M, \mathcal{O}_N)$ with almost complex structure I is a supermetric g which satisfies $g(IX \otimes IY) = g(X \otimes Y)$ for all vector fields X, Y in $T\mathbf{M}$. A *Kähler supermetric* is a supermetric, the derived *Kähler form* $J(X, Y) = g(X, IY)$ of which is closed: $dJ = 0$. As an example of a Kähler supermanifold see the discussion of the space $\mathbb{C}P^{3|4}$ in section 5.5. For more details on supergeometry, see e.g. [45].

B. Representations of $\mathfrak{su}(4)$ and $\mathfrak{u}(4|4)$

In this appendix, we briefly recall a few facts on the representation theory of $\mathfrak{su}(4)$ and give the necessary background on the superalgebra $\mathfrak{u}(4|4)$ and its subalgebra $\mathfrak{su}(4)$.

Representation of $\mathfrak{su}(4)$ in terms of Plücker and embedding coordinates

Consider the case $\mathbb{C}P^3 \cong \text{SU}(4)/\text{U}(3) \subset \mathbb{R}^{16}$. A representation of $\text{SU}(4)$ acting on functions written in terms of complex Plücker coordinates $a^i \in \mathbb{C}^4$ is given by

$$\mathcal{L}^{\hat{a}} = \bar{a}^i \lambda_{ij}^{\hat{a}} \frac{\partial}{\partial \bar{a}^j} - a^j \lambda_{ij}^{\hat{a}} \frac{\partial}{\partial a^i}, \quad (\text{B.1})$$

Schwinger construction for Lie superalgebras

Consider a Lie superalgebra having the even generators $\boldsymbol{\lambda}^a$ and the odd generators $\boldsymbol{\lambda}^\alpha$ satisfying the commutation relations

$$[\boldsymbol{\lambda}^a, \boldsymbol{\lambda}^b] = f^{ab}{}_c \boldsymbol{\lambda}^c, \quad [\boldsymbol{\lambda}^a, \boldsymbol{\lambda}^\alpha] = f^{a\alpha}{}_\beta \boldsymbol{\lambda}^\beta, \quad \{\boldsymbol{\lambda}^\alpha, \boldsymbol{\lambda}^\beta\} = f^{\alpha\beta}{}_a \boldsymbol{\lambda}^a. \quad (\text{B.11})$$

We summarize the generators into $\boldsymbol{\lambda}^A = (\boldsymbol{\lambda}^a, \boldsymbol{\lambda}^\alpha)$ and the commutation relations to

$$\{\{\boldsymbol{\lambda}^A, \boldsymbol{\lambda}^B\}\} = f^{AB}{}_C \boldsymbol{\lambda}^C, \quad (\text{B.12})$$

where $\{\cdot\}$ denotes the supercommutator. Assume furthermore that the generators are in a representation acting on an $(m|n)$ -dimensional supervector space. After introducing a set of m bosonic and n fermionic oscillators together with the corresponding annihilation and creation operators

$$\hat{\mathbf{a}}^I = (\hat{a}^i, \hat{\eta}^\alpha) \quad \text{and} \quad \hat{\mathbf{a}}^{I\dagger} = (\hat{a}^{i\dagger}, \hat{\eta}^{\alpha\dagger}), \quad (\text{B.13})$$

the Schwinger construction yields a representation of the Lie superalgebra by

$$\hat{\mathbf{L}}^A = \hat{\mathbf{a}}^{I\dagger} \boldsymbol{\lambda}_{IJ}^A \hat{\mathbf{a}}^J. \quad (\text{B.14})$$

Representation of $\mathfrak{u}(4|4)$ in terms of Plücker and embedding coordinates

Completely analogously to the representations of $\mathfrak{u}(4)$ in terms of coordinates describing $\mathbb{C}P^3$, one finds a representation of $\mathfrak{u}(4|4)$ in terms of coordinates describing $\mathbb{C}P^{3|4}$. We have

$$\mathcal{L}^{\hat{A}} = \bar{\mathbf{a}}^I \boldsymbol{\lambda}_{IJ}^{\hat{A}} \frac{\partial}{\partial \bar{\mathbf{a}}^J} - \mathbf{a}^J \boldsymbol{\lambda}_{IJ}^{\hat{A}} \frac{\partial}{\partial \mathbf{a}^I}, \quad (\text{B.15})$$

where $\mathbf{a}^I = (a^i, \eta^\alpha) \in \mathbb{C}^{4|4}$ and in terms of the real coordinates

$$\mathbf{x}^{\hat{A}} = \bar{\mathbf{a}}^I \boldsymbol{\lambda}_{IJ}^{\hat{A}} \mathbf{a}^J \quad (\text{B.16})$$

describing the embedding of $\mathbb{C}P^{3|4}$ in $\mathbb{R}^{32|32}$, the above generators read as

$$\mathcal{L}^A = -i\sqrt{2} f^{ABC} \mathbf{x}_B \frac{\partial}{\partial \mathbf{x}^C}, \quad \mathcal{L}^0 \sim \mathbf{x}^A \frac{\partial}{\partial \mathbf{x}^A}. \quad (\text{B.17})$$

Representations and Dynkin diagrams for $\mathfrak{psu}(4|4)$

The representations of supergroups are divided into two classes. In the first class, the representation space is spanned by an even basis, while in the second one, the basis is odd and they can be regarded as dual to each other. Representations are again labelled by highest weights, which are the eigenvalues of the generators of the Cartan subalgebra. The latter is generated by two copies of the Cartan subalgebra of $\mathfrak{su}(4)$ as well as $H_4 := \text{diag}(0, 0, 0, 1, 1, 0, 0, 0)$, cf. section 6.1. Accordingly, the Dynkin diagram has seven nodes:

$$\begin{array}{ccccccc} a_1 & & a_2 & & a_3 & & q_0 & & a_4 & & a_5 & & a_6 \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \otimes & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \end{array} \quad (\text{B.18})$$

It is now evident that a class I representation is dual to a class II representation provided that

$$a_i^I = a_{7-i}^{II} \quad \text{and} \quad q_0^I = q_0^{II}. \quad (\text{B.19})$$

Since the superdeterminant is no longer a polynomial of finite degree, there is no invariant totally antisymmetric ε -tensor for the $SU(m|n)$ supergroups. However, the duality between class I and II representation takes over the rôle of the duality between the representation and the corresponding conjugate representation. The latter arises from the interchange between covariant and contravariant indices corresponding to a contraction with an ε -tensor in the $SU(4)$ picture, as we saw above.

The second order Casimir operator on irreducible representations of $\mathfrak{u}(4)$

We constructed the algebra of functions on the fuzzy flag manifolds from spherical representations of the groups underlying the flag manifolds. These representations are described by Young diagrams, and given such a Young diagram with $m \leq 4$ rows of $n_1 \geq \dots \geq n_m \geq 0$ boxes, the eigenvalue of the second order Casimir operator on these representations reads as¹³

$$\frac{1}{2} \left(4 \sum_{i=1}^m n_i + \sum_{i=1}^m n_i(n_i + 1 - 2i) - \frac{(\sum_{i=1}^m n_i)^2}{4} \right). \quad (\text{B.20})$$

We will be mostly interested in the representations consisting of a row of $2L$ boxes and 2 rows of L boxes. For these diagrams, the above formula reduces to $L(L + 3)$. The other type of diagrams we will encounter consist of three rows with $a + b + a$, $a + b$ and a boxes, respectively. For them, the above formula reduces to $a^2 + a(3 + b) + \frac{1}{2}b(4 + b)$.

The second order Casimir operator on representations of $\mathfrak{u}(n|n)$

We can write the second order Casimir operator acting on the Hilbert space $\mathcal{H}_{1|0;4|4}^F$ of $\mathbb{C}P_F^{3|4}$ in terms of one set of oscillators using the Schwinger construction:

$$\begin{aligned} C_2 &= g_{AB} \hat{\mathbf{L}}^A \hat{\mathbf{L}}^B \\ &= \frac{n-1}{n} \hat{\mathbf{N}}_b (\hat{\mathbf{N}}_b + n) + \frac{1}{n} \hat{\mathbf{N}}_b^2 - \frac{n+1}{n} \hat{\mathbf{N}}_f (n - \hat{\mathbf{N}}_f) - \frac{1}{n} \hat{\mathbf{N}}_f^2 + (2\hat{\mathbf{N}}_b \hat{\mathbf{N}}_f + n\hat{\mathbf{N}}_f - n\hat{\mathbf{N}}_b) \\ &= \frac{1}{2} \hat{\mathbf{N}} (\hat{\mathbf{N}} - 1), \end{aligned}$$

where $\hat{\mathbf{N}}_b = \hat{a}^{i\dagger} \hat{a}^i$ and $\hat{\mathbf{N}}_f = \hat{\eta}^{\alpha\dagger} \hat{\eta}^\alpha$. For the other flag supermanifold we require more than one set of oscillators and the calculation is more complicated.

Fierz and super Fierz identities

Consider the Gell-Mann matrices λ^a , $a = 1, \dots, 15$ of $\mathfrak{su}(4)$ and extend them to the generators $\lambda^{\hat{a}}$ of $\mathfrak{u}(4)$ by adding $\lambda^0 = \mathbb{1}/\sqrt{4}$. We have the Fierz identity

$$\lambda_{ij}^{\hat{a}} \lambda_{kl}^{\hat{a}} = \delta_{il} \delta_{jk}, \quad (\text{B.21})$$

which trivially extends in the case of the antisymmetric tensor products $\lambda^{\hat{a}\hat{b}\dots} := \lambda^{\hat{a}} \wedge \lambda^{\hat{b}} \wedge \dots$ we defined in section 2.3 to

$$\begin{aligned} (\lambda^{\hat{a}\hat{b}})_{ij;kl} (\lambda^{\hat{a}\hat{b}})_{mn;pq} &= (\delta_{ip} \delta_{km} \delta_{jq} \delta_{ln})_{[ij][kl][mn][pq]}, \\ (\lambda^{\hat{a}\hat{b}\hat{c}})_{ijk;lmn} (\lambda^{\hat{a}\hat{b}\hat{c}})_{pqr;stu} &= (\delta_{is} \delta_{lp} \delta_{jt} \delta_{mq} \delta_{ku} \delta_{nr})_{[ijk][lmn][pqr][stu]}. \end{aligned} \quad (\text{B.22})$$

¹³This formula is given, e.g., in [46] with a different normalization of the Lie algebra generators.

Moreover, we can write

$$\mathrm{tr}(A_1 \lambda^{\hat{a}}) \lambda^{\hat{a}} = A_1 \quad \text{and} \quad \mathrm{tr}(A_1 \lambda^a) \lambda^a = A_1 - \frac{1}{4} \mathrm{tr}(A_1) \mathbb{1} \quad (\text{B.23a})$$

for any hermitian matrix $A_1 \in \mathrm{Mat}_4$ as well as

$$\begin{aligned} \mathrm{tr}(A_2 \lambda^{\hat{a}\hat{b}}) \lambda^{\hat{a}\hat{b}} &= A_2 \\ \mathrm{tr}(A_3 \lambda^{\hat{a}\hat{b}\hat{c}}) \lambda^{\hat{a}\hat{b}\hat{c}} &= A_3 \end{aligned} \quad (\text{B.23b})$$

for hermitian matrices $A_2 \in \mathrm{Mat}_4 \wedge \mathrm{Mat}_4$ and $A_3 \in \mathrm{Mat}_4 \wedge \mathrm{Mat}_4 \wedge \mathrm{Mat}_4$.

We can use the Killing form on $U(4|4)$ to establish the following super Fierz identity:

$$g^{\hat{A}\hat{B}} \lambda_{\hat{A}}^{IJ} \lambda_{\hat{B}}^{KL} = \delta^{IL} \delta^{JK} . \quad (\text{B.24})$$

To prove this formula quickly, one can e.g. extend the one for $\mathfrak{su}(n|n)$ given in [38]. From this Fierz identity, we can, as in the purely bosonic case, immediately derive the further identities

$$\begin{aligned} (\lambda^{\hat{A}\hat{B}})_{IJ;KL} (\lambda^{\hat{A}\hat{B}})_{MN;PQ} &= (\delta_{IP} \delta_{KM} \delta_{JQ} \delta_{LN}) \{IJ\} \{KL\} \{MN\} \{PQ\} , \\ (\lambda^{\hat{A}\hat{B}\hat{C}})_{IJK;LMN} (\lambda^{\hat{A}\hat{B}\hat{C}})_{PQR;STU} &= (\delta_{IS} \delta_{LP} \delta_{JT} \delta_{MQ} \delta_{KU} \delta_{NR}) \{IJK\} \{LMN\} \{PQR\} \{STU\} , \end{aligned}$$

where $\lambda^{\hat{A}\hat{B}}$ denotes again the graded antisymmetric tensor product

$$\lambda^{\hat{A}\hat{B}} = \lambda^{\hat{A}} \hat{\cap} \lambda^{\hat{B}} \quad (\text{B.25})$$

introduced in section 5.3.

Oscillator representation of internal isotropy subgroups

In the description of the various Graßmannians, we needed the following formula to describe the action of the internal part of the isotropy subgroup acting on the states of certain Fock spaces:

$$[(a_p^i)^\dagger \lambda_{pq}^A a_q^i, a_1^{[j_1 \dagger} \dots a_k^{j_k] \dagger}] = 0 . \quad (\text{B.26})$$

To prove this formula, consider the following form of the generators L_{pq} of $U(k)$:

$$L_{pq} = a_p^{i \dagger} a_q^i , \quad (\text{B.27})$$

where i is summed over $1, \dots, n$. The composite creation operators relevant in the description of $G_{k;n}$ can be written as

$$A_k^{i_1 \dots i_k \dagger} = \frac{1}{k!} \epsilon_{p_1 \dots p_k} a_{p_1}^{[i_1 \dagger} \dots a_{p_k}^{i_k] \dagger} . \quad (\text{B.28})$$

Then

$$\begin{aligned} [L_{pq}, A_k^{i_1 \dots i_k \dagger}] &= \frac{1}{k!} \epsilon_{p_1 \dots p_k} a_p^{i \dagger} [a_q^i, a_{p_1}^{[i_1 \dagger} \dots a_{p_k}^{i_k] \dagger}] \\ &= \frac{1}{k!} \sum_{j=1}^k \epsilon_{p_1 \dots p_k} a_p^{i \dagger} a_{p_1}^{[i_1 \dagger} \dots \delta^{i_j} \delta_{qp_j} \dots a_{p_k}^{i_k] \dagger} \\ &= \frac{k}{k!} \epsilon_{qp_2 \dots p_k} a_p^{[i_1 \dagger} a_{p_2}^{i_2 \dagger} \dots a_{p_k}^{i_k] \dagger} \\ &= \delta_{pq} A_k^{i_1 \dots i_k \dagger} , \end{aligned}$$

and thus the (traceless) generators of $SU(k)$ leave the composite creation operators invariant.

In the supersymmetric case, the same statement holds. Here, the creation and annihilation operators satisfy

$$\{\mathbf{a}_p^I, \mathbf{a}_q^{J\dagger}\} = \delta_{pq} \delta^{IJ} . \quad (\text{B.29})$$

Using the relations

$$\begin{aligned} \{\mathbf{a}, \mathbf{bc}\} &= \{\mathbf{a}, \mathbf{b}\} \mathbf{c} + (-1)^{\tilde{\mathbf{a}}\tilde{\mathbf{b}}} \mathbf{b} \{\mathbf{a}, \mathbf{c}\} , \\ \{\mathbf{ab}, \mathbf{c}\} &= \mathbf{a} \{\mathbf{bc}\} + (-1)^{\tilde{\mathbf{b}}\tilde{\mathbf{c}}} \{\mathbf{a}, \mathbf{c}\} \mathbf{b} , \end{aligned} \quad (\text{B.30})$$

we obtain

$$\begin{aligned} \{\mathbf{a}_p^I, \mathbf{A}_{2|0}^{JK\dagger}\} &= \{\mathbf{a}_p^I, \mathbf{a}_1^{J\dagger}\} \mathbf{a}_2^{K\dagger} + (-1)^{\tilde{I}\tilde{J}} \mathbf{a}_1^{J\dagger} \{\mathbf{a}_p^I, \mathbf{a}_2^{K\dagger}\} - (-1)^{\tilde{J}\tilde{K}} (J \leftrightarrow K) \\ &= \delta_{p1} \delta^{IJ} \mathbf{a}_2^{K\dagger} + \delta_{p2} (-1)^{\tilde{I}\tilde{J}} \delta^{IK} \mathbf{a}_1^{J\dagger} - (-1)^{\tilde{J}\tilde{K}} (J \leftrightarrow K) . \end{aligned} \quad (\text{B.31})$$

With this relation, we can conclude that

$$\begin{aligned} \{\mathbf{a}_p^{I\dagger} \mathbf{a}_q^I, \mathbf{A}_{2|0}^{JK\dagger}\} &= \mathbf{a}_p^{I\dagger} \{\mathbf{a}_q^I, \mathbf{A}_{2|0}^{JK\dagger}\} \\ &= \delta_{q1} (\mathbf{a}_p^{J\dagger} \mathbf{a}_2^{K\dagger} - (-1)^{\tilde{J}\tilde{K}} \mathbf{a}_p^{K\dagger} \mathbf{a}_2^{J\dagger}) + \delta_{q2} ((-1)^{\tilde{K}\tilde{J}} \mathbf{a}_p^{K\dagger} \mathbf{a}_1^{J\dagger} - \mathbf{a}_p^{J\dagger} \mathbf{a}_1^{K\dagger}) \\ &= \delta_{p1} \delta_{q1} \mathbf{A}_{2|0}^{JK\dagger} + \delta_{p2} \delta_{q2} \mathbf{A}_{2|0}^{JK\dagger} \\ &= \delta_{pq} \mathbf{A}_{2|0}^{JK\dagger} . \end{aligned} \quad (\text{B.32})$$

Similar relations can also be proven for the remaining cases of $\mathbf{A}_{2|4}$ and $\mathbf{A}_{3|4}$ by observing that, for example,

$$\mathbf{A}_{3|0}^{JKL\dagger} = \mathbf{A}_{2|0}^{JK\dagger} \mathbf{a}_3^{L\dagger} - (-1)^{\tilde{K}\tilde{L}} \mathbf{A}_{2|0}^{JL\dagger} \mathbf{a}_3^{K\dagger} + (-1)^{\tilde{J}\tilde{K} + \tilde{J}\tilde{L}} \mathbf{A}_{2|0}^{KL\dagger} \mathbf{a}_3^{J\dagger} \quad (\text{B.33})$$

and

$$\{\mathbf{a}_p^{I\dagger} \mathbf{a}_q^I, \mathbf{A}_{2|0}^{JK\dagger} \mathbf{a}_3^{L\dagger}\} = \{\mathbf{a}_p^{I\dagger} \mathbf{a}_q^I, \mathbf{A}_{2|0}^{JK\dagger}\} \mathbf{a}_3^{L\dagger} + \mathbf{A}_{2|0}^{JK\dagger} \{\mathbf{a}_p^{I\dagger} \mathbf{a}_q^I, \mathbf{a}_3^{L\dagger}\} \quad (\text{B.34})$$

$$= (\delta_{pq} - \delta_{p3} \delta_{q3}) \mathbf{A}_{2|0}^{JK\dagger} \mathbf{a}_3^{L\dagger} + \delta_{q3} \mathbf{A}_{2|0}^{JK\dagger} \mathbf{a}_p^{L\dagger} . \quad (\text{B.35})$$

Hence,

$$\{\mathbf{a}_p^{I\dagger} \mathbf{a}_q^I, \mathbf{A}_{3|0}^{JKL\dagger}\} = \delta_{pq} \mathbf{A}_{3|0}^{JKL\dagger} . \quad (\text{B.36})$$

References

- [1] J. Madore, *The fuzzy sphere*, *Class. Quant. Grav.* **9** (1992) 69;
- [2] J. Hoppe, *Quantum theory of a massless relativistic surface and a two-dimensional bound state problem*, PhD Thesis, MIT, 1982, <http://www.aei.mpg.de/~hoppe/>; A. P. Balachandran, *Quantum spacetimes in the year 1*, *Pramana* **59** (2002) 359 [hep-th/0203259].
- [3] H. Grosse, C. Klimcik, and P. Presnajder, *Towards finite quantum field theory in non-commutative geometry*, *Int. J. Theor. Phys.* **35** (1996) 231 [hep-th/9505175]; H. Grosse and A. Strohmaier, *Noncommutative geometry and the regularization problem of 4D quantum field theory*, *Lett. Math. Phys.* **48** (1999) 163 [hep-th/9902138]; B. P. Dolan, D. O'Connor, and P. Presnajder, *Matrix ϕ^4 models on the fuzzy sphere and their continuum limits*, *JHEP* **03** (2002) 013 [hep-th/0109084].

- [4] S. Baez, A. P. Balachandran, B. Ydri, and S. Vaidya, *Monopoles and solitons in fuzzy physics*, Commun. Math. Phys. **208** (2000) 787 [hep-th/9811169]; A. P. Balachandran and S. Vaidya, *Instantons and chiral anomaly in fuzzy physics*, Int. J. Mod. Phys. A **16** (2001) 17 [hep-th/9910129]; G. Alexanian, A. P. Balachandran, G. Immirzi, and B. Ydri, *Fuzzy CP^2* , J. Geom. Phys. **42** (2002) 28 [hep-th/0103023]; B. P. Dolan and C. Nash, *Chiral fermions and Spin(C) structures on matrix approximations to manifolds*, JHEP **07** (2002) 057 [hep-th/0207007].
- [5] A. P. Balachandran, X. Martin, and D. O'Connor, *Fuzzy actions and their continuum limits*, Int. J. Mod. Phys. A **16** (2001) 2577 [hep-th/0007030].
- [6] A. P. Balachandran, B. P. Dolan, J.-H. Lee, X. Martin, and D. O'Connor, *Fuzzy complex projective spaces and their star-products*, J. Geom. Phys. **43** (2002) 184 [hep-th/0107099].
- [7] B. Ydri, *Fuzzy physics*, hep-th/0110006.
- [8] B. P. Dolan and O. Jahn, *Fuzzy complex Grassmannian spaces and their star products*, Int. J. Mod. Phys. A **18** (2003) 1935 [hep-th/0111020].
- [9] S. Vaidya, *Perturbative dynamics on fuzzy S^2 and RP^2* , Phys. Lett. B **512** (2001) 403 [hep-th/0102212]; H. Grosse, C. Klimcik, and P. Presnajder, *Finite quantum field theory in noncommutative geometry*, Commun. Math. Phys. **180** (1996) 429 [hep-th/9602115].
- [10] S. Ramgoolam, *On spherical harmonics for fuzzy spheres in diverse dimensions*, Nucl. Phys. B **610** (2001) 461 [hep-th/0105006]; Y. Kimura, *Noncommutative gauge theory on fuzzy four-sphere and matrix model*, Nucl. Phys. B **637** (2002) 177 [hep-th/0204256]; B. P. Dolan and D. O'Connor, *A fuzzy three sphere and fuzzy tori*, JHEP **10** (2003) 060 [hep-th/0306231]; B. P. Dolan, D. O'Connor, and P. Presnajder, *Fuzzy complex quadrics and spheres*, JHEP **02** (2004) 055 [hep-th/0312190].
- [11] A. P. Balachandran, S. Kurkcuglu, and S. Vaidya, *Lectures on fuzzy and fuzzy SUSY physics*, hep-th/0511114.
- [12] C. Klimcik and P. Severa, *Open strings and D-branes in WZNW models*, Nucl. Phys. B **488** (1997) 653 [hep-th/9609112]; H. Garcia-Compean and J. F. Plebanski, *D-branes on group manifolds and deformation quantization*, Nucl. Phys. B **618** (2001) 81 [hep-th/9907183].
- [13] A. Y. Alekseev, A. Recknagel, and V. Schomerus, *Non-commutative world-volume geometries: Branes on $SU(2)$ and fuzzy spheres*, JHEP **09** (1999) 023 [hep-th/9908040].
- [14] R. C. Myers, *Dielectric-branes*, JHEP **12** (1999) 022 [hep-th/9910053].
- [15] S. P. Trivedi and S. Vaidya, *Fuzzy cosets and their gravity duals*, JHEP **09** (2000) 041 [hep-th/0007011]; S. R. Das, S. P. Trivedi, and S. Vaidya, *Magnetic moments of branes and giant gravitons*, JHEP **10** (2000) 037 [hep-th/0008203].

- [16] A. F. Kleppe and C. Wainwright, *Super coset space geometry*, hep-th/0610039.
- [17] G. Gotz, T. Quella, and V. Schomerus, *The WZNW model on PSU(1,1|2)*, hep-th/0610070.
- [18] P. S. Howe and G. G. Hartwell, *A superspace survey*, Class. Quant. Grav. **12** (1995) 1823.
- [19] E. Ivanov, L. Mezincescu and P. K. Townsend, *Fuzzy CP^(n|m) as a quantum superspace*, hep-th/0311159; E. Ivanov, L. Mezincescu and P. K. Townsend, *A super-flag Landau model*, hep-th/0404108; E. Ivanov, L. Mezincescu and P. K. Townsend, *Planar super-Landau models*, JHEP **0601** (2006) 143 [hep-th/0510019].
- [20] A. D. Popov and C. Saemann, *On supertwistors, the Penrose-Ward transform and N = 4 super Yang-Mills theory*, Adv. Theor. Math. Phys. **9** (2005) 931 [hep-th/0405123].
- [21] B. P. Dolan, I. Huet, S. Murray, and D. O'Connor, *Noncommutative vector bundles over fuzzy CP^N and their covariant derivatives*, hep-th/0611209.
- [22] M. Aganagic and C. Vafa, *Mirror symmetry and supermanifolds*, hep-th/0403192; S. P. Kumar and G. Policastro, *Strings in twistor superspace and mirror symmetry*, Phys. Lett. B **619** (2005) 163 [hep-th/0405236].
- [23] R. F. Picken, *The Duistermaat-Heckman integration formula on flag manifolds*, J. Math. Phys. **31** (1990) 616.
- [24] L. J. Mason and N. M. J. Woodhouse, *Integrability, self-duality, and twistor theory*, Clarendon, Oxford (1996).
- [25] A. Cucchieri, T. Mendes, and A. Pelissetto, *Pseudo-character expansions for U(N)-invariant spin models on CP^{N-1}*, Ann. Comb. **3** (1999) 287 [hep-lat/9810027].
- [26] P. B. Gilkey, J. V. Leahy, and J. Park, *Spinors, spectral geometry and Riemannian submersions*, Global Analysis Research Center of Seoul National University (1998).
- [27] S. Helgason, *Groups and geometric analysis*, Acad. Press (1984).
- [28] M. S. Dijkhuizen, *Some remarks on the construction of quantum symmetric spaces*, math.QA/9512225.
- [29] A. M. Perelomov, *Generalized coherent states and their applications*, Springer, Berlin (1986).
- [30] M. Chaichian, D. Ellinas, and P. Presnajder, *Path integrals and supercoherent states*, J. Math. Phys. **32** (1991) 3381. H. Grosse and P. Presnajder, *The construction on noncommutative manifolds using coherent states*, Lett. Math. Phys. **28** (1993) 239.

- [31] A. M. Perelomov, *Description of generalized coherent states closest to classical states (in Russian)*, Yad. Fiz. **29** (1979) 1688.
- [32] S. Kurkcuoğlu and C. Saemann, *Drinfeld twist and general relativity with fuzzy spaces*, hep-th/0606197.
- [33] Y. I. Manin, *Gauge field theory and complex geometry*, Grundlehren der mathematischen Wissenschaften, 289, Springer (1988).
- [34] S. Kleiman and D. Laksov, *Schubert Calculus*, Amer. Math. Monthly **79** (1972) 1061.
- [35] P. Cartier, C. DeWitt-Morette, M. Ihl, C. Saemann, and M. E. Bell, *Supermanifolds - Application To Supersymmetry*, math-ph/0202026. in: “Multiple facets of quantization and supersymmetry: Michael Marinov memorial volume”, Eds. M. Olshanetsky and A. Vainshtein, World Scientific (2002).
- [36] C. Klimcik, *An extended fuzzy supersphere and twisted chiral superfields*, Commun. Math. Phys. **206** (1999) 587 [hep-th/9903202].
- [37] I. Bars, B. Morel, and H. Ruegg, *Kac-Dynkin diagrams and supertableaux*, J. Math. Phys. **24** (1983) 2253.
- [38] I. Bars, *Supergroups and their representations*, Lectures Appl. Math. **21** (1983) 17.
- [39] E. Witten, *Perturbative gauge theory as a string theory in twistor space*, Commun. Math. Phys. **252** (2004) 189 [hep-th/0312171].
- [40] M. Aganagic and C. Vafa, *Mirror symmetry, D-branes and counting holomorphic discs*, hep-th/0012041.
- [41] B. S. DeWitt, *Supermanifolds*, 2nd ed., Cambridge University Press, Cambridge (1992).
- [42] P. Deligne, P. Etingof, D. Freed, L. Jeffrey, D. Kazhdan, J. Morgan, D. Morrison, and E. Witten, *Quantum fields and strings: A course for mathematicians*, AMS, Providence (1999).
- [43] V. G. Kac, *Representations of classical Lie superalgebras*, in: “Differential Geometrical Methods in Mathematical Physics II”, Lectures Notes in Mathematics 676, Springer (1978).
- [44] D. Leites, *Spectra of graded commutative rings*, Russian Math. Surveys **30** (1974) 209.
- [45] S. Varsaie, *Kählerian supermanifolds*, J. Math. Phys. **40** (1999) 6701.
- [46] D. J. Gross, *Two-dimensional QCD as a string theory*, Nucl. Phys. B **400** (1993) 161 [hep-th/9212149].