

Title	Equality of NMR Relaxation Times for Different Molecular Models
Creators	McConnell, James
Date	1987
Citation	McConnell, James (1987) Equality of NMR Relaxation Times for Different Molecular Models. (Preprint)
URL	https://dair.dias.ie/id/eprint/816/
DOI	DIAS-STP-87-19

EQUALITY OF NMR RELAXATION TIMES FOR DIFFERENT MOLECULAR MODELS
James McConnell, Dublin Institute for Advanced Studies, Dublin 4

Expressions for nuclear magnetic relaxation times and for dielectric permittivity derived for spherical molecules are applicable in rotational diffusion theory to linear and symmetric top molecules. This holds for nuclear magnetic relaxation by intramolecular dipolar interaction, quadrupolar interaction or anisotropic chemical shift, provided that in the latter two cases the interactions have axes of cylindrical symmetry which are parallel to the axes of rotational symmetry of the molecule.

The theorem is proved by deriving equations for ensemble average of the rotation operator $\langle R(t) \rangle$ associated with the molecule, and then showing that the dielectric permittivity and relaxation times depend only on $\langle R(t) \rangle_{00}$, where the suffixes denote the 00 - matrix element with respect to spherical harmonics as basis.

- 1) J. McConnell, Physica 138A, 367-381 (1986).